Pre-harvest sprouting in wheat: current status and future prospects

Charan Singh, Umesh Ravindra Kamble, Vikas Gupta*, Gyanendra Singh, Sonia Sheoran, Arun Gupta, Bhudeva Singh Tyagi, Pradeep Kumar, Chandra Nath Mishra, Gopalareddy Krishannapa, Santosh Kumar Bishnoi, Amit Kumar Sharma, Satish Kumar and Gyanendra Pratap Singh

ICAR- Indian Institute of Wheat and Barley Research, Karnal-132001, Haryana

Article history:

Received: 30 July, 2021

Revised: 02 Sept., 2021

Accepted: 17 Sept., 2021

Citation:

Singh C, UR Kamble, V Gupta, G Singh, S Sheoran, A Gupta, BS Tyagi, P Kumar, CN Mishra, K Gopalareddy, SK Bishnoi, AK Sharma, S, Kumar and GP Singh. 2021. Pre-Harvest Sprouting in Wheat: Current Status and Future Prospects. *Journal of Cereal Research* 13 (Spl-1): 1-22. <http://doi.org/10.25174/2582-2675/2021/114484>

*Corresponding author:

E-mail: vikasdwrr@gmail.com

© Society for Advancement of Wheat and Barley Research

Abstract

Pre-harvest sprouting (PHS) having adverse effects on both crop yields and quality, is one of the major constraints of wheat production in areas with high rainfall. In the north-eastern parts of India and also many areas of the world receiving rainfall during the late maturity stages of the crop affects both grain yields and quality. PHS trait is polygenic trait and affected by number of environmental factors. The available diversity in germplasm for the trait is very limited and for improving tolerance to PHS newer sources needs to be identified. The primary cause of pre-harvest sprouting is the breakdown or lacking of seed dormancy under humid and wet conditions along with the degradation of starch in the germinated seeds due to enhanced alpha amylase activity. This review describes the factors affecting the pre-harvest sprouting tolerance, genes associated with it, physiological, biochemical and molecular mechanism involved, screening technique and breeding approaches to develop PHS tolerant wheat genotypes.

Keywords: abscisic acid; dormancy; genes, QTLs; gibberellic acid; pre-harvest sprouting (PHS); wheat

1. Introduction

Wheat (*Triticum aestivum* L.), the widely cultivated prime cereal crop under different agro-climatic conditions throughout the world, provides about 20% of protein and 20% calories to the mankind. The 4th advance estimate of crop season 2020-21, projects 109.52 million tonnes production of wheat in India from an area of 34.24 million hectares (Anonymous, 2021). The advent of climate change and increasing human population pressure over the globe are the most critical challenges for the global food production. This has necessitated the breeders to breed high yielding as well as abiotic and biotic stress tolerant crop varieties for erratic climatic fluctuations to meet the future wheat demand of 140 million tonnes by 2050. In recent years, major wheat growing regions throughout the globe experienced extreme and unpredictable weather

conditions due to climate change. One of the major abiotic factors that limit wheat productivity is the pre-harvest sprouting (PHS); a phenomenon in which seeds germinate while intact on spike or on the mother plant before harvest (Wang *et al.*, 2019) due to rainfall and high humidity (Andreoli *et al.* 2006) at physiological maturity with breakage of dormancy (Patwa and Penning, 2020). PHS sometimes also characterized as *vivipary* (Fang and Chu, 2008), which occurs in a wide range of cereal crops including wheat. The major factors affecting the pre-harvest sprouting are environmental conditions, seed dormancy, seed coat permeability and color, α -amylase activity, endogenous hormones levels, functional proteins, genes, and quantitative trait loci and others (Gao *et al.*, 2013). Pre-harvest sprouting observed across all the major

wheat growing regions in the world (Cabral *et al.*, 2014), especially in the locations witnessing frequent rainfall and high humidity during late season or prior to harvest (Zhang *et al.*, 2017a). The PHS causes reduction in both wheat grain yield as well as end-use quality (Lin *et al.*, 2018). Pre-harvest sprouting have been reported in most of the cereals crop *viz.*, maize, wheat, rice, barley and sorghum (<http://www.fao.org/faostat/en/#compare>, 2019) from most of the regions of the world including Japan, China, India, the United States, Canada, Australia, North Africa, and throughout Europe (Biddulph *et al.*, 2007; Nakamura, 2018). It is reported as a world-wide problem, occurring once in 10 years in major wheat-producing areas throughout the globe (Olaerts *et al.*, 2018). In India, Eastern and far-eastern states namely Bihar, West Bengal, Jharkhand, Assam are more prone to erratic rainfall patterns frequently and aggravated the situation of PHS in wheat.

Wheat is used for the production of various food products *viz.*, breads, cookies, pasta, biscuits, cakes, noodles, and breakfast cereals (Olaerts *et al.*, 2018). However, sprouting of grains reduces the market value and the level of sprouting decides the suitability of the grain for food or feed industry (Cunha *et al.*, 2004). The price of sprouted wheat could be reduced by 20-50% (Simsek *et al.*, 2014; Sorrells and Sherman, 2007), and if having more than 4% sprouted grains treated as unacceptable for producers and consumers (Moot and Every, 1990; Sorenson and Wiersma, 2004). The average annual losses due to pre-harvest sprouting are approximately \$100 million in Canada, while about 24.91 million ha area of wheat is affected in China (Xiao *et al.*, 2002). Annual losses in yield and quality of wheat due to pre-harvest sprouting are around \$1 billion globally (Wahl and Rourke, 1994; Bewley *et al.*, 2006; Biddulph *et al.*, 2008; DePauw *et al.*, 2012; Nakamura, 2018; Shao *et al.*, 2018; Vetch *et al.*, 2019; Liton *et al.*, 2021). The main cause of pre-harvest sprouting is the breakdown or lacking seed dormancy under humid and wet conditions which leads to huge economic losses due to the reduction in grain weight and end-use quality (Zhang and Liu, 1989; Kulwal *et al.*, 2012; Kocheshkova *et al.*, 2017; Shao *et al.*, 2018). The pre-harvest sprouting severely downgrades grain quality due to alpha amylase activity which is mainly degrading starch in wheat grain. Therefore, pre-harvest sprouting is directly associated with reductions in both grain yield as well as quality (Lang *et*

al., 2021). Pre-harvest sprouting causes the reduction in quality of grains due to the degradation of starch and protein in germinated kernels (Flintham, 2000; Shorinola *et al.*, 2016). The increase in amylase activity results in the breakdown of grain carbohydrate reserves due to which bread quality of wheat is affected causing sticky crumb and collapsed loaves (Imtiaz *et al.*, 2008). Alpha-amylase activity has a direct effect on the quality of bread and pasta and adversely impacts the malting process (Pertene, 1964). Enhanced production of alpha-amylase in wheat grain, due to the pre-harvest sprouting, leads to decrease in Hagberg falling number and adverse effects on cooked pasta quality (Singh *et al.* 2014). Further, PHS causes significant yield reduction and decreases milling and baking quality of wheat grains (Groos *et al.*, 2002). The flour from sprouted wheat produces dough that is difficult to handle due to darker crust color, a sticky crumb texture, and poor slice ability (Ibrahim and Appolonia, 1979; Lorenz *et al.*, 1983; Lukow and Bushuk, 1984; Olaerts *et al.*, 2018).

The various products made from sprouted wheat flour are porous, sticky, and off-color and generally of poor bake quality (Sorenson and Wiersma, 2004). Grant *et al.* (1993) reported that sprouting caused higher cooking losses, decreased firmness and lower spaghetti stickiness values. Although, the effects of PHS on final food product depends on the amount of enzymes present and the breakdown of kernel starches, oils, and proteins (Sorenson and Wiersma, 2004), which negatively affect the quality of various products made from wheat flour namely noodles, Arabic flat-breads (Edwards *et al.*, 1989), breads, cookies, pies (Lorenz *et al.*, 1983). Therefore, research needs to be focused on the development of wheat varieties with enhanced degree of dormancy in PHS affected wheat production areas. In this review, the major focus will be on the factors responsible for pre-harvest sprouting, genes associated with PHS tolerance, physiological, biochemical, molecular mechanism and breeding strategies for the improvement of pre-harvest sprouting tolerance in wheat.

2. Factors affecting Pre-harvest sprouting (PHS)

Pre-harvest sprouting is mainly controlled by seed dormancy but also influenced by several other factors such as genetic control, red seed color (Gfeller and Svejda, 1960; Groos *et al.*, 2002), spike morphology

(King and Richard, 1984), physical barriers to water penetration (Gale, 1989) and environmental factors such as temperature and moisture (Argel and Humphreys, 1983; Ceccato *et al.*, 2011). Details of factors responsible for PSH in wheat are discussed below

2.1 Environment

Seed germination is the most vital step in the life cycle of plant and forms the basis of agricultural production. In contrast, seed dormancy is a phenomenon which prevents germination in spite of favourable environment and avoids losses due to untimely sprouting or germination (Nonogaki *et al.*, 2018). Although seed dormancy is regulated by number of genes, there exist evidences on the other hand that dormancy is also highly influenced by certain environmental factors. The climate change and its effect are evident in the form of extreme weather events. This leads to the occurrence of PHS more frequently which ultimately results in reduction of seed quality and food production shortages in cereals (Maity and Pramanik, 2013). The weather conditions before 14 days to physiological maturity severely affects seed dormancy (Strand, 1989) and continuous rain and high humidity along with higher temperature at maturity can cause PHS (Reddy *et al.*, 1985; Penfield and MacGregor, 2017). Rain enhances moisture on the spike and with an optimum temperature at the time of seed maturity creates suitable conditions for PHS prior to harvesting (Lunn *et al.*, 2002). The temperature and PHS in barley are positively correlated (Gualano and Benech-Arnold, 2009).

Drought like conditions coupled with high temperature enhanced PHS tolerance in white spring wheat variety Cunderdin, a PHS sensitive genotype to the levels similar to (DM2001) a dormant genotype (Biddulph *et al.*, 2005). Dry conditions with high-temperature before grain maturity leads to PHS tolerance and also found inverse relationship of low temperature with high dormancy rate in wheat (Thomason *et al.*, 2009). Certain environmental conditions related with temperature and rainfall at the time of seed maturity leads to the onset of PHS or LMA (late maturity alpha-amylase activity (Patwa and Penning, 2020). Hence, considering current climate change projections of enhanced temperature and untimely precipitation across the globe, risk of PHS is expected to increase and become a greater challenge in wheat producing areas in near future (Shorinola *et al.*, 2016).

Fig 1. Factors affecting Pre-harvest sprouting in wheat

2.2 Crop morphology

Pre-harvest sprouting (PHS) sensitivity is a complex trait which is controlled by genetic factors along with other factors, such as spike and plant morphology, germination inhibiting compounds in the bracts, stage of maturity of plants and environment. PHS is influenced by water-soluble germination inhibitors in seed coat and morphological structure of spikes *viz.*, erectness of spikes and openness of florets (King and Richards, 1984; Gatford *et al.*, 2002; Tan *et al.*, 2006). A number of simple morphological traits of the cereals can create an umbrella effect and so can be used to limit pre-harvest sprouting in cereals. The structure of the wax rather than its amount is considered as an important factor for repelling water entry (King, 1989). The presence of awns leads to uptake of water by the ear-heads and increase in PHS in wheat, but it still remains uncertain if this effect occurs under natural field conditions or the awn or the lemma or glume of awned varieties leads to such correlation (Ji *et al.*, 2018). Presence of epicuticular wax can resist entry of water into ear-head and also provide a physical barrier in reducing PHS but would have little effect on LMA (late maturity alpha-amylase) which is ensued by temperature fluctuations rather than moisture conditions. Barley with smooth and wild-type wax decreases PHS under artificial rainfall conditions as compared to non-glaucous, glossy mutants due to variations in water uptake by the earheads. An extremely glaucous spike phenotype should also be combined with awnless and erect spike phenotype which plays a major role in reducing water uptake (King and Von Wettstein-Knowles, 2000).

2.3 Seed coat colour

Although PHS is highly influenced by the certain environmental factors such as temperature and rain or

humidity at seed maturity, several physical, chemical and genetic factors could also influence the onset and severity of PHS. Physical attributes such as seed testa colour, presence/absence of awns and epicuticular wax are associated with differences in PHS sensitivity (Patwa and Penning, 2020). Several researchers concluded that seed dormancy is closely associated with red grain colour in wheat. Pre-harvest sprouting in wheat is associated with insufficient seed dormancy and it is reported that red seed coat is a traditional marker for resistance to pre harvest sprouting in wheat breeding (Yu-ichi et al. 2016). Many studies during last decade have associated red seed coat with higher level of seed dormancy and PHS tolerance. Catechin and proanthocyanidins (PAs) are synthesized through the flavonoid synthesis pathway are believed to be associated with red seed coat colour in wheat. Early cytogenetics studies suggested that three genes viz., *R-A1*, *R-B1* and *R-D1* on homeologous group- 3 chromosomes control seed coat colour and accumulation of catechin, a precursor of the red pigment that leads to prevention of seed germination having pleiotropic effect (Lin et al., 2016).

On an average, white grained wheat seeds have been reported to be more vulnerable to PHS than red colored wheat seeds, although variation is seen in both groups for resistance (Andreoli et al., 2006). However, PHS occurs occasionally in red seed coat varieties, implicating that the red-grain genotypes alone do not contribute towards resistance against PHS (Flintham, 2000). The varietal differences exist for pre-harvest sprouting (PHS) tolerance and associated grain characteristics viz., falling number (FN), α -amylase activity, degree of sprouting (DS) under field sprouting (FS) and spike wetting conditions (Ji et al., 2018). Awnless varieties appear to have better sprouting tolerance than awned varieties under intensive sprouting conditions, but found to be comparatively less evident than seed coat colour. The white seeds mutants of 'Chinese Spring' and 'AUS1490' showed increased germination under wet conditions, indicating that R genes are associated with PHS tolerance (Li et al., 2016). However, contribution of R genes to PHS resistance remains unsolved puzzle. Additional genetic effects like α -amylase activities, endogenous hormones levels, genes and QTLs, independent of seed coat pigmentation are known to be involved in the control seed dormancy in white and red colored seeds as reported by several researchers.

2.4 Seed dormancy

Dormancy is the inhibition of germination of morphologically ripe and healthy seeds even under optimum conditions of temperature, light and moisture (Chouard et al., 1960). Dormancy is a quantitative trait regulated by multiple genes, is strongly influenced by environmental conditions (Nakamura, 2018). There are two main environmental factors viz., temperature and moisture which affects pre-harvest sprouting in wheat especially during the late maturity stage (Hilhorst, 1995; Yanagisawa et al., 2005; Gao et al., 2006), because low temperature and high moisture both are able to break dormancy and promote sprouting of seed (Argel et al., 1983; Ceccato et al., 2011). Therefore, seed dormancy has been considered as major factor that determines pre-harvest sprouting resistance in wheat (Bewley and Black, 1982; Mares and Mrva, 2001; Finch-Savage and Leubner-Metzger, 2006; Lan et al., 2005; Lin et al., 2008; Yang et al., 2011). Dormancy is typically measured by either a germination index (ranging from 0.0 to 1.0) (Reddy et al., 1985) or germination resistance index (ranging from 0 to 50) calculated on hand threshed grains imbibed on filter paper.

Many researchers have reported dormancy as a key genetic constituent of PHS resistance (Sun et al., 2012; Yang et al., 2014). Mechanism of seed dormancy can also be classified as seed-coat imposed dormancy and embryo related dormancy. Former is associated with seed coat inhibitory compounds (Himi et al., 2002) and the later involves the crosstalk of plant hormones i.e. ABA (abscisic acid), GA (gibberellin), auxin etc. (Finch-Savage et al., 2006; Liu et al., 2013a). Seed coat imposed dormancy is involved particularly in seed survival mechanism. Dormancy is affected by both genetic and environmental factors (Cavangh et al., 2008). Red-grained wheat genotypes exhibit a wide range of seed dormancy and are more resistant to PHS as they contained dominant alleles, whereas white-grained cultivars are susceptible to PHS as they lack seed dormancy at maturity (Freed et al., 1976; Gfeller and Svejda, 1960; Everson et al., 1961; McEwan et al., 1980).

2.5 Genes associated with PHS tolerance

There are several genes which are identified and are found to be associated with PHS tolerance in wheat. Liu et al. (2013) observed *TaPHS1*, major gene determining

pre-harvest sprouting resistance and suggested that the pre-harvest sprouting resistance was independent of grain color (qualitative). However, seed color genes may modify the level of pre-harvest sprouting (quantitative) in either pre-harvest sprouting resistant or susceptible genotype groups as determined by *TaPHS1*. Therefore, *TaPHS1* is a highly valuable pre-harvest sprouting resistance gene for breeding white wheat cultivars. Himi and Noda (2005) reported that *R* genes regulate the expressions of the *CHS*, *CHI*, *F3H* and *DFR* genes that encode enzymes in the early steps of flavonoid synthesis. In fact the *Tamby10* genes appeared to be located in the same regions as the *R* loci, and *R*, *TaDFR* and *Tamby10* were proven to be *Myb* type transcription factors (Himi and Noda, 2005; Himi *et al.*, 2011; Bi *et al.*, 2014). It is reported in literature that red-grained wheat is more tolerant to sprouting than white-grained wheat (Probert, 2000; Warner *et al.*, 2000; Groos *et al.*, 2002). In wheat, three *TaSdr* genes viz., *TaSdr-A1*, *TaSdr-B1*, and *TaSdr-D1* have been cloned, and are involved in seed dormancy; among which, *TaSdr-B1* on chromosome 2B was observed to play a vital role in regulating seed dormancy (Zhang *et al.*, 2017b; Zhang *et al.*, 2014). Barrero *et al.* (2015) reported another gene *MKK3-A* (mitogen-activated protein kinase 3), also called *TaMKK3-A*, to be located on chromosome 4AL as a candidate gene of *Phs-A1* locus which is associated with duration of seed dormancy (Torada *et al.*, 2016). Another gene, *TaVp1* extensively studied in wheat was linked with seed dormancy and pre-harvest sprouting resistance (Yang *et al.*, 2007a; Chang *et al.*, 2010; Yang *et al.*, 2014; Chang *et al.*, 2011; Hattori *et al.*, 1992; Nakamura and Toyama 2001; Xia *et al.*, 2008; McKibbin *et al.*, 2002). This gene was mapped about 30 cM from *R* loci on homologous group 3L chromosomes (Bailey *et al.*, 1999). Further, there are reports that the wheat has several dormancy QTLs, including *QPhs.ocs-3A.1* and *Phs1* (Gong *et al.*, 2014; Mori *et al.*, 2005; Torada *et al.*, 2005; 2008) (Table 1).

3. Physiological, biochemical and molecular mechanism involved in pre-harvest sprouting

PHS tolerance / resistance is associated with a variety of developmental, physiological and morphological characteristics of the spike and seed, including seed coat color and permeability, seed dormancy alpha-amylase activity and plant growth hormone levels (abscisic acid, gibberellin and auxin). Seed dormancy is the most

important genetic element influencing PHS resistance; hence researchers have focused on understanding the molecular mechanism of seed dormancy as a way to improve PHS resistance in wheat breeding programmes.

Grain color (GC) is an important genetic factor affecting the brightness of flour and is also associated with seed dormancy and PHS resistance. It is controlled by the *R-1* gene series distally located on long arms of chromosomes 3A, 3B and 3D (Metzger and Silbaugh, 1970). Enzymes like DFR (dihydroflavonol-4-reductase), CHI (chalcone flavanone isomerase), F3H (flavanone 3-hydroxylase), and CHS (chalcone synthase) are expressed only in immature red grains and are nearly repressed in the grains of white wheat (Mir *et al.*, 2012; Thomas and Ougham, 2014). *Myb*-type *Tamby10-1* transcription factors control anthocyanin production and the red pigment of wheat grain by up-regulating the structural genes encoding DFR, CHI, F3H and CHS in flavonoid biosynthesis pathway (Ali *et al.*, 2019). Groos *et al.* (2002) identified four QTLs for both PHS and GC resistance, three of which were close to *R* genes and one was mapped on chromosome 5AS using a RIL population. Later on, Zhou *et al.* (2017) detected three main QTLs for PHS resistance, one on chromosome 5D and other two loci co-located with *Tamby10-1* genes on chromosomes 3A and 3D by Genome wide association mapping studies (GWAS). Recently, Lang *et al.*, (2021) mapped the PHS resistance gene *PHS-3D* from synthetic hexaploid wheat to a 2.4 Mb presence-absence variation (PAV) region and found that its resistance effect was attributed to the pleiotropic *Myb10-D* by integrated omics and functional analyses. Therefore, it is possible to breed PHS-tolerant white wheat by using CRISPR / Cas9 (a gene-editing technology) for altering GC-related genes keeping in view the other dormancy-related QTLs besides those provided by the *R-1* genes of the red grained parent used for such editing (Ali *et al.*, 2019).

3.1 α - Amylase activity

The α -amylase enzyme widely exists and participates in many physiology processes in plants, which can hydrolyze α -1, 4-glycosidic bond in the saccharides. The expression of α -amylase was involved in plant metabolism and could affect the germination rate, cold tolerance and production of seed (Autio *et al.*, 2001) and is strictly controlled by the phytohormones ABA and gibberellin. During grain development ABA inhibits the amylase expression, four

isozymes of α -amylase in wheat have been identified affecting PHS, namely malt- α -amylase (α -amylase-1) located on homologous chromosomes - 6, green- α -amylase (α -amylase-2) located on homologous chromosomes - 7, α -amylase-3 on chromosome -5 and α -amylase-4 has two members on homologous chromosomes -2 and 3 (Gale and Ainsworth, 1984; Zhang *et al.*, 2017c). Among all the three genomes, B genome harbors genes for α -amylase-1 and α -amylase-2. GA3 seemed to be involved in regulation of expression level of α -amylase-1 and α -amylase-2 (Marchylo *et al.*, 1983).

3.2 Plant growth hormones

The growth hormones, such as ABA and GA play important roles in regulation of dormancy and germination in wheat. ABA induces dormancy and GA stimulates seed germination (Kucera *et al.*, 2005; Finkelstein *et al.*, 2008). Several studies have reported the regulatory mechanisms of other hormones like ethylene, jasmonate brassinosteroids, and auxin in controlling seed dormancy, germination and PHS resistance (Liu *et al.*, 2013b; Kim *et al.*, 2019; Ju *et al.*, 2019). Light and temperature, also affect the dormancy and germination by disturbing the balance between ABA and GA levels in cereal crops (Gubler *et al.*, 2008; Lzydorczyk *et al.*, 2017). ABA is an essential hormone that promotes seed dormancy, seed maturation and tolerance to desiccation. During imbibition ABA levels increases upto 2.5-fold in dormant wheat and remains unchanged in non-dormant grains (Ried *et al.*, 1990). ABA level is regulated by its synthesis and catabolism. Zeaxanthin epoxidase (ZEP)/ABA1 and 9-cis-epoxycarotenoid dioxygenase (NCED) are two key enzymes involved in ABA synthesis and PHS regulation. ABA catabolism is mainly the result of hydroxylation by ABA 8'-hydroxylase (ABA8'OH)/cytochrome p450 monooxygenase 707A (CYP707A) (Okamoto *et al.*, 2006). The two genes NCED and CYP707A play important roles in germination and dormancy by controlling the ABA level in seeds. MYB96 and ABI4 can increase ABA levels by affecting the expression of NCED2/6 and CYP707A1/2, respectively (Shu *et al.*, 2013; Lee *et al.*, 2015). Chono *et al.*, (2013) reported the importance of higher embryonic ABA levels in inducing seed dormancy during seed maturation phase in wheat *via* mutational analysis of two homologs of *TaABA8'OH1* i.e. *TaABA8'OH1A* and *TaABA8'OH1D*. The resource of TILLING (Targeting Induced Local Lesions In

Genome) mutants, like Kronos and Cadenza, have been developed in wheat. The exome sequences of Kronos and Cadenza mutants have been re-sequenced using Illumina NGS that can be used to screen for mutations in pre-harvest sprouting and dormancy related genes (Nakamura *et al.*, 2011).

GA is another key plant hormone that plays a significant function in seed dormancy and germination control (Finch-Savage *et al.*, 2006). GA breaks seed dormancy and promotes germination mostly by overcoming the mechanical restraint imposed by the aleurone or testa and stimulating the growth of the embryo (Debeaujon and Koornneef, 2000). It also regulates the expression of amylase synthesis genes, which are involved in seed germination and starch hydrolysis in the endosperm. The bioactive GA content in plants is controlled by an equilibrium between its synthesis and inactivation, mediated principally by GA2ox genes (GA2-oxidase encoding), GA3ox (GA3-oxidase encoding), and GA20ox (encoding GA 20-oxidase), respectively (Yamaguchi *et al.*, 2008). In wheat, six GA metabolism genes, *TaKS*, *TaKO1*, *TaKA01*, *TaGA20ox1*, *TaGA2ox1*, and *TaGA2ox8*, were detected by GeneChip in dry dormant and after-ripened grains, indicating that they might participate in dormancy regulation (Liu *et al.*, 2013b). In a number of cultivated species, including wheat, rice, and barley many genes were identified which encoded these enzymes and their expression played an important role in dormancy and germination by adjusting seed GA levels (Pearce *et al.*, 2015). The role of GA in regulating seed germination and dormancy was shown by the transcription level of these gene orthologs due to post-ripening as well as in non-dormant and dormant cereal crops. In imbibed post-ripened barley and wheat seeds, for example, the dormancy loss was demonstrated to be related with an enhanced expression of the *TaGA3ox* and *TaGA20ox* genes and to a higher bioactive *GA1* content (Liu *et al.*, 2013c; Gubler *et al.*, 2008; Kashiwaruka *et al.*, 2016). GA signals in plants are perceived by the soluble receptor protein gene *GID1* (Gibberellin insensitive dwarf 1), that was first mapped in rice (Ueguchi-Tanaka *et al.*, 2007). Reduced height (*Rht*)-1 in wheat is an ortholog of *GAI* (Peng *et al.*, 1999). Notably, *Rht*-B1b (also known as *Rht1*) and *Rht*-D1b (also known as *Rht2*) were used to reduce lodging and to increase grain yield during the Green Revolution (Eshed and Lippman, 2019). Two intragenic *Rht*-B1c suppressor

alleles (*Rht-B1c.23* and *Rht-B1c.26*) reported by Van De Velde *et al.*, (2017) showed improved PHS resistance and have no negative effects on grain yield or quality.

3.3 QTLs Identified for PHS resistance

PHS is a quantitative trait controlled by multiple QTLs or genes that are important for breeding PHS-resistant varieties. Recent advances in the genomics of cereal crops have led to the identification of several genes involved in PHS resistance. PHS resistance QTLs have been reported almost on all wheat chromosomes accounting varying degree of resistance (Tai *et al.*, 2021; Zhou *et al.*, 2017). In wheat, more than 250 QTLs for PHS resistance have been identified in ~40 mapping populations that included hexaploid synthetic wheats, tetraploid durum wheats and diploid einkorn wheats (Gupta *et al.*, 2020). The PHS resistance genes on chromosomes 3A, 3B and 3D regions which encode a Myb transcription factor (*TaMyb10*) are considered to be tightly linked or pleiotropic with red seed coat color determined by dominant R alleles (Himi *et al.*, 2011). A QTL on chromosome 3AS encode for Mother of Flowering Time (*TaMFT*: was cloned which promotes seed dormancy in wheat embryos (Nakamura *et al.*, 2011; Lei *et al.*, 2013; Liu *et al.*, 2013c). A total of 188 QTLs or loci from 40 previous studies associated with resistance to PHS in wheat have been reported (Tai *et al.*, 2021). In a recent study, three *Ae. tauschii*-derived quantitative trait loci (QTLs), *QDor.3D.1*, *QDor.3D.2*, and *QDor.3D.3*, were detected on chromosome 3DL using four simple sequence repeats (SSR) markers and developed 10 Kompetitive allele-specific PCR (KASP) markers (He *et al.*, 2021). Furthermore, multiple potential genes for PHS resistance have been found using comparative genomics or transcriptome analysis, including *TaSdr-A1* and *TaSdr-B1* on chromosome 2AS and 2BS, *TaPHS1* and *TaMFT* on chromosome 3AS, *TaVp-1* and *Tamyb10* on group- 3 chromosomes, and two tandem genes, *PM19-A1/A2* encoding ABA-inducible Plasma Membrane 19 proteins and *TaMKK3-A* on 4AL (Ali *et al.*, 2019). Many gene-specific markers, including SSRs (*Xgwm15*, *Xgwm894* & *Xgwm937*), STMS markers (*Xwmc104*, *Xwmc397*, and *Xwmc468*), and STS markers (*Vp1-B2* and *Vp1-B3*), were produced for the *Vp1* and can be used to identify PHS resistance in various genotypes. Ogonnaya *et al.* (2007) discovered that the *Xgwm894* and *Xgwm937* markers were significantly linked with PHS resistance and could

be exploited to improve PHS resistance in wheat. PHS resistance is governed by both epistatic and additive effects that are influenced by environmental factors. To better understand the complicated genetic structure of QTL, the interaction between QTL epistasis (QXQ) and the environment (Q X E, Q X Q X E) for PHS resistance was explored (Liu *et al.*, 2010). Rapid advances in molecular technologies, such as NGS technologies (Brenchley *et al.*, 2012), and ongoing chromosomal-based and wheat whole genome sequencing projects (International Wheat Genome Sequencing Consortium, IWGSC) (Clavijo *et al.*, 2017; IWGSC 2018), would provide new opportunities for identification and functional analysis of the candidate genes controlling PHS resistance. Liu *et al.* (2010) designed the *TaPHS1*-SNP1 marker, which may be utilized as a diagnostic marker in breeding to identify the *TaPHS1* resistance allele. Rasheed *et al.* (2016) identified five KASP-based functional marker assays for four PHS resistance genes: SDR SNP for *TaSdr-B1*, *TaMFT*-1617R for *TaMFT-A1*, *TaMFT*-721J for *TaMFT-A1*, *Vp1B1*-83 IND for *TaVp-1B* and *Vp1B1*-193 IND for *TaVp-1B*. Furthermore, the CAPS (cleaved amplified polymorphism sequence) marker *Sdr2A*, which is located between the *Xgwm95* and *Xgwm372* markers, has been established and can be used for identifying PHS resistant genotypes (Zhang *et al.*, 2017).

Genome-wide association (GWA) studies have also facilitated the discovery of variant sites regulating PHS (Table 1). Using GWAS, more than 250 MTA (marker trait associations) for PHS and related traits have been identified. Among known QTLs, 30 QTLs were also shown to be stable over environments; these were considered potentially useful for improvement of PHST in wheat (Gupta *et al.*, 2020). Recently, Tai *et al.* (2021) reported 66 meta-QTLs (MQTLs) distributed on all 21 wheat chromosomes out of 188 valid QTLs for PHS resistance from 40 articles. A major QTL for PHST was introgressed into an elite Indian wheat cv. Lok1 that is PHS susceptible using marker-assisted backcross breeding (Gautam *et al.*, 2021). These PHST (pre-harvest sprouting tolerant) lines were also pyramided with one gene each for high grain protein content (*Gpc-B1*) and leaf rust resistance (*Lr24*). Recently, Wang *et al.* (2021) assessed the distribution of three causal single nucleotide polymorphisms in *TaPHS1* at bases -222, +646, and +666 in 725 Chinese wheat accessions.

Table 1. List of QTL/genes identified for PHS resistance

Trait	Population	QTLs	Chromosome	Nearest Marker	QTL name	Reference
PHS and GC	RILs	5	3AL	<i>Xffb293</i>		Groos <i>et al.</i> (2002)
			3BL	<i>Xgwm403</i>		
			3BL	<i>Xbcd131</i>		
			3DL	<i>Xgwm3</i>		
			5AS	<i>Xbcd1871</i>		
PHS and SD	RILs	3	3A	<i>Xpsr394-Xgwm5</i>	<i>taVp1</i>	Osa <i>et al.</i> (2003)
			3A	<i>Xcdo345</i>	<i>QPhs.ocs-3A.1</i>	
			3A	<i>Xpsp3050</i>	<i>QPhs.ocs-3A.2</i>	
PHS	DHLs	4	2B	<i>Xbarc55-Xwmc474</i>	<i>QPhs.cnl-2B.1</i>	Munkvold <i>et al.</i> (2009)
			2D	<i>Xwmc111-WxPt-999</i>	<i>QPhs.cnl-2D.1</i>	
			3D	<i>7Xbarc1161</i>	<i>QPhs.cnl-3D.1</i>	
			6D	<i>Xcfd37-Xbarc196</i>	<i>QPhs.cnl-6D.1</i>	
PHS and GC	RILs	6	3AL	<i>Xwmc559-1</i>		Lin <i>et al.</i> (2016)
			3AL	<i>Tamyb10-A1-74</i>		
			3AL	<i>Tamyb10-A1-74</i>		
			3DL	<i>BS00067163_51</i>		
			3DL	<i>Tamyb10-D1-93</i>		
			1A/1D/3A/5B	<i>Xbarc148</i>		
PHS and GC	DHLs	5	3B	<i>Xbarc77-Xwmc30</i>	<i>QGi.crc-3B</i>	Fofana <i>et al.</i> (2009)
			3D	<i>7Xwmc552-Xwmc533</i>	<i>QGi.crc-3D</i>	
			3A	<i>Xcfa2193-Xwmc594</i>	<i>QSi.crc-3A</i>	
			5D	<i>Xgwm469-Xcfd10</i>	<i>QSi.crc-5D</i>	
			3D	<i>Xwmc11-Xcfd223</i>	<i>QCli.crc-3D</i>	
PHS	RILs		2AL	<i>Xgwm1045-Xgwm296</i>	<i>QPhs.ccsu-2A.5</i>	Mohan <i>et al.</i> (2009)
			3AL	<i>Xgwm153-Xgwm155</i>	<i>QPhs.ccsu-3A.1</i>	
			3BL	<i>Xgwm1005-Xgwm980</i>	<i>QPhs.ccsu-3B.6</i>	
PHS	DHLs	1	5D	<i>XCfD40-XBARC1097</i>	<i>qPhs5D.1</i>	Zhu <i>et al.</i> (2010)
PHS	RILs	3	1A	<i>Xwmc611-Xwmc333</i>	<i>QPhsd.spa-1A.1</i>	Knox <i>et al.</i> (2012)
			2A	<i>Xgwm515-Xgwm425</i>	<i>QPhsd.spa2A.1</i>	
			7B	<i>Xgwm297-Xwmc532</i>	<i>QPhsd.spa-7B.1</i>	
PHS	DHLs	4 + 18 KASP markers developed	3B	19 SNPs flanking the QTL	<i>QSi.crc-3B</i>	Cabral <i>et al.</i> (2014)
			4A	12 SNPs flanking the QTL	<i>QGi.crc-4A</i>	
			7B	10 SNPs flanking the QTL	<i>QSi.crc-7B</i>	
			7D	04 SNPs flanking the QTL	<i>QFn.crc-7D</i>	
PHS	DHLs	5	1A	<i>wPt-6274</i>	<i>QPhs.spa-1A</i>	Singh <i>et al.</i> (2014)
			1B	<i>Xwmc191</i>	<i>QPhs.spa-1B</i>	
			5B	<i>wPt-6910-wPt-7400</i>	<i>QPhs.spa-5B</i>	
			7A	<i>Xcfa2174</i>	<i>QPhs.spa-7A</i>	
			7B	<i>Xwmc606</i>	<i>QPhs.spa-7B</i>	
PHS and SD	RILs	1	2B	<i>Xwmc477-Xbarc55</i>	<i>Sdr2B</i>	Zhang <i>et al.</i> (2014)
PHS and SD	RILs	1	4A	<i>wsnph_Ex_c66324_64493429-CD920298</i>	<i>4A-1</i>	Barrero <i>et al.</i> (2015)
PHS and SD	RILs	4	4A	<i>GBS212432-GBS10994</i>	<i>QPhs.pseru-4A.1</i>	Lin <i>et al.</i> (2015)
			4B	<i>7Xbarc20-Xwmc238</i>	<i>QPhs.pseru-4B.1</i>	
			5A	<i>TTM_199619-TTM_1259</i>	<i>QPhs.pseru-5A.1</i>	
			5B	<i>7Xbarc346-2-TTm_62137_50</i>	<i>QPhs.pseru-5B.1</i>

PHS	RILs	6	3A	<i>TaMFT</i>	<i>QDor-3A</i>	Cao <i>et al.</i> (2016)
			4A	<i>Cfa2256</i>	<i>QDor-4A</i>	
			1B	<i>Xbarc181</i>	<i>QDor-1B</i>	
			7B	<i>UCW99</i>	<i>QDor-7B</i>	
			4A	<i>Cfa2256</i>	<i>QAwn-4A</i>	
			6B	<i>Xwmc397</i>	<i>QAwn-6B</i>	
PHS and SD	RILs	1	2A	<i>Xgwm95-Xgwm372</i>	<i>Sdr2A</i>	Zhang <i>et al.</i> (2017a)
PHS and SD	Back cross population	3	2D	<i>Xwmc503</i>	<i>QDor-2D</i>	Dale <i>et al.</i> (2017)
			3D	<i>Xcfd22</i>	<i>QDor-3D</i>	
			3D	<i>Vp1-4</i>	<i>TaVp1</i>	
PHS	717 Chinese Wheat landraces	3	3A	<i>7AX-111578083</i>	<i>QTL1</i>	Zhou <i>et al.</i> (2017)
			3D	<i>3DArT-seq & 5 SNPs</i>	<i>QTL2</i>	
			5D	<i>AX-109028892</i>	<i>QTL3</i>	
PHS	CM32 and SHW-L1	5 eQTLs	2A	<i>AX-110609678</i>	<i>eQABI4.15DPA.2A.1</i>	Xiao <i>et al.</i> (2021)
			2D	<i>AX-110515525</i>	<i>eQABI4.20DPA.2D.1</i>	
			2D	<i>AX-111690676</i>	<i>eQABI4.20DPA.2D.2</i>	
			3B	<i>AX-95660238</i>	<i>eQABI4.20DPA.3B.1</i>	
			4A	<i>AX-86175059</i>	<i>eQABI4.20DPA.4A.1</i>	
			4A	<i>AX-110113739</i>	<i>eQABI4.20DPA.4A.2</i>	
PHS	CSSL	3QTL	3DL	SSR&KASP	<i>QDor.3D.1</i>	He <i>et al.</i> (2021)
			3DL	SSR&KASP	<i>QDor.3D.2,</i>	
			3DL	SSR&KASP	<i>QDor.3D.3</i>	
PHS	SHW-L1			KASP	<i>QPHS.sicau-3D</i> (PHS-3D)	Lang <i>et al.</i> (2020)
PHS	211 DH + 167 US winter wheat variety	4 + KASP	2AS	KASP-222	<i>Qphs.hwmg-2A.1</i>	Shao <i>et al.</i> (2018)
			3AS	KASP-314	<i>Qphs.hwmg-3A.1</i>	
			3BS	KASP765,471	<i>Qphs.hwmg-3B.1</i>	
			5AL	KASP8426	<i>Qphs.hwmg-5A.1</i>	
PHS	SHW-L1, Chuanmai 32, AS225, AS60	2	3DL	<i>AX-94415259</i>	<i>qPHS.sicau-3D</i>	Yang <i>et al.</i> (2019)
			1BS	<i>AX-94924265</i>	<i>qPHS.sicau-1B</i>	
PHS	RILs	5	1D	BS00022188_51 Kukri_c10485_1346	<i>QPhs.umb-1D</i>	Liton <i>et al.</i> (2021)
			4A.2	BS00068243_51 BS00037019_51	<i>QPhs.umb-4A</i>	
			6B.1	RAC875_c63209_154 BobWhite_c1509_622	<i>QPhs.umb-6B</i>	
			6D	Jagger_c211_207 IAAV7251	<i>QPhs.umb-6D</i>	
			7A	Ku_c8523_166	<i>QPhs.umb-7A</i>	
				Kukri_c38390_218		

4. Breeding approaches for pre-harvest sprouting tolerance in wheat

The development of pre-harvest sprouting resistance lines offers basis for the enrichment of genetic resources in wheat breeding programs; accordingly, introducing genes from wheat-related species could provide an effective way to significantly improve wheat grain quality and its resistance to pre-harvest sprouting (He *et al.*, 2021). Wheat germplasm contains an appreciable amount of variation

for pre-harvest sprouting characteristics which could be further exploited to develop pre-harvest sprouting resistance varieties (Wang *et al.*, 2020). The plant breeding strategy for introducing tolerance to pre-harvest sprouting in wheat is to incorporate major genes from different wild relatives or derived lines into modern breeding lines (Andreoli *et al.*, 2006). Details of exotic landraces, varieties and lines exhibiting PHS tolerance are presented below (Table 2).

Table 2. Details of wild relatives / land races/ lines identified having PHS tolerance in wheat

Wild relatives/ land races/ lines identified with PHS tolerance	Country	References
<i>Aegilops tauschii</i>	China	Liu <i>et al.</i> (1998)
<i>Triticum turgidum</i> and <i>Aegilops tauschii</i> (DD)	China	Liu <i>et al.</i> (1998); Yu <i>et al.</i> (2014)
Synthetic hexaploid wheat genotypes (SHW-L1)	China	Yang <i>et al.</i> (2014)
Chinese landraces viz., Xiaoyuhua, Yongchuanbaike and Baiyuhua	China	Xiao <i>et al.</i> (2002)
Hongshangtou and Wanxianbaimaizi	China	Yang <i>et al.</i> (2009); Chang <i>et al.</i> (2010)
RL4137	Canada	Bassoi and Flintham, (2005);
BRS177	Brazil	Flintham, (1993)
RL4137	Canada	DePauw <i>et al.</i> (2012)
Konde, Kumpa and Windy cultivars	Australia	Jimenez <i>et al.</i> (2017)

The resistance to pre-harvest sprouting in crops is a complex trait which is conditioned by both genetic and environmental factors (Derera *et al.*, 1977, Torada *et al.*, 2008; Barrero *et al.*, 2015). Therefore, the breeding approaches which can be used for the improvement of pre-harvest sprouting improvement in wheat are presented below (Fig. 2).

Fig. 2. Breeding approaches for PHS tolerance

Breeding varieties with high pre-harvest sprouting resistance have important implications for reducing yield loss and improving grain quality (Ali *et al.*, 2019). Therefore, breeding for genetically improved wheat with

proper seed dormancy is the most effective approach to protect wheat cultivars from pre-harvest sprouting damages (Liu *et al.*, 2008). The wild wheat species (*A. tauschii*) are more dormant (conferring PHS resistance) than domesticated wheat indicating that artificial selection has contributed to the reduction of dormancy in modern wheat (Dong *et al.*, 2015; Volis, 2016). The red grained wheat usually shows more resistance to pre-harvest sprouting as the red colour genes on the long arm of group 3 chromosomes can have pleiotropic effects on pre-harvest sprouting resistance (Groos *et al.*, 2002). Breeding for pre-harvest sprouting resistance in wheat based on phenotypic selection is challenging, because pre-harvest sprouting is expressed as a quantitative character that is influenced by environment and genotype x environment interactions (Anderson *et al.*, 1993). Therefore, selecting for pre-harvest sprouting resistant genotypes on a phenotypic basis at early stages of segregation may not be efficient (Singh *et al.*, 2012).

Several quantitative trait loci or genomic regions affecting pre-harvest sprouting resistance or seed dormancy in wheat have been identified in different gene pools *via* linkage with molecular markers. Therefore, marker-assisted selection was applied in many breeding programs as pre-harvest sprouting resistance genes were cloned on chromosomes 3A (Nakamura *et al.*, 2011; Liu *et al.*, 2013c) and 4A (Torada *et al.*, 2016), and also QTLs for pre-harvest sprouting resistance identified across the genome. The use of molecular markers for marker-assisted selection could be helpful for direct identification of favorable or deleterious alleles in diverse groups of genotypes (Lazo *et al.*, 2004). The first QTL found for grain dormancy in wheat was reported in 1993 (Anderson *et al.*, 1993). Molecular markers have been developed for the causal SNPs in both the promoter and coding region of the TaPHS1 gene (Liu *et al.*, 2015), and the causal SNP for the TaMKK gene (Torada *et al.*, 2016). These markers can be used either to identify germplasm carrying these two genes, or to select resistant lines in breeding materials. The TaPHS1 gene has been successfully transferred through marker assisted selection approach to increase pre-harvest sprouting resistance (Kottearachchi *et al.*, 2006; Gupta *et al.*, 2008). Singh *et al.* (2014) also used simple sequence repeat (SSR) and Diversity Arrays Technology markers (DART) on Double Haploid (DH) Canadian adapted durum wheat. Kumar *et al.* (2010) developed individual

plant by marker assisted breeding and phenotyped for pre-harvest sprouting tolerance to verify the effect of introgression of pre-harvest sprouting tolerance QTL. However, due to some undesirable traits from the linkage drag, these QTLs have not been widely used in wheat breeding.

Metabolomics is considered as one of the useful approach in understanding the dynamic metabolic changes since metabolomic analysis measures the metabolite composition of cells and its response to external factors. Liu *et al.* (2015) used transgene anti-trx-s byre-programming of 's' metabolism in seeds especially at later seed development (dough development, seed maturing) and post-harvest ripening stages using metabolomes. They concluded that sugars, organic acids, amino acids, choline metabolites and fatty acids of the transgenic and wild-type (control) wheat differed significantly. After 30-days post-harvest ripening, most metabolites in transgenic seeds had higher levels than in controls including amino acids, sugars, organic acids, fatty acids and choline metabolites. These clearly indicated that transgene reduced metabolic activities of mature seeds responsible for pre-harvest sprouting.

Breeding for pre-harvest sprouting resistance for various environments is challenging due to its polygenic nature, poor understanding of the trait, non-standardization of phenotyping techniques and non-availability of suitable donors having seed dormancy. In order to improve pre-harvest sprouting resistance, more information on PHS resistance genetic architecture, PHS resistance pathways and gene regulations, genetic-by-environment interactions, based user-friendly markers and efficient selection method is required.

5. Methods to measure pre-harvest sprouting in wheat

The evaluation of pre-harvest sprouting resistance under field conditions depends on the presence of weather conditions conducive for seed sprouting after physiological maturity; thus, pre-harvest sprouting phenotypic data may not be repeatable in different environments (Graybosch *et al.*, 2013; Kato *et al.*, 2001). Furthermore, environment and genotype $\times$ environment interactions influence tolerance to PHS (Kato *et al.*, 2001). Therefore, pre-harvest sprouting damage can be assessed by exposing spikes to artificial rain or through germination tests or by natural/ artificial

weathering of the spike (Okuyama *et al.*, 2020) under controlled conditions. Following tests are used to measure Pre-harvest sprouting in wheat.

5.1 Germination test

Germination testing is a useful and easy method for assessing pre-harvest sprouting resistance of different wheat genotypes. Wu and Carver (1999) demonstrated that percentage of germination had good association with field assessment of sprout damage. Percentage of germination was negatively associated with seed dormancy level or sprouting resistance (Biddulph *et al.*, 2008). The genotypes with a higher percentage of sprouted grains resulted in lower falling number values; therefore, these results permitted to suggest that in those cases where the falling number test could not be performed, the germination percentage could be used as a selection tool for pre-harvest sprouting (Okuyama *et al.*, 2020). Further, association between grain germination percentage and falling number allows the wheat breeding programs to have a useful criterion for the selection, early discarding of unsuitable lines and planning a new cycle of hybridization for resistance to pre-harvest sprouting.

5.2. Falling number test

Falling Number test was developed by Hagberg (1960) as a rapid, high throughput method for determining alpha-amylase activity in grain and later adopted by industry as a test for sprout-damaged grain (Best and Muller, 1991). The low falling numbers values result from high levels of the enzyme alpha-amylase (Yu *et al.*, 2015). The falling number test has its limitations in use for breeding owing to the need of an expensive instrument, and the much time-consuming process as compared to spike-wetting tests (Martinez *et al.*, 2018).

6. Screening techniques and approaches for Indian conditions

Although all the methods used for the phenotyping pre-harvest sprouting are time-consuming and labor intensive, under natural condition of field screening is much easier. Under natural conditions of North East India, where pre monsoon rainfall during normal maturity time of wheat is a normal phenomenon, field screening may be effectively done. Such attempt is known to have done in Shillongani, Assam, India for screening wheat genotypes for PHS tolerance. Moreover, as already discussed, identification

of the molecular markers associated with genes/ QTLs for pre-harvest sprouting may lead to successful breeding of tolerant varieties. Marker-assisted selection is a better way to evaluate pre-harvest sprouting (He *et al.*, 2021). However, the success of marker-assisted selection depends on the identification of molecular markers tightly linked with quantitative trait loci (QTLs) for pre-harvest sprouting resistance (Gao *et al.*, 2013; Kulwal *et al.*, 2012; Mares and Mrva, 2014). Thus, the identification of reliable markers closely linked to pre-harvest sprouting resistance QTLs is crucial for marker-assisted selection method (Kulwal *et al.*, 2012, Gao *et al.*, 2013; Anderson *et al.*, 1993). Kulwal *et al.*, (2005) conducted QTL analysis for pre-harvest sprouting tolerance involving chromosome group-3A using a trait-specific inter-varietal recombinant inbred mapping population derived from a cross SPR8198 (a PHS tolerant genotype) and HD2329 (a PHS susceptible cultivar), and identified a major QTL on 3A- chromosome for pre-harvest sprouting tolerance which explained up to 78.03% of phenotypic variation. Therefore, it could be suggested that the bread wheat genotypes, SPR8198), which carry the desired allele of the major QTL detected might be exploited for developing pre-harvest sprouting tolerant varieties of wheat.

A combination of linkage analysis and association mapping could be the best approach for detecting maximum number of marker-trait associations to be used for molecular breeding (Jaiswal *et al.*, 2012). However, only a few reports on QTL analysis for tolerance to pre-harvest sprouting are available at national level. Department of Genetics and Plant Breeding, Ch. Charan Singh University, Meerut, UP, India has made an excellent progress for identification of QTLs associated with pre-harvest sprouting tolerance in hexaploid wheat, and development pre-harvest sprouting tolerant wheat genotype through marker assisted selection (Kulwal *et al.*, 2005). A major QTL (*QPhs.ccsu-3A.1*) for tolerance to pre-harvest sprouting was mapped on the long arm of wheat chromosome 3A (3AL) using a bi-parental population involving one parental genotype with red grain. This QTL (*QPhs.ccsu-3A.1*), later on was successfully transferred into the cultivar, HD2329 using marker-assisted selection (Kumar *et al.*, 2010).

7. Conclusions and future prospects

Breeding varieties for pre-harvest sprouting tolerance is very difficult owing to its genetic complexities,

environmental effects and over all poor understanding of the basis and inheritance of the trait. PHS is controlled by polygenes and is difficult to score in the segregating generations. Considering the availability of less variability for the trait, it is important to screen large number of germplasm including wild progenitors to identify new donors for this trait. These identified sources will be used in developing improved tolerant breeding material with added variability for the trait. Few genes/QTLs that have been cloned and linked markers developed could be useful for the transfer of this trait through marker assisted breeding programmes. Back cross breeding could be effectively applied for introgression of major QTLs identified for PHS tolerance. With the availability of next generation sequence data, it is now a day's possible to dissect the traits to few Kb regions to enable cloning of the genes concerned. The new Omics technologies would be very useful in the near future to further elucidate the underlying mechanisms contributing to PHS tolerance. Alternatively, variability available in other species/ genera may also be explored which could pave the way for development of transgenic/ cisgenic wheat in near future that are resistant to pre harvest sprouting.

Conflict of Interest

Authors declare that they have no conflict of interest.

Ethical Compliance Statement

NA

Author's Contribution

CS, URK, VG: Collection of literature, Conceptualization, Compilation, Writing original draft, Correspondence; GS, SS, AG, BST, PK: Editing, Data Compilation and analysis; CNM, GK, SKB, AKS, SK, GPS: Final editing, Proof Reading

8. References

1. Ali A, J Cao, H Jiang, C Chang, HP Zhang, SW Sheikh and C Ma. 2019. Unraveling molecular and genetic studies of wheat (*Triticum aestivum* L.) resistance against factors causing pre-harvest sprouting. *Agronomy* 9(3): 117.
2. Anderson JA, MES Sorrells and SD Tanksley. 1993. RFLP analysis of genomic regions associated with resistance to pre-harvest sprouting in wheat. *Crop Science* 33: 453–459.

3. Andreoli C, C Bassoi-MI and D Brunetta. 2006. Genetic control of seed dormancy and pre-harvest sprouting in wheat. *Scientific Agriculture* **63**(6): 564–566.
4. Anonymous. 2021. Progress Report of AICRP on Wheat and Barley 2020-21, Crop Improvement. eds: G Singh, BS Tyagi, A Gupta, SK Singh, S Kumar, H Khan, V Gupta, K Gopalareddy, AK Sharma, CN Mishra, C Singh, HM Mamrutha, K Venkatesh, UR Kamble, R Tiwari, A Verma and GP Singh. ICAR-Indian Institute of Wheat and Barley Research, Karnal, Haryana, India. p.227.
5. Argel PJ and LR Humphreys. 1983. Environmental effects on seed development and hard seededness in *Stylosanthes hamata* cv. Verano. I. Temperature. *Australian Journal of Agricultural Research* **34**: 261–270.
6. Autio K, T Simoinen, T Suortti, M Salmenkallio-Marttila, K Lassila and A Wilhelmson. 2001. Structural and Enzymic Changes in Germinated Barley and Rye. *Journal of the Institute of Brewing*. **107**: 19–25.
7. Bailey PC, RS McKibbin, JR Lenton, MJ Holdsworth, JE Flintham and MD Gale. 1999. Genetic map locations for orthologous Vp1 genes in wheat and rice. *Theoretical and Applied Genetics* **98**: 281–284.
8. Barrero JM, C Cavanagh, KL Verbyla, JFG Tibbits, AP Verbyla, BE Huang, GM Rosewarne, S Stephen, P Wang and A Whan, P Rigault, MJ Hayden and F Gubler. 2015. Transcriptomic analysis of wheat near-isogenic lines identifies PM19-A1 and A2 as candidates for a major dormancy QTL. *Genome Biology* **16**: 93.
9. Bassoi MC and J Flintham. 2005. Relationship between grain colour and pre-harvest sprouting-resistance in wheat. *Pesquisa Agropecuária Brasileira*. **40**: 981–988.
10. Best S and R Muller. 1991. Use of the hagberg falling number apparatus to determine malt and barley quality. *Journal of the Institute of Brewing*. **97**: 272–278.
11. Bewley JD, and M Black. 1982. Viability and longevity. In *Physiology and Biochemistry of Seeds in Relation to Germination*, eds J. D. Bewley and M. Black (Berlin: Springer), 1–59.
12. Bewley JD, M Black and P Halmer. 2006. The encyclopedia of seeds science. In *Technology and Uses*; CABI Publishing: Oxfordshire, UK, 2006; p. 528.
13. Bi HH, YW Sun, YG Xiao and LQ Xia. 2014. Characterization of DFR allelic variations and their associations with pre-harvest sprouting resistance in a set of red-grained Chinese wheat germplasm. *Euphytica* **195**: 197–207.
14. Biddulph TB, DJ Mares and JA Plummer et al. 2005. Drought and high temperature increases pre-harvest sprouting tolerance in a genotype without grain dormancy. *Euphytica* **143**: 277–283.
15. Biddulph TB, JA Plummer and Setter TL et al. 2007. Influence of high temperature and terminal moisture stress on dormancy in wheat (*Triticum aestivum* L.). *Field Crops Research* **103**: 139–153.
16. Biddulph TB, JA Plummer, TL Setter and DJ Mares. 2008. Seasonal conditions influence dormancy and pre-harvest sprouting tolerance of wheat (*Triticum Aestivum* L.) in the field. *Field Crops Research* **107**(2): 116–128.
17. Brenchley R, M Spannagl, M Pfeifer, GLA Barker, R D-Amore, AM Allen, N McKenzie, M Kramer, A Kerhornou and D Bolser et al. 2012. Analysis of the bread wheat genome using whole-genome shotgun sequencing. *Nature* **491**: 705.
18. Cabral AL, MC Jordan, CA McCartney, FM You, DG Humphreys, R MacLachlan and CJ Pozniak. 2014. Identification of candidate genes, regions and markers for pre-harvest sprouting resistance in wheat (*Triticum aestivum* L.). *BMC Plant Biology* **14**: 340–351.
19. Cao L, K Hayashi, M Tokui, M Mori, H Miura and K Onishi. 2016. Detection of QTLs for traits associated with pre-harvest sprouting resistance in bread wheat (*Triticum aestivum* L.). *Breeding Science* **66**(2): 260–70.
20. Cavanagh C, M Morell, I Mackay and W Powell. 2008. From mutations to magic: Resources for gene discovery, validation and delivery in crop plants. *Current Opinion and Plant Biology* **11**: 215–221.
21. Ceccato DV, HD Bertero and D Batlla. 2011. Environmental control of dormancy in Quinoa (*Chenopodium Quinoa*) seeds: two potential genetic

- resources for pre-harvest sprouting tolerance. *Seed Science Research* **21**: 133–141.
22. Chang C, HP Zhang, QX Zhao, JM Feng, HQ Si, J Lu and CX Ma. 2011. Rich allelic variations of Viviparous-1A and their associations with seed dormancy/pre-harvest sprouting of common wheat. *Euphytica* **179**: 343–353.
23. Chang C, JM Feng, HQ Si, B Yin, HP Zhang and CX Ma. 2010. Validating a novel allele of viviparous-1 (Vp-1Bf) associated with high seed dormancy of Chinese wheat landrace, Wanxianbaimaizi. *Molecular Breeding* **25**: 517–525.
24. Chono M, H Matsunaka, M Seki, M Fujita, C Kiribuchi-Otobe and S Oda, et al. 2013. Isolation of a wheat (*Triticum aestivum* L.) mutant in ABA8'-hydroxylase gene: effect of reduced ABA catabolism on germination inhibition under field condition. *Breeding Science* **63**: 104–115.
25. Chouard P. 1960. Vernalization and its Relations to Dormancy. *Annual Review in Plant Physiology* **11**: 191–238.
26. Cunha GR, JLF Pires and A Pasinato. 2004. Introdução ao problema da germinação na pré-colheita em trigo no Brasil. In: Cunha, G.R. and Pires, J.L.F., Eds., *Germinação pré-colheita em trigo*. Embrapa, Passo Fundo, 11-20.
27. Dale Z, H Jie, H Luyu, Z Cancan, Z Yun, S Yaru and Suoping L. 2017. An Advanced Backcross Population through Synthetic Octaploid Wheat as a “Bridge”: Development and QTL Detection for Seed Dormancy. *Frontiers in Plant Science* **8**: 2123.
28. Debeaujon I, KM Leon-Kloosterziel and M Koornneef. 2000. Influence of the Testa on Seed Dormancy, Germination, and Longevity in Arabidopsis. *Plant Physiology* **122**: 403–414.
29. Depauw RM, P Hucl, RE Knox, AK Singh, SL Fox and DG Humphreys. 2012. Developing standardized methods for breeding pre-harvest sprouting resistant wheat, challenges and successes in Canadian wheat. *Euphytica* **188**: 7–14.
30. Derera NF, GM Bhatt and GJ McMaster. 1977. On the problem of pre-harvest sprouting of wheat. *Euphytica* **26**: 299–308.
31. Dong ZD, J Chen, T Li, F Chen and DQ Cui. 2015. Molecular survey of Tamyb10- 1 genes and their association with grain colour and germinability in Chinese wheat and *Aegilops tauschii*. *Journal of Genetics* **94**: 453–459.
32. Edwards RA, AS Ross, DJ Mares, FW Ellison and JD Tomlinson. 1989. Enzymes from rain-damaged and laboratory-germinated wheat effects on product quality. *Journal of Cereal Science* **10**: 157–167.
33. Eshed, Y and ZB Lippman 2019. Revolutions in agriculture chart a course for targeted breeding of old and new crops. *Science* **366**: (6466).
34. Everson E. 1961. Varietal variation for dormancy in mature wheat. *Michigan Agricultural Experimental Station Quarter Bulletin*. **43**: 820–829.
35. Fang J and C Chu. 2008. Absciscic acid and the pre-harvest sprouting in cereals. *Plant Signaling and Behavior* **3**(12): 1046–1048.
36. Finch-Savage WE and G Leubner-Metzger. 2006. Seed dormancy and the control of germination. *New Phytologist* **171**(3): 501–523.
37. Flinham JE. 1993. Grain colour and sprout-resistance in wheat. In: Walker-Simmons, M.K.; Ried, J.L. (Ed.). Pre-harvest sprouting in cereals. 6th Ed. Saint Paul: American Association of Cereal Chemists p.30–36.
38. Flinham JE. 2000. Different genetic components control coat imposed and embryo-imposed dormancy in wheat. *Seed Science Research* **10**(1): 43–50.
39. Fofana B, DG Humphreys, G Rasul, S Cloutier, A Brule-Babel, S Woods, OM Lukow and DJ Somers. 2009. Mapping quantitative trait loci controlling pre-harvest sprouting resistance in a red 9 white seeded spring wheat cross. *Euphytica* **165**: 509–521.
40. Freed RD, EH Everson, K Ringlund and M Gullord. 1976. Seed coat color in wheat and the relationship to seed dormancy at maturity. *Cereal Research Communication* **4**: 147–149.
41. Gale MD and CC Ainsworth. 1984. The relationship between alpha-amylase species found in developing and germinating wheat grain. *Biochemistry and Genetics* **22**: 1031–1036.
42. Gale MD. 1989. The genetics of pre-harvest sprouting in cereals, particularly in wheat. In: Derera NF (ed)

- Pre-harvest field sprouting in cereals. *CRC, Boca Raton*.
43. Gao JB, JM Pan, QW Lu, JF Pan, MG Xu. 2006. Influencing factors of wheat sprout in field and its prevention measures. *Seed Science and Technology* **25**: 75–77.
 44. Gao X, CH Hu, HZ Li, YJ Yao, M Meng, J Dong, WC Zhao, QJ Chen, XY Li. 2013. Factors affecting pre-harvest sprouting resistance in wheat (*Triticum aestivum* L.): A Review. *Journal of Animal and Plant Science* **23**(2): 556–565.
 45. Gatford KT, RF Eastwood and GM Halloran. 2002. Germination inhibitors in bracts surrounding the grain of *Triticum tauschii*. *Functional Plant Biology* **29**: 881–890.
 46. Gautam T, Kumar K, Agarwal P, Tyagi S, Jaiswal V, Gahlaut V, Kumar S, Prasad P, Chhuneja P, Balyan HS and Gupta PK. 2021. Development of white-grain pre-harvest sprouting tolerant and pyramided protein-rich leaf rust resistant wheats using molecular breeding. DOI: <https://doi.org/10.21203/rs.3.rs-284315/v1>.
 47. Gfeller F and F Svejda. 1960. Inheritance of post-harvest seed dormancy and kernel color in spring wheat lines. *Canadian Journal of Plant Science* **40**: 1–6.
 48. Gong X, C Li, M Zhou, Y Bonnardeaux and G Yan. 2014. Seed dormancy in barley is dictated by genetics, environments and their interactions. *Euphytica* **197**: 355–368.
 49. Grant LA, JW Dick and DR Shelton 1993. Effects of drying temperature, start damage, sprouting, and additives on spaghetti quality characteristics. *Cereal Chemistry* **70**(3): 676–684.
 50. Graybosch RA, St Amand P, Bai G (2013). Evaluation of genetic markers for prediction of pre-harvest sprouting tolerance in hard white winter wheats. *Plant Breeding* **132**: 359–366.
 51. Groos C, G Gay, MR Perretant, L Gervais, M Bernard, F Dedryver and G Charmet. 2002. Study of the relationship between pre-harvest sprouting and grain color by quantitative trait loci analysis in a whitexred grain bread-wheat cross. *Theoretical and Applied Genetics* **104**(1): 39–47.
 52. Gualano NA and RL Benech-Arnold. 2009. Predicting pre-harvest sprouting susceptibility in barley: Looking for “sensitivity windows” to temperature throughout grain filling in various commercial cultivars. *Field Crops Research* **114**: 35–44.
 53. Gubler F, T Hughes, P Waterhouse and J Jacobsen. 2008. Regulation of dormancy in barley by blue light and after-ripening: Effects on abscisic acid and gibberellin metabolism. *Plant Physiology* **147**: 886–896.
 54. Gupta PK, HS Balyan, S Kumar, PK Kulwal, N Kumar, RR Mir, A Kumar, A Mohan and KV Prabhu 2008. QTL analysis and marker assisted selection for improvement in grain protein content and pre-harvest sprouting tolerance in bread wheat. in: Appels, R., Eastwood, R., Lagudah, E., Langridge, P., Mackay, M., McIntyre, L., Sharp, P. (Eds). Proceedings of 11th International Wheat Genet Symposium, Brisbane Australia, Sydney University Press p. 1–3.
 55. Gupta, PK, HS Balyan, S Sharma and R Kumar 2020. Genetics of yield, abiotic stress tolerance and biofortification in wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics*, **133**(5), 1569–1602.
 56. Hagberg S. 1960. A rapid method for determining alpha-amylase activity. *Cereal Chemistry* **37**: 218–222.
 57. Hattori T, V Vasil, L Rosenkrans, LC Hannah, DR McCarty and IK Vasil. 1992. The Viviparous-1 gene and abscisic acid activate the C1 regulatory gene for anthocyanin biosynthesis during seed maturation in maize. *Gene Development* **6**: 609–618.
 58. He J, Zhang D, X Chen, Y Li, M Hu, S Sun, Q Su, Y Su, S Li. 2021. Identification of QTLs and a Candidate Gene for Reducing Pre-Harvest Sprouting in *Aegilops tauschii*-*Triticum aestivum* Chromosome Segment Substitution Lines. *International Journal of Molecular Sciences* **22**: 3729.
 59. Hilhorst HWM. 1995. A critical update on seed dormancy. I. Primary dormancy. *Seed Science Research* **5**(1): 61–73.
 60. Himi E and K Noda. 2005. Red grain colour gene (R) of wheat is a Mybtype transcription factor. *Euphytica* **143**(3): 239–242.
 61. Himi E, DJ Mares, A Yanagisawa and K Noda. 2002. Effect of grain colour gene (R) on grain dormancy

- and sensitivity of the embryo to abscisic acid (ABA) in wheat. *Journal of Experimental Botany* **53**: 1569–1574.
62. Himi E, M Maekawa, H Miura and K Noda. 2011. Development of PCR markers for Tamyb10 related to R-1, red grain color gene in wheat. *Theoretical and Applied Genetics* **122**: 1561–1576.
63. <http://www.fao.org/faostat/en/#compare> (3/9/2019). Food and Agriculture Organization of the United Nations.
64. Ibrahim Y and B D-Appolonia. 1979. Sprouting in hard red spring wheat. *Bakers Digest* **53**(5): 17–19.
65. Imtiaz M, FC Ogbonnaya, J Oman and M Van Ginkel. 2008. Characterization of quantitative trait loci controlling genetic variation for pre-harvest sprouting in synthetic backcross-derived wheat lines. *Genetics* **178**(3): 1725–1736.
66. Jaiswal V, RR Mir, A Mohan, HS Balyan and PK Gupta. 2012. Association mapping for pre-harvest sprouting tolerance in common wheat (*Triticum aestivum* L.). *Euphytica* **188**: 89–102.
67. Ji T, B Penning and BK Baik. 2018 Pre-harvest sprouting resistance of soft winter wheat varieties and associated grain characteristics. *Journal of Cereal Science* **83**: 110–115.
68. Jiménez N, D Mares, K Mrva, C Lizana, S Contreras and AR Schwembe. 2017. Susceptibility to pre-harvest sprouting of Chilean and Australian elite cultivars of common wheat. *Crop Science* **57**: 462–474.
69. Ju L, Y Jing, P Shi, J Liu, J Chen, J Yan, J Chu, KM Chen, J Sun. 2019. JAZ proteins modulate seed germination through interaction with ABI5 in bread wheat and Arabidopsis. *New Phytologist* **223**: 246–260.
70. Kashiwakura YI, D Kobayashi, Y Jikumaru, Y Takebayashi, E Nambara, M Seo, Y Kamiya, T Kushiro and N Kawakami. 2016. Highly Sprouting-Tolerant Wheat Grain Exhibits Extreme Dormancy and Cold Imbibition-Resistant Accumulation of Abscisic Acid. *Plant Cell Physiology* **57**: 715–732.
71. Kato K, W Nakamura, T Tabiki, H Miura and S Sawada. 2001. Detection of loci controlling seed dormancy on group 4 chromosomes of wheat and comparative mapping with rice and barley genomes. *Theoretical and Applied Genetics* **102**(6): 980–985.
72. Kim KH, SH Shin, HS Lee, JY Kim, JH Son, CS Kang, JC Park and CS Park. 2014. Relationship between pre-harvest sprouting and functional markers, *TaVp-1A*, *TaSdr-B1* and *TaPHS1* in Korean wheat cultivars. *Research on Crops* **15**: 604–612.
73. Kim SY, KM Warpeha, SC Huber. 2019. The brassinosteroid receptor kinase, BRI1, plays a role in seed germination and the release of dormancy by cold stratification. *Journal of Plant Physiology* **241**, 153031.
74. King R and R Richards. 1984. Water uptake in relation to pre-harvest sprouting damage in wheat: Ear characteristics. *Australian Journal of Agricultural Research* **35**: 327–336.
75. King RW and P von Wettstein-Knowles. 2000. Epicuticular waxes and regulation of ear wetting and pre-harvest sprouting in barley and wheat. *Euphytica* **112**: 157–166.
76. King RW. 1989. Physiology of sprouting resistance. In: Derera NF (ed) Pre-harvest field sprouting in cereals. CRC Press Inc, Boca Ratan, pp 27–60.
77. Knox RE, FR Clarke, JM Clarke, SL Fox, RM DePauw and AK Singh 2012. Enhancing the identification of genetic loci and transgressive segregants for pre-harvest sprouting resistance in a durum wheat population. *Euphytica* **186**:193–206.
78. Kocheshkova AA, PY Kroupin, MS Bazhenov, GI Karlov, AA Pochtovyy and VP Upelnik et al. 2017. Pre-harvest sprouting resistance and haplotype variation of *ThVp-1* gene in the collection of wheat-wheatgrass hybrids. *PLoS ONE* **12**(11): e0188049.
79. Koornneef M, L Bentsink and H Hilhorst. 2002. Seed dormancy and germination. *Current Opinion in Plant Biology* **5**: 33–36.
80. Kottarachchi NS, N Uchino, K Kato and H Miura. 2006. Increased grain dormancy in white-grained wheat by introgression of pre-harvest sprouting tolerance QTLs. *Euphytica* **152**(3): 421–428.
81. Kucera B, MA Cohn and G Leubner-Metzger. 2005. Plant hormone interactions during seed dormancy release and germination. *Seed Science Research* **15**: 281–307.
82. Kulwal P, G Ishikawa, D Bensch, Z Feng, LX Yu, A Jadhav, S Mehete and ME Sorrells. 2012. Association

- mapping for pre-harvest sprouting resistance in white winter wheat. *Theoretical and Applied Genetics* **125**: 793–805.
83. Kulwal PL, N Kumar, A Gaur, P Khurana, JP Khurana, AK Tyagi, HS Balyan and PK Gupta. 2005. Mapping of a major QTL for pre-harvest sprouting tolerance on chromosome 3A in bread wheat. *Theoretical and Applied Genetics* **111**(6): 1052–1059.
 84. Kumar J, RR Mir, N Kumar, A Kumar, A Mohan and KV Prabhu et al. 2010. Marker-assisted selection for pre-harvest sprouting tolerance and leaf rust resistance in bread wheat. *Plant Breeding* **129**: 617–621.
 85. Lan XJ, YM Wei, DC Liu, ZH Yan and YL Zheng. 2005. Inheritance of seed dormancy in Tibetan semi-wild wheat accession Q1028. *Journal of Applied Genetics* **46**(2): 133–138.
 86. Lang J, Y Fu1, Y Zhou, M Cheng, M Deng, M Li, T Zhu, J Yang, X Guo, L Gui, L Li, Z Chen, Y Yi, L Zhang, M Hao, L Huang, C Tan, G Chen, Q Jiang, P Qi, Z Pu, J Ma, Z Liu, Y Liu, M-C Luo, Y Wei, Y Zheng, Y Wu, D Liu and J Wang. 2021. Myb10-D confers PHS-3D resistance to pre-harvest sprouting by regulating NCED in ABA biosynthesis pathway of wheat. *New Phytologist* **230**: 1940–1952.
 87. Lazo GR, S Chao, DD Hummel, H Edwards, CC Crossman, N Lui, DE Matthews, VL Carollo, DL Hane and FM You et al. 2004. Development of an expressed sequence tag (EST) resource for wheat (*Triticum aestivum* L.): EST generation, unigene analysis, probe selection and bioinformatics for a 16,000-locus bin-delineated map. *Genetics* **168**: 585–593.
 88. Lee HG and PJJJ Seo. 2015. The MYB96-HHP module integrates cold and abscisic acid signaling to activate the CBF-COR pathway in Arabidopsis. *Plant Journal* **82**: 962–977.
 89. Lei L, X Zhu, S Wang, M Zhu, BF Carver and L Yan. 2013. TaMFT-A1 is associated with seed germination sensitive to temperature in winter wheat. *PLoS One* **8**: e73330.
 90. Lin M, DD Zhang, SB Liu, GR Zhang, JM Yu, AK Fritz and GH Bai. 2016. Genome-wide association analysis on pre-harvest sprouting resistance and grain color in US winter wheat. *BMC Genomics* **17**: 794.
 91. Lin M, S Cai, S Wang, S Liu, G Zhang and G Bai. 2015. Genotyping- by-sequencing (GBS) identified SNP tightly linked to QTL for pre-harvest sprouting resistance. *Theoretical and Applied Genetics* **128**(7): 1385–1395.
 92. Lin M, S Liu, G Zhang and G Bai 2018. Effects of TaPHS1 and TaMKK3-A genes on wheat pre-harvest sprouting resistance. *Agronomy* **8**: 210.
 93. Lin R, RD Horsley and PB Schwarz. 2008. Associations between caryopsis dormancy, α -amylase activity, and pre-harvest sprouting in barley. *Journal of Cereal Science* **48**: 446–456.
 94. Liton MMA Uzzal, A Curt. Cartney M, Colin W, Hiebert, S Kumar, C Mark, Jordan, Belay T and Ayele 2021. Identification of loci for preharvest sprouting resistance in the highly dormant spring wheat RL4137. *Theoretical and Applied Genetics* **134**: 113–124
 95. Liu DC, XJ Lan, ZR Wang, YL Zheng, YH Zhou, JL Yang and C Yen. 1998. Evaluation of *Aegilops tauschii* Cosson for pre-harvest sprouting tolerance. *Genetic Resources and Crop Evolution* **45**: 495–498.
 96. Liu S, S Cai, R Graybosch, C Chen and G Bai. 2008. Quantitative trait loci for resistance to pre-harvest sprouting in US hard white winter wheat Rio Blanco. *Theoretical and Applied Genetics* **117**: 691–699.
 97. Liu S, SK Sehgal, J Li, M Lin, HN Trick, J Yu, BS Gill and G Bai. 2013a. Cloning and characterization of a critical regulator for pre-harvest sprouting in wheat. *Genetics* **195**(1): 263–273.
 98. Liu S, SK Sehgal, M Lin, J Li, HN Trick, BS Gill, G Bai. 2015. Independent mis-splicing mutations in TaPHS1 causing loss of pre-harvest sprouting (PHS) resistance during wheat domestication. *New Phytologist* **208**: 928–935.
 99. Liu X, H Zhang, Y Zhao, Z Feng, Q Li, H-Q Yang, S Luan, J Li, Z-H He. 2013c. Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in *Arabidopsis*. *Proceedings of the National Academy of Sciences*. **110**: 15485–15490.
 100. Liu Y, N Ye, R Liu, M Chen, J Zhang. 2010. H_2O_2 mediates the regulation of ABA catabolism and GA biosynthesis in Arabidopsis seed dormancy

- and germination. *Journal of Experimental Botany* **61**: 2979–2990.
101. Liu A, F Gao, Y Kanno, MC Jordan, Y Kamiya, M Seo and BT Ayele. 2013b. Regulation of wheat seed dormancy by after-ripening is mediated by specific transcriptional switches that induce changes in seed hormone metabolism and signaling. *PLoS One* **8**: e56570.
102. Lorenz K, P Roewe-Smith, K Kulp and Bates L. 1983. Pre-harvest sprouting of winter wheat. II. Amino acid composition and functionality of flour and flour fractions. *Cereal Chemistry* **60**(5): 360–366.
103. Lukow OM and W Bushuk. 1984. Influence of germination on wheat quality. 1. Functional bread making and biochemical properties. *Cereal Chemistry* **61**(4): 336–339.
104. Lunn GD, P Kettlewell, BJ Major and RK Scott. 2002. Variation in dormancy duration of the U.K. wheat cultivar Hornet due to environmental conditions during grain development. *Euphytica* **126**: 89–97.
105. Lzydorczyk C, TN Nguyen, S Jo, S Son, PA Tuan and BT Ayele. 2017. Spatiotemporal modulation of abscisic acid and gibberellin metabolism and signaling mediates the effects of suboptimal and supraoptimal temperatures on seed germination in wheat (*Triticum aestivum* L.). *Plant Cell Environment* **41**: 1022–1037.
106. Maity A and P Pramanik. 2013. Climate change and seed quality: an alarming issue in crop husbandry. *Current Science* **105**:1336–1338.
107. Marchylo BA, JE Kruger, AW Macgregor. 1983. Production of multiple forms of α -amylase in germinated, incubated, whole, de-embryonated wheat kernels. *Cereal Chemistry* **61**: 305–310.
108. Mares D and K Mrva. 2001. Mapping quantitative trait loci associated with variation in dormancy in Australian wheat. *Australian Journal Agriculture Research* **52**: 1257–1265.
109. Mares DJ and K Mrva. 2014. Wheat grain pre-harvest sprouting and late maturity alpha-amylase. *Planta* **240**: 1167–1178.
110. Martinez SA, J Godoy, M Huang, ZW Zhang, AH Carter, KA Garland Campbell and CM Steber. 2018. Genome-wide association mapping for tolerance to pre-harvest sprouting and low falling numbers in wheat. *Frontiers in Plant Science* **9**: 141.
111. McEwan JM. 1980. The sprouting reaction of stocks with single genes for red grain colour derived from hilgendorf 61 wheat. *Cereal Research Communications* **8**: 261–264.
112. McKibbin RS, MD Wilkinson, PC Bailey, JE Flintham, LM Andrew, PA Lazzeri, MD Gale, JR Lenton and MJ Holdsworth. 2002. Transcripts of Vp-1 homeologues are misspliced in modern wheat and ancestral species. *Proceedings of the National Academy of Sciences* **99**:10203–10208.
113. Metzger RJ and BA Silbaugh. 1970. Location of genes for seed coat color in hexaploid wheat (*Triticum aestivum* L.). *Crop Science* **10**: 495–496.
114. Mir RR, M Zaman-Allah, N Sreenivasulu, R Trethowan and RK Varshney. 2012. Integrated genomics, physiology and breeding approaches for improving drought tolerance in crops. *Theoretical and Applied Genetics* **125**: 625–645.
115. Mohan A, P Kulwal, R Singh, V Kumar, RR Mir, J Kumar, M Prasad, HS Balyan and PK Gupta. 2009. Genome-wide QTL analysis for pre-harvest sprouting tolerance in bread wheat. *Euphytica* **168** (3): 319–329.
116. Moot DJ and D Every. 1990. A comparison of bread baking, falling number, α -amylase assay and visual method for the assessment of pre-harvest sprouting in wheat. *Journal of Cereal Science* **11**(3): 225–234.
117. Mori M, N Uchino, M Chono, K Kato and H Miura. 2005. Mapping QTLs for grain dormancy on wheat chromosome 3A and the group 4 chromosomes, and their combined effect. *Theoretical and Applied Genetics* **110**(7): 1315–1323.
118. Munkvold JD, J Tanaka, D Benscher and ME Sorrells. 2009. Mapping quantitative trait loci for pre-harvest sprouting resistance in white wheat. *Theoretical and Applied Genetics* **119**: 1223–1235.
119. Nakamura S and T Toyama. 2001. Isolation of a VP1 homologue from wheat and analysis of its expression in embryos of dormant and non-dormant cultivars. *Journal of Experimental Botany* **52**: 875–876.

120. Nakamura S, F Abe, H Kawahigashi, K Nakazono, A Tagiri, T Matsumoto, S Utsugi, T Ogawa, H Handa, H Ishida, M Mori, K Kawaura, Y Ogihara and H Miura. 2011. A wheat homolog of MOTHER OF FT AND TFL1 acts in the regulation of germination. *Plant Cell* **23**: 3215–3229.
121. Nakamura S. 2018. Grain dormancy genes responsible for preventing pre-harvest sprouting in barley and wheat. *Breeding Science* **68**: 295–304.
122. Nonogaki H, JM Barrero and C Li 2018. Editorial: Seed Dormancy, Germination, and Pre-harvest Sprouting. *Frontiers in Plant Science* **9**: 1783. doi: 10.3389/fpls.2018.01783
123. Ogbonnaya FC, M Imtiaz and RM Depauw. 2007. Haplotype diversity of pre-harvest sprouting QTLs in wheat. *Genome* **50**: 107–118.
124. Okamoto M, A Kuwahara, M Seo, T Kushiro, T Asami, N Hirai, Y Kamiya, T Koshihara and E Nambara. 2006. Which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in Arabidopsis. *Plant Physiology* **141**: 97–107.
125. Okuyama LA, CR Riede and MM Kohli. 2020. Association between falling number and grain characteristics to evaluate pre-harvest sprouting in wheat. *Journal of Experimental Biology and Agricultural Sciences* **8**(6): 758–764.
126. Olaerts H, Vandekerckhove L and CM Courtin. 2018. A closer look at the bread making process and the quality of bread as a function of the degree of pre-harvest sprouting of wheat (*Triticum aestivum*). *Journal of Cereal Science* **80**: 188–197.
127. Osa M, K Kato, M Mori, C Shindo, A Torada and H Miura. 2003. Mapping QTLs for seed dormancy and the Vp1 homologue on chromosome 3A in wheat. *Theoretical and Applied Genetics* **106**(8): 1491–1496.
128. Patwa N and BW Penning. 2020. Environmental Impact on cereal crop grain damage from pre-harvest sprouting and late maturity alpha-amylase. R. Roy chowdhury et al. (eds.), *Sustainable Agriculture in the Era of Climate Change* 23–41.
129. Pearce S, AK Huttly, IM Prosser, YD Li, SP Vaughan, B Gallova, A Patil, JA Coghill, J Dubcovsky and P Hedden and AL Phillips. 2015. Heterologous expression and transcript analysis of gibberellin biosynthetic genes of grasses reveals novel functionality in the GA3ox family. *BMC Plant Biology* **15**: 130.
130. Penfield S and DR MacGregor. 2017. Effects of environmental variation during seed production on seed dormancy and germination. *J Exp Bot.* 2017 Feb 1. **68**(4): 819–825. doi: 10.1093/jxb/erw436. PMID: 27940467.
131. Peng J, DE Richards, NM Hartley, GP Murphy, KM Devos, JE Filntham, J Beales, LJ Fish, AJ Worland, F Pelica and D Shudhakar et al. 1999. 'Green revolution' genes encode mutant gibberellin response modulators. *Nature* **400**: 256–261.
132. Perten H. 1964. Application of the falling number method for evaluating alpha-amylase activity. *Cereal Chemistry* **41**: 127–139.
133. Probert RJ. 2000. The role of temperature in the regulation of seed dormancy and germination. *Seeds* **261**: 292.
134. Rasheed A, W Wen, F Gao, S Zhai, H Jin, J Liu, Q Guo, Y Zhang, S Dreisigacker, X Xia and Z HE. 2016. Development and validation of KASP assays for genes underpinning key economic traits in bread wheat. *Theoretical and Applied Genetics* **129**: 1843–1860.
135. Reddy LV, RJ Metzger and TM Ching. 1985. Effect of temperature on seed dormancy of wheat. *Crop Science* **25**(3): 455–458.
136. Ried JL and MK Walker-Simmons. 1990. Synthesis of abscisic acid-responsive, heat-stable proteins in embryonic axes of dormant wheat grain. *Plant Physiology* **93**: 662–667.
137. Shao M, G Bai, TW Rife, J Poland, M Lin, S Liu, H Chen, T Kumssa, A Fritz, H Trick, Y Li and G Zhang. 2018. QTL mapping of pre-harvest sprouting resistance in a white wheat cultivar Danby. *Theoretical and Applied Genetics* **131**: 1683–1697.
138. Shao M, G Bai, TW Rife, J Poland, M Lin, S Liu, H Chen, T Kumssa, A Fritz, H Trick and Y Li. 2018. QTL mapping of pre-harvest sprouting resistance in a white wheat cultivar Danby. *Theoretical and Applied Genetics* **131**: 1683–1697. <https://doi.org/10.1007/s00122-018-3107-5>.

139. Shorinola O, N Bird, J Simmonds, S Berry, T Henriksson, P Jack, P Werner, T Gerjets, D Scholefield, B Balcáková and M Valárik. 2016. The wheat Phs-A1 pre-harvest sprouting resistance locus delays the rate of seed dormancy loss and maps 0.3 cM distal to the PM19 genes in UK germplasm. *Journal of Experimental Botany* **67**: 4169–4178.
140. Shu K, H Zhang, S Wang, M Chen, Y Wu, S Tang, C Liu, Y Feng, X Cao and Q Xie. 2013. ABI4 regulates primary seed dormancy by regulating the biogenesis of abscisic acid and gibberellins in Arabidopsis. *PLoS Genetics* **9**: e1003577.
141. Silverstone AL, H-S Jung, A Dill, H Kawaide, Y Kamiya and T-P Sun. 2001. Repressing a repressor gibberellin-induced rapid reduction of the RGA protein in Arabidopsis. *The Plant Cell* **13**: 1555–1566.
142. Simsek S, JB Ohm, H Lu, M Rugg, W Berzonsky, MS Alamri and M Mergoum. 2014. Effect of pre-Harvest sprouting on physicochemical properties of starch in wheat. *Foods* **3**: 194–207.
143. Singh AK, RE Knox, JM Clarke, FR Clarke, A Singh, RM DePauw and RD Cuthbert. 2014. Genetics of pre-harvest sprouting resistance in a cross of Canadian adapted durum wheat genotypes. *Molecular Breeding* **33**: 919–929.
144. Singh R, P Hucl, M Baga and RN Chibbar. 2012. Validation of Molecular Markers for Pre-harvest Sprouting Resistance in Bread Wheat. *Cereal Research Communications* **40**(2): 194–203.
145. Sorenson B and J Wiersma. 2004. Sprout damaged wheat, crop insurance and quality concerns. *Minn. Crop News Arch.* (2004); Available online: <http://web.archive.org/web/20130205033819/http://www.extension.umn.edu/cropEnews/2004/04MNCN31.htm> (accessed on 2 April 2014).
146. Sorrells M and J Sherman. 2007. Facts: Pre-Harvest Sprouting. MAS Wheat. Available online: <https://maswheat.ucdavis.edu/> (accessed on 2 January 2019).
147. Strand E. 1989. Studies on seed dormancy in small grain species. I Barley II Wheat. *Norway Journal of Agricultural Sciences* **3**: 101–115.
148. Sun YW, HD Jones, Y Yang, S Dreisigacker, SM Li, XM Chen, PR Shewry, LQ Xia. 2012. Haplotype analysis of viviparous-1 gene in CIMMYT elite bread wheat germplasm. *Euphytica* **186**: 25–43.
149. Tai L, HJ Wang, XJ Xu, WH Sun, L Ju, WT Liu and KM Chen. 2021. Pre-harvest sprouting in cereals: genetic and biochemical mechanisms. *Journal of Experimental Botany* **72**(8): 2857–2876.
150. Tan MK, PJ Sharp, MQ Lu and N Howes. 2006. Genetics of grain dormancy in white wheat. *Australian Journal of Agricultural Research* **57**: 1157–1165.
151. Thomas H and H Ougham. 2014. The stay-green trait. *Journal of Experimental Botany* **65**: 3889–3900.
152. Torada A, M Koike, S Ikeguchi and I Tsutsui. 2008. Mapping of a major locus controlling seed dormancy using backcrossed progenies in wheat (*Triticum aestivum* L.). *Genome* **51**(6): 426–432.
153. Torada A, M Koike, T Ogawa, Y Takenouchi, K Tadamura, J Wu, T Matsumoto, K Kawaura, YA Ogiwara. 2016. Causal gene for seed dormancy on wheat chromosome 4A encodes a map kinase kinase. *Current Biology* **26**(6): 782–787.
154. Torada A, S Ikeguchi and M Koike. 2005. Mapping and validation of PCR-based markers associated with a major QTL for seed dormancy in wheat. *Euphytica* **143**: 251–255.
155. Ueguchi-Tanaka M, M Nakajima, E Katoh et al. 2007. Molecular interactions of a soluble gibberellin receptor, GID1, with a rice DELLA protein, SLR1, and gibberellin. *The Plant Cell* **19**: 2140–2155.
156. Van De Velde K, PM Chandler, D Van Der Straeten and A Rohde. 2017. Differential coupling of gibberellin responses by Rht-B1c suppressor alleles and Rht-B1b in wheat highlights a unique role for the DELLA N-terminus in dormancy. *Journal of Experimental Botany* **68**: 443–455.
157. Vetch JM, RN Stougaard, JM Martin and MJ Giroux. 2019. Review: revealing the genetic mechanisms of pre-harvest sprouting in hexaploid wheat (*Triticum aestivum* L.). *Plant Science* **281**: 180–185.
158. Volis S. 2016. Seed heteromorphism in *Triticum dicoccoides*: association between seed positions within a dispersal unit and dormancy. *Oecologia* **181**: 401–412.

159. Wahl TI and AD O'Rourke. 1994. The economics of sprout damage in wheat. *Agribusiness* **10**(1): 27–41.
160. Wang X, H Liu, G Liu, M-S Mia, Kadambot-HM Siddique and G Yan. 2019. Phenotypic and genotypic characterization of near-isogenic lines targeting a major 4BL QTL responsible for pre-harvest sprouting in wheat. *BMC Plant Biology* **19**: 348 <https://doi.org/10.1186/s12870-019-1961-1>
161. Wang X, H Liu, HM Kadambot-Siddique and G Yan. 2021. Transcriptomic profiling of wheat nearisogenic lines reveals candidate genes on chromosome 3A for pre-harvest sprouting resistance. *BMC Plant Biology* **21**: 53 <https://doi.org/10.1186/s12870-021-02824-x>
162. Warner RL, DA Kudrna, SC Spaeth and SS Jones. 2000. Dormancy in white-grain mutations of Chinese Spring wheat (*Triticum aestivum* L.). *Seed Science Research* **10**(1): 51–60.
163. Wu J and BF Carver. 1999. Sprout damage and pre-harvest sprout resistance in hard white winter wheat. *Crop Science* **39**(2): 441–447.
164. Xia LQ, MW Ganai, PR Shewry, ZH He, Y Yang and MS Roder. 2008. Exploiting the diversity of Viviparous-1 gene associated with pre-harvest sprouting tolerance in European wheat varieties. *Euphytica* **159**: 411–417.
165. Xiao C, Y Liu, W Chen, J Yang, M Cheng, C Watt, J Cheng, Z Wang, Z Tan M Li and J Wang. 2021. Characterization and expression quantitative trait loci analysis of TaABI4, a pre-harvest sprouting related gene in wheat. *Seed Science Research* 1–11. <https://doi.org/10.1017/S0960258521000015>.
166. Xiao SH, XY Zhang, CS Yan and H Lin. 2002. Germplasm improvement for pre-harvest sprouting resistance in Chinese white-grained wheat: an overview of the current strategy. *Euphytica* **126**: 35–38.
167. Yanagisawa A, T Nishimura, Y Amano, A Torada and S Shibata. 2005. Development of winter wheat with excellent resistance to pre-harvest sprouting and rain damage. *Euphytica* **143**(3): 313–318.
168. Yang J, C Tan, J Lang, H Tang, M Hao, Z Tan, H Yu, Y Zhou, ZH Liu, ML Li and Y Zhou. 2019. Identification of qPHS.sicau-1B and qPHS.sicau-3D from synthetic wheat for pre-harvest sprouting resistance wheat improvement. *Molecular Breeding* **39**: 132–143.
169. Yang J, Y Liu, Z Pu, L Zhang, Z Yuan, G Chen, Y Wei, Y Zheng, D Liu and JR Wang. 2014. Molecular characterization of high pI α -amylase and its expression QTL analysis in synthetic wheat RILs. *Molecular Breeding* **34**: 1075–1085.
170. Yang JH, YX Yu, JS Cheng, XL Tan and WP Shen. 2011. Study on the pre harvest sprouting tolerance in *Triticum aestivum* ssp. *Yunnanense* King. *Journal of Triticeae Crops* **31**: 747–752 (in Chinese).
171. Yang Y, CL Zhang, SX Liu, YQ Sun, JY Meng and LQ Xia. 2014. Characterization of the rich haplotypes of viviparous-1A in Chinese wheats and development of a novel sequence-tagged site marker for pre-harvest sprouting resistance. *Molecular Breeding* **33**: 75–88.
172. Yang Y, XM Chen, ZH He, M Roder and LQ Xia. 2009. Distribution of Vp-1 alleles in Chinese white-grained landraces, historical and current wheat cultivars. *Cereal Research Communication* **37**: 169–177.
173. Yang Y, YZ Ma, ZS Xu, XM Chen, ZH He, Z Yu, M Wilkinson, HD Jones, PR Shewry and LQ Xia. 2007. Isolation and characterization of Viviparous-1 genes in wheat cultivars with distinct ABA sensitivity and pre-harvest sprouting tolerance. *Journal of Experimental Botany* **58**: 2863–2871.
174. Yu-ichi K, Daisuke K, Yusuke J, Yumiko T, Eiji N, Mitsunori S, Yuji K, Tetsuo K, Naoto K. 2016. Highly sprouting-tolerant wheat grain exhibits extreme dormancy and cold imbibition-resistant accumulation of abscisic acid. *Plant and Cell Physiology*. **57** (4): 715–732.
175. Yu M, GY Chen, LQ Zhang, YX Liu, DC Liu, JR Wang, ZE Pu, L Zhang, XJ Lan, YM Wei and CJ Liu. 2014. QTL Mapping for important agronomic traits in synthetic hexaploid wheat derived from *Aegilops tauschii* ssp. *tauschii*. *Journal of Integrated Agriculture* **13**: 1835–1844.
176. Yu N, R Laurenz, L Siler, P Ng, E Souza and J Lewis. 2015. Evaluation of alpha-amylase activity and falling number around maturity for soft white and soft red wheat varieties in michigan. *Cereal Research Communications* **43**(4): 672–681.

177. Zhang D, J He, L Huang, C Zhang, Y Zhou, Y Su and S Li. 2017a. An advanced backcross population through synthetic octaploid wheat as a “bridge”: development and QTL detection for seed dormancy. *Frontiers in Plant Science* **8**: 2123.
178. Zhang HF and RZC Liu. 1989. Studies on pre-harvest sprouting resistance in winter wheat and its determination. *Acta Agronomica Sinica*. **15**: 116–122.
179. Zhang Q and C Li. 2017c. Comparisons of copy number, genomic structure, and conserved motifs for α -amylase genes from barley, rice and wheat. *Frontiers in Plant Science* **8**: 1727.
180. Zhang Y, X Miao, X Xia and Z He. 2014. Cloning of seed dormancy genes (*TaSdr*) associated with tolerance to pre-harvest sprouting in common wheat and development of a functional marker. *Theoretical and Applied Genetics* **127**(4): 855–866.
181. Zhang Y, X Xia and Z He. 2017b. The seed dormancy allele *TaSdr-A1a* associated with pre-harvest sprouting tolerance is mainly present in Chinese wheat landraces. *Theoretical and Applied Genetics* **130**: 81–89.
182. Zhou SH, L Fu, QH Wu, JJ Chen, YX Chen, JZ Xie, ZZ Wang, GX Wang, DY Zhang, Y Liang and Y Zhang. 2017. QTL mapping revealed *TaVp-1A* conferred pre-harvest sprouting resistance in wheat population Yanda 1817xBeinong 6. *Journal of Integrated Agriculture* **16**: 435–444.
183. Zhou Y, H Tang, MP Cheng, KO Dankwa, ZX Chen, ZY Li, S Gao, YX Liu, QT Jiang, XJ Lan, ZE Pu, YM Wei, YL Zheng, LT Hickey and JR Wang. 2017. Genome-wide association study for pre-harvest sprouting resistance in a large germplasm collection of Chinese wheat landraces. *Frontiers Plant Science* **8**: 401.
184. Zhu ZL, B Tian, B Liu, QG Xie and JC Tian. 2010. Quantitative trait loci analysis for pre-harvest sprouting using intact spikes in wheat (*Triticum aestivum* L.). *Shandong Agriculture Science* **6**: 19–23.