

## Influence of propylene glycol on conception rate in cows

ZEKI ERISIR<sup>1</sup> and ATILLA YILDIZ<sup>2</sup>

University of Firat, Elazig 23119 Turkey

Received: 23 July 2015; Accepted: 21 August 2015

### ABSTRACT

The aim of the study was to evaluate effect of propylene glycol (PG) drenched during the first 7 days after fixed-time artificial insemination (FTAI) on plasma progesterone (P<sub>4</sub>) and insulin levels, and on pregnancy rate in Holstein cows in summer. Ovsynch program was performed in 40 healthy Holstein cows. Cows were distributed randomly into two groups as propylene glycol (group PG) (20) and control group (group C) (20). Group PG and group C were daily administered with 500 ml of PG and water, respectively for 7 days from day 0 to 7 post insemination. Pregnancy diagnosis was carried out 60 days after insemination via rectal examination. Cows exposed to moderate heat stress due to temperature-humidity index ( $74.4 \pm 1.9$  mean THI). Conception rate for cows in group PG increased 2.9 fold when was compared with group C. On day 7, the insulin levels were higher in group PG than group C. It was concluded that PG administered once daily from day 0 to 7 after FTAI increased insulin and P<sub>4</sub> concentrations and the conception rate in Holstein cows. The rise in insulin and P<sub>4</sub> levels depending on the drenched PG are likely related with the increases of conception rate.

**Key words:** Fertility, Heat stress, Insulin, Progesterone, Propylene glycol

Fertility in lactating cows is decreased during the hot seasons of the year. Summer-calved cows had greater number of services per conception and lower conception rates than cows calved in other seasons (Ghavi Hosseini-Zadeh *et al.* 2013). The lactating cow at an air temperature of 30°C is at risk of infertility because of heat stress (Peter 2007). Heat stress has two major effects on reproduction. First, heat stressed cows demonstrate estrus less highly than other cows. Secondly, conception rate is decreased during heat stress (Hansen 2007). Thatcher (1974) showed that conception rates are depressed at environmental temperatures above the critical temperature of 21 °C. Heat stress affects the development and the function of the corpus luteum and causes a decrease in P<sub>4</sub> levels (Ullah *et al.* 1996). Therefore, the FTAI and various hormonal protocols to improve the use of artificial insemination are widely used in the dairy farms to prevent the negative effects on reproductive performance of high temperature environment. However, the ability of these strategies to improve the fertility is limited. Use of FTAI protocols can eliminate problems of estrus detection that heat stress caused. But this was not adequate to reinstate herd pregnancy rates to a level seen in cool weather due to the serious outcomes of heat stress for embryogenesis (Hansen and Arechiga 1999). Hence, alternative techniques to FTAI protocols that improve pregnancy rates for a systematic breeding system

should be developed. Leroy *et al.* (2006) showed that cleavage rate and blastocyst development were severely reduced *in vitro* in a low glucose environment vs. a physiologically normal glucose environment. The use of glycogenic supplements during the estrous cycle increased the synthesis of P<sub>4</sub>. Oral drenching of PG raises insulin levels which affect the developing corpus luteum directly. (Miyoshi *et al.* 2001). Insulin is required for maintenance of P<sub>4</sub> synthesis, which eases lipoprotein using in bovine luteal cells (Poff *et al.* 1998). Luteal function and conception rate improves in cows treated with insulin subcutaneously (Selvaraju *et al.* 2002). Propylene glycol is a substance used in prevention and treatment of ketosis in dairy cows (Rakkwamsuk *et al.* 2010). There are studies where oral drenching of PG has an increased conception rate in cows (McArt *et al.* 2012, Slobodanka *et al.* 2012). However, other groups have reported no differences in the success of pregnancy at first artificial insemination for cows drenched with PG (Chagas *et al.* 2007, Miyoshi *et al.* 2001). Hidalgo *et al.* (2004), using PG for embryo transfer, found that the group of cows treated had higher 60-day pregnancy rates than the control group. Propylene glycol improved *in vitro* embryonic development rate (Ponsart *et al.* 2014). However, the results of using PG are controversial for the success of conceiving and need more investigation. Thus, it can be speculated that PG drenched in the first seven days post insemination are useful for the establishment of pregnancy by increasing plasma P<sub>4</sub> and insulin concentrations in dairy cows during the hot season. Therefore, the present investigation was undertaken to evaluate the effect of PG

Present address: <sup>1</sup>Faculty of Veterinary Medicine (zerisir@hotmail.com), <sup>2</sup>Veterinary Department (ayildiz1@firat.edu.tr), Sivrice Vocational School.

drenched during the first seven days after FTAI on plasma  $P_4$  and insulin levels, and on conception rate in Holstein cows during summer.

## MATERIALS AND METHODS

**Study area, animals and treatments:** The study was conducted on 40 healthy Holstein cows at a private dairy farm located in Elazig province of Turkey in summer season. Elazig region is between the latitude of 38° 40'N and longitude of 39° 13'E, at an altitude of 1,093 m. Ambient temperature (AT) (°C) and relative humidity (RH) were obtained from reports prepared at Turkish State Meteorological Service. The mean temperature was  $29.4 \pm 0.5^\circ\text{C}$  with in the period of the experiment. The relative humidity is  $29.7 \pm 2.3\%$ . The temperature and humidity index (THI) was obtained using the formula:  $\text{THI} = (0.8 \times \text{AT} + (\text{RH}\%/100) \times (\text{AT} - 14.4) + 46.4)$  (Thom 1959). Cows with detectable puerperal complications following calving were not included in the experiment. The body condition score (point scale from 1 to 5) were 2.5–3.0 units. The mean body weight was  $517.0 \pm 34.57$  kg. The parity of cows ranged between 2 and 3. During the experiment, all the animals were kept under the similar feeding and managemental conditions. Average daily milk production for the farm was between 20 to 23 kg/cow during the study period. Lactating cows were milked twice daily. All cows were examined by rectal palpation and vaginoscopic examination at week 6 post-partum. There were no clinically detectable disorders in their reproductive tracts. On 45<sup>th</sup> day of lactation, Ovsynch protocol was done with the administration of GnRH (Busereline acetate). The administration of PGF2 $\alpha$  analogue D- Cloprostenol (25 mg, Dinoprost tromethamine) was done on day 8 after first administration and second GnRH administration was on day 10 after first administration. Cows were artificially inseminated after 18h following second GnRH administration. At insemination (day 0), cows were distributed randomly into two groups as group PG and control group (group C). The randomization was stratified by parity, body weight and BCS. Propylene glycol (group PG) and water (group C) were supplied via drench in a volume of 500 ml once daily from day 0 to 7 post insemination. The diagnosis of pregnancy was done on 60<sup>th</sup> day after insemination via rectal examination.

**Sample collection and biochemical assays:** In all animals, blood samples to be assayed for  $P_4$  and insulin were obtained from the jugular vein using heparinized vacutainer tubes just before the application on day 0 (oestrus) and one hours after the administration of PG or water on day 7. Samples were later centrifuged at  $1,500 \times g$  for 15 min for plasma separation, which was then stored at  $-20^\circ\text{C}$  until being analyzed. Insulin concentrations were measured by RIA in solid phase, using the insulin kit, having a 0.05 ng/ml sensibility and a 9.5% intra assay variation coefficient. Plasma  $P_4$  concentration was determined using a commercial solid-phase RIA kit. The sensitivity of the assay was 0.1 ng/ml and the intra assay CV was 5.6%.

**Statistical analysis:** The conception rates of groups were compared with the Odds ratio (OR). The reference category had an OR = 1. An adjusted OR > 1.0 indicates that the probability of the increased conception rate because of the treatment was, compared with cows in the reference category. Insulin and  $P_4$  concentrations were compared using a *t*-test to determine the effects of treatment. Results were considered statistically and declared significant at  $P < 0.05$ .

## RESULTS AND DISCUSSION

Mean values of THI were equal to  $74.4 \pm 1.9$ . Cows in the study exposed to moderate heat stress according to the temperature-humidity index ( $74.4 \pm 1.9$  mean THI).

Plasma concentrations of insulin and  $P_4$  on Day 0 and Day 7 are presented in Table 1. On day 7, the insulin levels were higher in group PG ( $P < 0.05$ ) than group C. The insulin levels on Day 7 for cows in group PG were statistically higher compared to ones on day 0 (before treatment) ( $P < 0.05$ ). In the current study, we obtained that drenching of PG increased insulin plasma concentrations, which is in accordance with other previous studies that reported an increase in the concentrations of insulin following administration of PG (Christensen *et al.* 1997, Miyoshi *et al.* 2001). Propylene glycol may stimulate insulin secretion directly (Studer *et al.* 1993). Likewise, it is possible that the PG stimulates pancreatic insulin secretion indirectly by PG metabolizes to propionate (Webb *et al.* 1999).

The average  $P_4$  concentration was higher ( $P < 0.05$ ) in group PG when compared with group C on Day 7 after insemination. Plasma  $P_4$  concentrations increased significantly on day 7 after PG drenching. This was similar to the findings of Miyoshi *et al.* (2001) who found a significant rise in milk  $P_4$ . Elevation of insulin by PG drenching affects the developing corpus luteum directly (Miyoshi *et al.* 2001). Insulin is a requisite to maintain  $P_4$  synthesis and ease lipoprotein using in bovine luteal cells (Poff *et al.* 1998). Cows treated with insulin subcutaneously display rises in  $P_4$  concentrations (Selvaraju *et al.* 2002). Likewise, insulin-like growth factor-I (IGF-I) stimulates  $P_4$  production by cultured bovine luteal cells (Chakravofy *et al.* 1993). Also, drenching of PG is known to rise IGF-I

Table 1. Mean levels and standard deviations ( $\pm$ SD) of plasma progesterone ( $P_4$ ) and insulin concentrations for cows in group C and group PG

| Parameters             | Days | Group C<br>n=20      | Group PG<br>n=20      |
|------------------------|------|----------------------|-----------------------|
| Progesteron<br>(ng/mL) | 0    | $0.67 \pm 0.19^A$    | $0.69 \pm 0.17^A$     |
|                        | 7    | $2.89 \pm 0.37^{Ba}$ | $3.38 \pm 0.41^{Bb}$  |
| Insulin<br>(ng/mL)     | 0    | $15.69 \pm 0.43$     | $15.63 \pm 0.51^A$    |
|                        | 7    | $14.78 \pm 0.37^a$   | $19.87 \pm 0.59^{Bb}$ |

a,bValues with different superscript letters within a row differ significantly at  $P < 0.05$ . A,BValues with different superscript letters within a column differ significantly at  $P < 0.05$ .

Table 2. The conception rates of cows in group C and group PG

| Groups   | n  | Conception rate<br>(%,n/n) | Odds ratio | 95% Confidence interval | P values |
|----------|----|----------------------------|------------|-------------------------|----------|
| Group C  | 20 | 30.0<br>(6)                | Referent   |                         |          |
| Group PG | 20 | 55.0<br>(11)               | 2.9        | 0.78 to 10.47           | P=0.11   |

(Grummer *et al.* 1994). Change in insulin is closely associated to IGF-I levels (O'Callaghan and Boland 1999). In a direct way, IGF-I stimulates corpus luteum growth and steroidogenesis (Alvarez *et al.* 2000). That's why,  $P_4$  levels could raise in reply to insulin and IGF-I, whose levels also would be more elevated after a PG drenching.

Mean conception rate at first insemination was 55.0% (11/20) in group PG and 30.0% (6/20) in group C (Table 2). Cows in group PG had 2.9 times more conception rate than control ones. Supplementation of PG from day 0 to day 7 improved conception rate at a rate of 25.0%. The present finding is in accordance with León *et al.* (2010) who also recorded higher conception rate in Holstein cows cured with a glycogenic diet (glycerol). Hidalgo *et al.* (2004) observed that the group of cows treated using mono propylene glycol for embryo transfer had higher 60-day conception rates than group C. The mechanism through which PG act on conception rate has been completely understood; nevertheless, it could be stated by the effects of insulin and  $P_4$  on early embryo development (Block *et al.* 2011). Due to the fact that adding insulin to a culture media promotes embryo cell mitosis and the ratio of embryos reaching the blastocyst stage (Augustin *et al.* 2003). Therefore, concentrations of tau interferon (IFN- $\tau$ ) were higher in cows fed regimes prepared to create elevated insulin levels (Mann *et al.* 2003). Likewise, the increased  $P_4$  levels during the early stages of pregnancy may contribute to a progress of embryonic development causing to a raised level of IFN- $\tau$  production and a related rise in conception rate (Morris and Diskin 2008). The inadequate IFN- $\tau$  synthesis is one of the reasons of conception failure. (Diskin and Morris 2008). Therefore, PG improves embryo capacity to prompt maternal recognition of pregnancy, which influences embryo health and ability to establish pregnancy and results in an increase in conception rate. In addition,  $P_4$  has an important role in regulating changes in the uterine environment for supporting early embryonic growth and development. Early  $P_4$  stimulation changes endometrial secretions and proceeds to development of conceptus (Butler 2003).

In conclusion, the results of the present study indicate that drenching of PG at a dose rate of 500 ml/day from days 0 to 7 after insemination increases insulin and  $P_4$  concentrations and the conception rate in Holstein cows. The rise in insulin and  $P_4$  concentrations after drenching PG leads to increase in the conception rate. This application can provide a contribution to improve low conception rates

due to moderate heat stress occurred especially in the summer.

## REFERENCES

- Alvarez P, Spicer L J, Chase C C Jr, Payton M E, Hamilton T D, Stewart R E, Hammond A C, Olson T A and Wettemann R P. 2000. Ovarian and endocrine characteristics during an estrous cycle in Angus, Brahman, and Senepol cows in a subtropical environment. *Journal of Animal Science* **78** (5): 1291–1302.
- Augustin R, Pocar P, Wrenzycki C, Niemann H and Fischer B. 2003. Mitogenic and antiapoptotic activity of insulin bovine embryos produced in vitro. *Reproduction* **126** (1): 91–99.
- Block J, Hansen PJ, Loureiro B and Bonilla L. 2011. Improving post-transfer survival of bovine embryos produced in vitro: Actions of insulin-like growth factor-1, colony stimulating factor-2 and hyaluronan. *Theriogenology* **76** (9): 1602–09.
- Butler W R. 2003. Energy balance relationships with follicular development, ovulation and fertility in postpartum dairy cows. *Livestock Production Science* **83** (2–3): 211–18.
- Chagas L M, Gore P S J, Meier S, Macdonald K A and Verkerk G A. 2007. Effect of monopropylene glycol on luteinizing hormone, metabolites, and postpartum anovulatory intervals in primiparous dairy cows. *Journal of Dairy Science* **90** (3): 1168–75.
- Chakravofty A, Joslyn M I and Davis J S. 1993. Characterization of insulin and insulin-like growth factor I actions in the bovine luteal cell: regulation of receptor tyrosine kinase activity phosphatidylinositol-3-kinase, and deoxyribonucleic acid synthesis. *Endocrinology* **133** (3):1331–40.
- Christensen J O, Grummer R R, Rasmussen F E and Bertics S J. 1997. Effect of method of delivery of propylene glycol on plasma metabolites of feed-restricted cattle. *Journal of Dairy Science* **80** (3): 563–68.
- Diskin M G and Morris D G. 2008. Embryonic and early foetal losses in cattle and other ruminants. *Reproduction in Domestic Animals* **43** (Suppl 2): 260–67.
- Ghavi Hossein-Zadeh N, Mohit A and Azad N. 2013. Effect of temperature-humidity index on productive and reproductive performances of Iranian Holstein cows. *Iranian Journal of Veterinary Research* **14** (2): 106–12.
- Grummer R R, Winkler J C, Bertics S J and Studer V A. 1994. Effect of propylene glycol dosage during feed restriction on metabolites in blood of prepartum Holstein heifers. *Journal of Dairy Science* **77** (12): 3618–23.
- Hansen P J. 2007. Hidden Factors Affecting Fertility. WCDS Advances in Dairy Technology 19: 339–349 (<http://www.wcds.ca/proc/2007/Manuscripts/Peter2.pdf>)
- Hansen P J and Are'chiga C F 1999. Strategies for managing reproduction in the heat-stressed dairy cow. *Journal of Animal Science* **77** (Suppl 2): 36–50.
- Hidalgo C O, Gomez E, Prieto L, Duque P, Goyache F, Fernandez L, Fernandez I, Facal N and Diez C. 2004. Pregnancy rates and metabolic profiles in cattle treated with propylene glycol prior to embryo transfer. *Theriogenology* **62** (3–4): 664–76.
- León A O, Ceróna J H and Gutiérreza C G. 2010. Holstein Glycerol supplementation after artificial insemination increases conception rate in Holstein cows. *Revista Mexicana de Ciencias Pecuarias* **48** (1): 69–74.
- Leroy J L M R, Vanholder T, Opsomer G, Van Soom A and de Kruif A. 2006. The in vitro development of bovine oocytes after maturation in glucose and  $\beta$ -hydroxybutyrate concentrations associated with negative energy balance in dairy

- cows. *Reproduction in Domestic Animals* **41** (2): 119–23.
- Mann GE, Green MP, Sinclair KD, Demmers KJ, Fray MD, Gutierrez CG, Garnsworthy PC and Webb R. 2003. Effects of circulating progesterone and insulin on early embryo development in beef heifers. *Animal Reproduction Science* **79** (1–2): 71–79.
- McArt J A A, Nydam D V and Oetzel G R. 2012. A field trial on the effect of propylene glycol on displaced abomasum, removal from herd, and reproduction in fresh cows diagnosed with subclinical ketosis. *Journal of Dairy Science* **95** (5): 2505–12.
- Miyoshi S, Pate J L and Palmquist D L. 2001. Effects of propylene glycol drenching on energy balance, plasma glucose, plasma insulin, ovarian function and conception in dairy cows. *Animal Reproduction Science* **68** (1–2): 29–43.
- Morris D and Diskin M. 2008. Effect of progesterone on embryo survival. *Animal* **2** (8): 1112–19.
- O’Callaghan D and Boland M P. 1999. Nutritional effects on ovulation, embryo development and the establishment of pregnancy in ruminants. *Animal Science* **68** (2): 299–314.
- Poff J P, Fairchild D L and Condon W A. 1998. Effects of antibiotics and medium supplements on steroidogenesis in cultured cow luteal cells. *Journal of Reproduction and Fertility* **82** (1): 135–43.
- Ponsart C, Gamarra G, Lacaze S and Ponter A A. 2014. Nutritional status of donor cows: insulin related strategies to enhance embryo development. *Animal Reproduction* **11** (3): 195–98.
- Rakkwamsuk T, Rungruang S, Choothesa A and Wensing T. 2010. Performance of periparturient dairy cows fed either by alfalfa hay or peanut hay in total mixed ration: A field trial in Thailand. *African Journal of Agricultural Research* **5** (12): 1430–38.
- Selvaraju S, Agarwal S K, Karche S D, Srivastava S K, Majumdar A C and Shanker U. 2002. Fertility responses and hormonal profiles in repeat breeding cows treated with insulin. *Animal Reproduction Science* **73** (3–4): 141–49.
- Slobodanka V M, Pavlović V, Gvozdić D, Jovičević M, Đukić M and Stepanović P. 2012. Propylene glycol energy supplementation during periparturient period in dairy cows and reproduction efficiency parameters. *Acta Veterinaria (Beograd)* **62** (1–2): 249–60.
- Studer V A, Grummer R R, Bertics S J and Reynolds C K. 1993. Effect of prepartum propylene glycol administration on periparturient fatty liver in dairy cows. *Journal of Dairy Science* **76** (10): 2931–39.
- Thatcher W W. 1974. Effects of season, climate and temperature on reproduction and lactation. *Journal of Dairy Science* **57** (3): 360–68.
- Thom EC. 1959. The discomfort index. *Weatherwise* **12** (2): 57–59.
- Ullah G, Fuquay J W, Keawkhong T, Clark B L, Pogue D E and Murphey E J. 1996. Effect of gonadotropin-releasing hormone at estrus on subsequent luteal function and fertility in lactating Holsteins during heat stress. *Journal of Dairy Science* **79** (11): 1950–53.
- Webb R, Campbell B K, Garverick H A, Gong J G, Gutierrez C G and Armstrong D G. 1999. Molecular mechanism regulating follicular recruitment and selection. *Journal of Reproduction and Fertility. Supplement* **54**: 33–48.