

Contents

Effects of biological and chemical sources of fertilizers on sunflower yield and its components	63
<i>Maryam Zamanian and Mohammad Yazdandoost</i>	
Targeted yield based integrated nutrient management of cabbage in torripsamment	75
<i>S.R. Yadav, B.S. Meena, Anirudh Choudhari and Sunil Kumar</i>	
Boron management in green gram (<i>Vigna radiata</i> L. Wilczek) under custard apple (<i>Annona squamosa</i> L.) based agri-horti system in semi-arid region	79
<i>Raj Kumar, J.K. Singh, Alok Kumar Singh, Srishti Dipriya Minz and Nasam Midhun Kumar</i>	
Conjunctive application of organic and inorganic fertilizers to improve soil fertility and wheat (<i>Triticum aestivum</i> L.) productivity in semi-arid region	85
<i>Narendar Kumar Meena, A.M. Latore, M.K. Singh, Prashant Sharma and Kamlesh Verma</i>	
Effect of different grazing management practices on soil health in <i>Lasiurus indicus</i> grasslands of arid western plain	93
<i>Nav Raten Panwar, Sharmila Roy, Mahesh Kumar and C.B. Pandey</i>	
Techno-economic analysis of inclined solar dryer for carrot (<i>Daucus carota</i> L.) drying	99
<i>Surendra Poonia, A.K. Singh and Dilip Jain</i>	
Genetic diversity of groundnut (<i>Arachis hypogaea</i> L.) revealed by RAPD and ISSR markers	109
<i>Papi Reddy, Pritesh Sabara, Shital M. Padhiyar, Kulkarni, G.U., J.V. Kheni and Rukam S. Tomar</i>	
Effect of hormonal application on physio-biochemical characters to improve seed yield of Sewan grass (<i>Lasiurus indicus</i> Henrard)	117
<i>Maharaj Singh, K. Venkatesan and N.K. Sinha</i>	
Pollen biology of <i>Grewia optiva</i> drummond genotypes: An important agroforestry tree of north western Himalayas	123
<i>Saresh N.V., Archana Verma, Asu Singh Godara, Dharmendera Meena and Arjun Lal Bijarnia</i>	
Genetic variability studies in fennel (<i>Foeniculum vulgare</i>) in arid western Rajasthan	135
<i>Mahipal Jat and Santosh Choudhary</i>	
Assessment of pattern and sustainability of livestock diversification in Odisha	141
<i>Urmi Pattanayak and Kalu Naik</i>	
Short Communication	
Should we be concerned about climate change?	149
<i>Bachir Khezzani</i>	
Modification of traditional tractor-drawn seed drill for arid region crops	151
<i>A.K. Singh, Dinesh Mishra, Prem Veer Gautam, Surendra Poonia and Dilip Jain</i>	



Boron Management in Green gram (*Vigna radiata* L. Wilczek) under Custard Apple (*Annona squamosa* L.) based Agri-horti System in Semi-arid Region

Raj Kumar, J.K. Singh, Alok Kumar Singh^{1*}, Srishti Dipriya Minz and Nasam Midhun Kumar¹

Institute of Agricultural Sciences, Banaras Hindu University, Varanasi 221 005, India

¹Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Solan 173 230, India

Received: December 2021

Abstract: An experiment was conducted at the Agricultural Research Farm, RGSC, BHU, Barkachha, Mirzapur (UP) on sandy clay loam soil during kharif 2016-17 to evaluate the effect of three levels of Boron (0.75, 1.50 and 2.25 kg ha⁻¹) and foliar spray of two levels of Solubor (0.2 and 0.3%) on green gram (*Vigna radiata* L. Wilczek) under custard apple (*Annona squamosa* L.) based agri-horti system. The growth and yield parameters were significantly affected by different levels of Boron and Solubor. Data revealed that application of Boron @ 2.25 kg ha⁻¹ gave maximum values of growth parameters, yield attributes, grain and stover yield, harvest index as well as content and removal of N and S by green gram over rest of the levels. Similarly, the application of solubor @ 0.3% ha⁻¹ exhibited higher in all growth parameters, yield attributes, yield, N and S content and its removal over rest of the levels. Moreover, the maximum net return (Rs. 60505 ha⁻¹) was also observed under the application of boron (2.25 kg ha⁻¹) along with solubor (0.3% ha⁻¹).

Key words: Agri-horti system, green gram, custard apple, boron, yield.

The agroforestry system creates a more integrated, diverse, productive, profitable, healthy and sustainable land-use system (Sharma *et al.* 2017b) in addition to the reclamation of degraded, wastelands (Sharma *et al.*, 2017a), soil water conservation (Singh *et al.*, 2020) and climate change adaptation and mitigation (Verma *et al.*, 2021). Agri-horticultural system is a modified traditional cropping system of India that utilizes the maximum capability of land as well as growing season and simultaneously enhances the productivity of land. Pulse crop along with Guava, Bael, Aonla, Custard apple and Karonda etc. are more commonly grown in Vindhyan region under the agri-horticultural system (Minz *et al.*, 2021). India is the largest producer and consumer of pulses in the world accounting for 25% of total world production and 27% of world's consumption. Green gram [*Vigna radiata* (L.) Wilczek] commonly known as "mung bean" is one of the important short duration pulse crops grown in India. It ranks third among all the pulses grown in India after chickpea and pigeonpea (Tamang *et al.*, 2015). The seeds of green gram are highly nutritious with protein, carbohydrates, minerals and vitamins. Proteins are rich in lysine, leucine and

threonine but poor in methionine, tryptophan, tyrosine. Every 100 g of edible portion of green gram seed contains 75 mg calcium, 4.5 mg phosphorus, 24.5 g protein and 348 kilo calories energy (Meena *et al.*, 2013). Boron is one of the most widely applied micronutrients. Solubor tetra borate (Na₂B₄O₇·5H₂O) is the most common B source containing 20% B. Solubor is a highly concentrated, soluble B source that can be used as foliar in form of liquid or dust. It is also used in liquid and suspension fertilizers. Solubor is preferred to borax because it dissolves more readily (Alloway, 2009). Solubor is a specially developed product for speedy and economic correction of boron deficiency in fruit trees, vegetables and other crops (Havlin *et al.*, 2014). Further, boron is an essential plant nutrient and primary function of boron is in plant cell wall structural integrity. Boron also provide cross links between cell wall polysaccharides for providing structure to the cell wall-important for cell expansion, regulation of hydrogen ion (H⁺) transport retention of cellular Ca⁺², and control of lignin production following cell expansion. Considering the above facts in view the present investigation is carried out to find the effect of Boron management in green gram (*Vigna radiata* L. Wilczek) under custard apple (*Annona squamosa* L.) based agri-horti system.

*E-mail: alokkj63@gmail.com

Materials and Methods

The field experiment was carried out during kharif season of 2016-17 at the Rajiv Gandhi South Campus Agricultural Research Farm, Barkachha (BHU), Mirzapur, located in the Vindhyan region of Mirzapur (25°10'' latitude, 82°37'' longitude and 427 meters above sea level) UP, India. Mirzapur falls to the sub-humid climate in a semi-arid belt. The normal period for the onset of monsoon in this region is June's third week and lasts until the end of September or sometimes extends to the first week of October. From December to mid-February, winter showers are often experienced. Overall, however, March to May is dry. On average, the major fraction (75%) of the total annual rainfall is received from June to September. The monthly mean maximum and minimum temperature during the period of experiment, ranges from 25.3°C to 36.7°C and 10.5°C to 27.3°C, respectively.

The soil of the experimental field was sandy clay loam in texture. The experimental soil was low in available nitrogen (147.7 kg ha⁻¹), organic carbon (0.36%), as well as available sulphur (12.2 kg ha⁻¹) and medium in phosphorus (14.2 kg ha⁻¹) and potassium (218.5 kg ha⁻¹). The experimental crop (green gram) was sown as an allay cropping in 9 years old custard apple which was planted in August, 2007 at the spacing of 5 x 5 meter. The experiment was conducted in randomized block design having Boron (0.75 kg ha⁻¹) + 0.2% Solubor for foliar spray, Boron (1.50 kg ha⁻¹) + 0.2% Solubor for foliar spray, Boron (2.25 kg ha⁻¹) + 0.2% solubor for foliar spray, Boron (0.75 kg ha⁻¹) + 0.3% Solubor for foliar spray, Boron (1.50 kg ha⁻¹) + 0.3% Solubor for foliar spray, Boron (2.25 kg ha⁻¹) + 0.3% Solubor for foliar spray, and Control (No Boron and Solubor) with variety (HUM-16) Malviya Jankalyani of green gram in three replications. The experiment consists of 7 treatment combinations. The gross and net plot size of experimental unit was 3 x 3 m² and 2.25 x 2.0 m², respectively. Row to row and plant to plant spacing was 30 cm and 10 cm, respectively. Observations of growth parameters were recorded at an interval of 20 days, i.e. 20th, 40th days after sowing and at maturity. Similarly, yield attributing characters were also recorded at harvesting. After harvesting, seed and stover samples were processed and analysed for N content

(Colorimetric method), S content (Turbidimetric method) and nutrient removal. Data obtained from various observations were analysed statistically by adopting the appropriate "Analysis of Variance" method (Gomez and Gomez, 1984). The treatment differences were tested by 'F' test of significance on the basis of null hypothesis. Critical differences were worked out at 5% level of probability where 'F' test was significant.

Results and Discussion

The experimental results revealed that application of 0.3% Solubor and 2.25 kg boron ha⁻¹ recorded significantly higher growth parameters viz. plant height, root nodules plant⁻¹, trifoliolate leaves plant⁻¹, branches plant⁻¹ and dry matter accumulation as compared to other treatments (Table 1). This may be due to important role of boron in tissue differentiation and carbohydrate metabolism. It is also a constituent of cell membrane and maintenance of conducting tissue with regulatory effect on other element. The similar result also observed by Kaisher *et al.* (2010) in green gram when they found that 30 kg S and 5 kg B ha⁻¹ was best in respect to plant height, number of branches plant⁻¹, number of pod plant⁻¹, number of seeds pod⁻¹, 1,000-seed weight, seed yield and protein content

Increase in nodule number and nodule development also observed with the application of 2.25 kg B ha⁻¹. Improvement in nodule development with the application of boron was due to its role in formation of nodule in leguminous plant concerned with precipitation of excess cation, buffer action and regulatory effect on other nutrient element. Such beneficial effect of boron with better edaphic environment available to the crop, might have improved all the growth parameters. Similar finding have also been reported by Shamsuddoha *et al.* (2011), Mathew *et al.* (2013), Parry *et al.* (2016). Similarly highest yield and yield attributing characters were also observed by the application of 0.3% Solubor and 2.5 kg B ha⁻¹ (Table 1 and 2). The beneficial effect of boron on yield attributes may be due to flower development, pollen grain formation, pollen viability, pollen tube growth for proper pollination and seed development. Similar results were reported by Mathew *et al.* (2013) in Sesamum. The findings also supported by the result of Kumar *et al.*

Table 1. Effect of Boron and Solubor on growth and yield attributes of green gram under custard apple based agri-horti system

Treatment	Plant height (cm)	Trifoliolate leaves (no. plant ⁻¹)	Root nodules (no. plant ⁻¹)	Primary branch/plant	Dry matter accumulation plant ⁻¹ (g)	Pods/plant (No.)	Pods length (cm)	Grains pod ⁻¹ (No.)	Test weight (g)
Levels of Solubor (20% B)									
0.2%	49.68	6.66	5.06	6.02	5.06	15.22	7.47	6.61	38.27
0.3%	51.23	10.97	5.22	6.27	5.22	15.64	7.72	6.79	38.73
SEm	0.26	0.95	0.14	0.13	0.14	0.19	0.14	0.08	0.31
CD (P=0.05)	0.56	2.08	NS	NS	NS	NS	NS	0.17	NS
Levels of Boron (kg ha ⁻¹) through Borax (10.5% B)									
0.75	48.22	6.47	4.92	5.92	4.92	14.72	7.25	6.40	36.82
1.50	50.10	6.68	5.11	5.93	5.11	15.30	7.55	6.63	38.58
2.25	53.05	10.29	5.40	6.58	5.40	16.28	7.98	7.07	39.00
SEm	0.31	1.17	0.17	0.15	0.17	0.23	0.13	0.10	0.38
CD (P=0.05)	0.68	2.54	0.37	0.33	0.37	0.50	0.29	0.21	0.83
Control (No solubor and No Boron)	45.47	6.13	4.63	5.83	4.63	11.67	6.83	6.03	34.03
Rest of the treatment	50.46	8.81	5.14	6.14	5.14	15.43	7.59	6.70	38.50
SEm	0.09	0.41	0.05	0.04	0.05	0.07	0.04	0.03	0.11
CD (P=0.05)	0.31	1.37	0.17	0.15	0.17	0.23	0.13	0.10	0.38

(2006) in cowpea, Singh *et al.* (2015) in mung bean and Parry *et al.* (2016), Singh *et al.* (2008) in blackgram. With increasing levels of Solubor from 0.2% to 0.3% ha⁻¹, removal of nitrogen and sulphur by seed and stover were significantly increased. Similar results were reported by Islam *et al.* (2006), Halwai *et al.* (2016).

The data on relative economics of various treatments revealed that the maximum net return (Rs. 73775 ha⁻¹) was recorded with application of 0.3% Solubor and 2.25 kg B ha⁻¹ to green gram under custard apple based agri-horti system, while the control (no Solubor and no boron) showed (Rs. 69206 ha⁻¹) value of net

Table 2. Effect of Boron and Solubor on yield and yield attributes of green gram under custard apple based agri-horti system

Treatment	Grain yield (kg ha ⁻¹)	Stover yield (kg ha ⁻¹)	Harvest index (%)	Nitrogen removal (kg ha ⁻¹)	Sulphur removal (kg ha ⁻¹)
Levels of Solubor (20% B)					
0.2%	764	2489	23.5	61.26	5.63
0.3%	789	2466	24.2	69.42	5.76
SEm ±	8.11	26.31	0.08	1.56	0.04
CD (P=0.05)	17.68	NS	NS	3.39	0.08
Levels of Boron (kg ha ⁻¹) through Borax (10.5% B)					
0.75	738	2426	23.7	61.44	5.39
1.50	753	2440	23.2	63.82	5.53
2.25	838.77	2567.04	24.6	70.76	6.17
SEm ±	9.94	32.22	0.10	1.91	0.05
CD (P=0.05)	21.65	70.21	0.21	4.15	0.10
Control (No solubor and No Boron)	710.27	2409.87	22.3	50.33	4.45
Rest of the treatment	776.78	2477.62	23.9	65.34	5.70
SEm ±	2.91	11.16	0.03	2.06	0.02
CD (P=0.05)	9.94	34.37	0.10	6.34	0.05

Table 3. Effect of Boron and Solubor on bio economics of green gram and custard apple based agri-horti system

Treatment	Cost of cultivation (Rs. ha ⁻¹)	Gross income (Rs. ha ⁻¹)				Net return (Rs ha ⁻¹)	Benefit : cost ratio
		Grain	Stover	Custard apple (1306 kg fruits ha ⁻¹)	Total		
Boron (0.75 kg ha ⁻¹) + 0.2% Solubor for foliar spray	27784	38774.67	6095.48	52246	97116.15	69332	2.50
Boron (1.50 kg ha ⁻¹) + 0.2% Solubor for foliar spray	28498	37819.43	6125.64	52246	96191.07	67693	2.38
Boron (2.25 kg ha ⁻¹) + 0.2% solubor for foliar spray	29212	42621.97	6446.20	52246	101314.17	72103	2.47
Boron (0.75 kg ha ⁻¹) + 0.3% Solubor for foliar spray	28042	39571.31	6034.47	52246	97851.78	69810	2.49
Boron (1.50 kg ha ⁻¹) + 0.3% Solubor for foliar spray	28756	38956.84	6073.51	52246	97276.35	68520	2.38
Boron (2.25 kg ha ⁻¹) + 0.3% Solubor for foliar spray	29470	44609.93	6389.02	52246	103244.96	73775	2.50
No Boron and Solubor	26754	36933.87	6024.67	52246	95204.54	68450.54	2.55

Production cost (Rs. ha⁻¹): Seed 52, Stover 2.50, fruits of custard apple 40, seasonal cost of production of custard apple Rs. 6227 ha⁻¹ for weeding, pest control and harvesting were included.

return but the benefit: cost ratio of control (no solubor and no boron) is higher (2.55) over other treatments (Table 3). This was due to the fact that application of 0.3% Solubor and 2.25 kg B ha⁻¹ exhibited increase in the seed and Stover yields and the cultivation cost was lower in control. These results were confirmed with the finding of Ram and Kumar (2009).

Conclusion

From this study it is concluded that significant increase in growth characters, yield and yield attributes of green gram were recorded with the application of 0.3% Solubor and 2.25 kg B ha⁻¹. Similarly maximum nitrogen removal and net return is also resulted in the application of 0.3% Solubor and 2.25 kg B ha⁻¹. Hence the green gram intercropped in alley of custard apple with application of 0.3% Solubor and 2.25 kg B ha⁻¹ proved to be most remunerative for the soils of semi-arid region.

References

Alloway, T.P., Gathercole, S.E., Kirkwood, H.J. and Elliott, J.G.. 2009. The working memory rating scale: A classroom-based behavioral assessment of working memory. *Learning and Individual Differences* 19(2): 242-245.

Gomez, K.A. and Gomez, A.A. 1984. *Statistical Procedures for Agricultural Research* (2nd Ed.). John wiley and Sons, New York. 680 p.

Halwai, M., Sharma, P.K., Shahi, S.K. and Singh, Y.V. 2016. Effect of boron and sulphur on yield and quality of mung bean (*Vigna radiata*) in Red

Soil of Mirzapur. *Indian Journal of Agriculture and Allied Science* 1(2): 60-64.

Havlin, J.L., Tisdale, S.L., Nelson, W.L. and Beaton, J.D. 2014. *Soil Fertility and Fertilizers: An Introduction to Nutrient Management*. 8th Edition, ISBN: 978-81-203-4868-4. pp. 290-291.

Islam, M.N., Hoque, S. and Islam, A. 2006. Effect of P × S interaction on nutrient concentration and yield of wheat, rice and mung bean. *Journal of Indian Society of Soil Science* 54(1): 86-91.

Kaisher, M.S., Rahamn, M.A., Amin, M.H.A. and Amarullah, A.S.M. 2010. Effect of sulphur and boron on the seed yield and protein content of mung bean (*Vigna radiata* L. Wilzek). *Bangladseh Research Publication Journal* 3: 1181-1186.

Kumar, A., Prasad, S. and Kumar, S.B. 2006. Effect of boron and sulphur on performance of gram (*Cicer arietinum*). *Indian Journal of Agronomy* 51(1): 57-59.

Mathew, J., George, S. and Indira, M. 2013. Effect of sulphur and boron on the performance of sesame (*Sesamum Indicum* L.) in Onattukara Sandy Soil of Kerala. *Indian Journal of Agriculture Research* 47(3): 214-219.

Meena, D.K., Das, P., Kumar, S., Mandal, S.C., Prusty, A.K., Singh, S.K., Akhtar, M.S., Behera, B.K., Kumar, K., Pal, A.K. and Mukherjee, S.C. 2013. Beta-glucan: An ideal immunostimulant in aquaculture (A Review). *Fish physiology and biochemistry* 39(3): 431-457.

Minz, S.D., Singh, A.K., Kumar, N.M. and Singh, B.K. 2021. effect of crop geometry and nitrogen management on growth attributes of pearl millet (*Pennisetum glaucum* L.) under guava based agri-horti system. *The Pharma Innovation Journal* 10(9): 2191-2195.

Parry, F.A., Chattoo, M.A., Magray, M., Gainie, S.A., Dar, Z.M. and Masood, A. 2016. Effect of different levels of sulphur and boron on growth and nodulation of garden pea (*Pisum sativum* L.). *Legume Research* 39(3): 466-469.

Ram, S.N. and Kumar, S. 2009. Performance of annona (*Annona squamosa*) based horti-pasture system relation to legumes intercropping and fertility levels under rainfed condition. *Indian Journal of Agronomy* 79(7): 493-496.

Shamsuddoha, A.T.M., Anisuzzaman, M., Suttradhar, G.N.C., Hakim, M.A. and Bhuiyan, M.A.I. 2011. Effect of sulphur and boron on nutrients in mung bean (*Vigna radiata* L.) and soil health. *Plant Resource Management* 2(2): 224-229.

Sharma, P., Singh, M.K. and Tiwari, P. 2017a. Agroforestry: A land degradation control and mitigation approach. *Bulletin of Environment, Pharmacology and Life Sciences* 6(5): 312-317.

Sharma, P., Singh, M.K., Tiwari, P. and Verma, K. 2017b. Agroforestry systems: Opportunities and challenges in India. *Journal of Pharmacognosy and Phytochemistry* 1(sp): 953-957.

Singh, A.K., Aisha, S., Singh, B.K. and Minz, S.D. 2020. Economic transformation through agroforestry in global scenario. In *Agriculture Development*

and Economic Transformation in Global Scenario (Ed. R.K. Rao), pp. 19-26. Mahima Research Foundation and Social Welfare, Varanasi.

Singh, M., Sirothia, P., Bhat, A.M., Tiwari, P. and Namdeo, A. 2015. Growth and yield of mung bean (*Vigna radiata* L.) in response to the application of sulphur and boron under rainfed condition. *The Bioscan* 10(4): 1665-1669.

Singh, R.P., Besen, J., Yadav, P.K., Singh, S.N., Singh, R.K. and Singh, J. 2008. Integrated of sulphur and molybdenum on growth, yield and quality of blackgram (*Vigna mungo* L.). *Legume Research* 31(3): 214-217.

Tamang, A., Ghosh, S.K., Garain, S., Alam, M.M., Haeberle, J., Henkel, K., Schmeisser, D. and Mandal, D. 2015. DNA-assisted β-phase nucleation and alignment of molecular dipoles in PvdF Film: A realization of self-poled bioinspired flexible polymer Nanogenerator for portable electronic devices. *ACS Applied Materials & Interfaces* 7(30): 16143-16147.

Verma, K., Sharma, P., Kumar, D., Vishwakarma, S.P. and Meena, N.K. 2021. Strategies sustainable management of agroforestry in climate change mitigation and adaptation. *International Journal of Current Microbiology and Applied Sciences* 10(01): 2439-2449.

