

## Effect of Organic and Inorganic Nutrition on Fertility Status of Soil and Yield of Vegetable Cowpea

Annu Kanwar\* and S.R. Sharma

Department of Soil Science and Agriculture Chemistry, SKN College of Agriculture, Jobner 303 329, India

Received: March 2014

**Abstract:** A field experiment was conducted during kharif, 2012 on loamy sand soil, to study the effect of organic and inorganic nutrition on fertility status of soil and yield of vegetable cowpea [*Vigna unguiculata* (L.) Walp.] var. RCV-7. The treatments consisted four organic manure (control, FYM @ 10 t ha<sup>-1</sup>, vermicompost @ 5 t ha<sup>-1</sup> and poultry manure @ 5 t ha<sup>-1</sup>) and five levels of inorganic nutrients (control, elemental sulphur @ 20 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup> + ferrous sulphate @ 50 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup> + ferrous sulphate @ 50 kg ha<sup>-1</sup> + zinc sulphate @ 25 kg ha<sup>-1</sup>). The experiment was laid out in randomized block design and replicated thrice. The levels of available nitrogen, phosphorus, potassium, sulphur, molybdenum, iron and zinc in soil at crop harvest were maximum and significantly higher under vermicompost @ 5 t ha<sup>-1</sup> (V<sub>5</sub>) over control. While significantly maximum organic carbon 0.278% was recorded with the application of FYM @ 10 t ha<sup>-1</sup> (F<sub>10</sub>) and under the M<sub>3</sub> (S + Mo + Fe). Under mineral nutrients treatments (M<sub>0</sub>-M<sub>4</sub>), the maximum available N, P, K, S, Mo, Fe and Zn in soil were recorded with the application of S + Mo + Fe + Zn (M<sub>4</sub>). The results revealed that the application of vermicompost @ 5 t ha<sup>-1</sup> and combined application of S + Mo + Fe were found significantly superior in increasing the green pod yield of cowpea over control.

**Key words:** Mineral nutrient, organic manure, green pod yield, cowpea.

Organic manure contains both macro and micro-nutrients. Improvement in available nutrient status of the soil with the incorporation of FYM alone or in combination with chemical fertilizers could be attributed to the slow decomposition of organic manure producing acids and enhancing soil biological activity. These in turn provide congenial soil physical conditions, conserve soil nitrogen and increase the availability of other nutrients. The mineralization of nutrients in the rhizosphere improves crop growth and provides a better source-sink relationship by enhancing synthesis and allocation of metabolites to reproductive organs. The application of vermicompost not only adds plant nutrients and growth regulators to the soil, but also increases aeration water retention capacity, microbial population, humic substances of the soil. Incorporation of FYM or poultry manure alone or along with chemical fertilizer improves soil biological activity, which in turn provides a congenial physical condition and improved availability of nutrient in the rhizosphere. The present investigation was undertaken to study the effect of organic

and inorganic nutrition on fertility status of the soil and yield of vegetable cowpea cv. RCV-7.

### Materials and Methods

The field experiments were conducted at S.K.N. College of Agriculture, Jobner, during 2012-2013 using cowpea cv. RCV-7. The soil was low in available nitrogen, medium in available phosphorus, sufficient in available potassium and deficient in sulphur, molybdenum, iron and zinc (134.90, 15.47, 190 kg ha<sup>-1</sup> and 7.97, 0.12, 5.34 and 0.42 mg kg<sup>-1</sup>, respectively). The soil was low in organic carbon (0.24%).

Treatments included four levels of organic nutrition (control, FYM 10 t ha<sup>-1</sup>, vermicompost 5 t ha<sup>-1</sup> and poultry manure 5 t ha<sup>-1</sup>) and five levels of inorganic nutrition (control, elemental sulphur @ 20 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup> + ferrous sulphate @ 50 kg ha<sup>-1</sup>, elemental sulphur @ 20 kg ha<sup>-1</sup> + ammonium molybdate @ 1.0 kg ha<sup>-1</sup> + ferrous sulphate @ 50 kg ha<sup>-1</sup> + zinc sulphate @ 25 kg ha<sup>-1</sup>) were replicated thrice and laid out

\*E-mail: punamsharma.soilscience@gmail.com

Table 1. Effect of organic and inorganic nutrition on available status of N, P, K and S, and Fe, Mo, Zn and OC in soil after crop harvest

Treatments	Available content (kg ha <sup>-1</sup> )				Available content			
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	S	Fe (mg kg <sup>-1</sup> )	Mo (mg kg <sup>-1</sup> )	Zn (mg kg <sup>-1</sup> )	OC (%)
<b>Organic</b>								
C <sub>0</sub> (control)	135.01	15.45	152.60	7.78	5.99	0.121	0.495	0.241
F <sub>10</sub> (FYM 10 t ha <sup>-1</sup> )	147.08	18.06	174.25	8.25	6.68	0.128	0.535	0.269
V <sub>5</sub> (vermicompost 5 t ha <sup>-1</sup> )	158.4	20.10	190.25	8.65	7.28	0.138	0.570	0.277
P <sub>5</sub> (poultry manure 5 t ha <sup>-1</sup> )	162.01	20.85	195.40	8.78	7.41	0.142	0.585	0.278
SEm±	3.35	0.66	5.56	0.12	0.20	0.003	0.011	0.004
CD (P=0.05)	9.58	1.88	15.93	0.33	0.58	0.008	0.032	0.011
<b>Inorganic nutrients</b>								
M <sub>0</sub> (control)	129.91	14.67	150.41	7.72	5.35	0.120	0.458	0.211
M <sub>1</sub> (S)	140.48	17.17	164.35	8.18	6.21	0.127	0.498	0.225
M <sub>2</sub> (S + Mo)	152.43	19.33	185.24	8.28	7.02	0.142	0.547	0.240
M <sub>3</sub> (S+Mo+Fe)	163.66	21.31	204.02	8.39	7.76	0.144	0.600	0.256
M <sub>4</sub> (S+Mo+Fe+Zn)	166.64	21.11	209.11	8.51	7.85	0.148	0.669	0.273
SEm±	3.74	0.73	6.22	0.13	0.23	0.003	0.012	0.004
CD (P=0.05)	10.80	2.12	17.96	0.37	0.66	0.009	0.036	0.013

in Randomized Block Design. Recommended doses of nitrogen @ 20 kg ha<sup>-1</sup> and phosphorus @ 40 kg ha<sup>-1</sup> were applied at sowing time.

The green pod yield of each plot was computed by sum of yield and of pod per plot of all pickings and recorded as total pod yield kg ha<sup>-1</sup>. To assess the fertility status of soil, the soil sample (0-15 cm depth) from each plot at crop harvest was taken. The samples were passed through 2 mm plastic sieve to avoid metallic contamination. The recommended dose of N, P and K was applied in the whole field prior to sowing. Whereas, sulphur and FYM were applied about three weeks (21 days) before sowing of the crop. The vermicompost and poultry manure were mixed manually in the randomly selected beds just before sowing. Mo, Fe and Zn were applied through ammonium molybdate, ferrous sulphate and zinc sulphate, respectively, through broadcasting as per treatment in randomly selected beds.

## Results and Discussion

### *Effect of organic nutrition on soil available nutrient status*

The application of organic manure significantly influenced the available contents of N, P, K, S, Mo, Fe and Zn in the soil at crop harvest (Table 1). The nutrient contents were maximum under the treatment P<sub>5</sub> (poultry manure @ 5 t ha<sup>-1</sup>). The treatment P<sub>5</sub> was found to be statistically at par with the treatment V<sub>5</sub> (vermicompost @ 5 t ha<sup>-1</sup>).

The higher availability of mineral nutrients in soil due to application of manures (vermicompost) could be ascribed to mineralization of manures, reduction in fixation and complexing properties of decomposition products of manures with micronutrients (Reddy and Reddy, 1998). Higher levels of mineral nutrients in vermicompost treated plots could also be attributed due to chelating action of organic compounds released during decomposition of organic manures, which protect these cations from fixation, precipitation, oxidation and leaching (Yadav and Kumar, 1998) of nutrients at harvest. The increase in availability of nutrients at harvest of the crop may also be due to enhanced microbial activity and nitrogen fixation by the crop, cyclic transformation of insoluble micronutrients (Mann *et al.*, 1978), enhanced mobility (Varalakshmi *et al.*, 2005), solubilization of native forms of nutrients.

### *Effect of mineral nutrition on soil available nutrient status of N, P, K, S, Mo, Fe, Zn*

The application of mineral nutrients significantly increased the available mineral contents of soil at harvest of the crop (Table 1). The maximum contents of N, P, K and Fe were recorded under the treatment M<sub>3</sub>, whereas S, Mo and Zn were maximum and significantly higher under the treatment M<sub>1</sub>, M<sub>2</sub> and M<sub>4</sub>, respectively. The improvement in nutrient status of the soil may be ascribed to more biomass (leaves, roots and dead cell of

Table 2. Nutrient composition of vermicompost and poultry manure

Nutrients	Vermicompost	Poultry manure
N (%)	1.74	1.30
P (%)	0.89	1.80
K (%)	0.86	0.80

microbes) added to soil by legumes (Hegde and Sarati, 1978), increase in symbiotic nitrogen fixation, increase in soil biomass and microbial activity. The poultry manure was rich in phosphorus (Table 2). The increase in available content of the nutrients may also be due to direct addition of these nutrients in the fields of the experimental crop. Synergism between nitrogen and iron, phosphorus and molybdenum and positive interaction between Mo and Zn may also be responsible for increase in available content of these nutrients. Similar findings were also reported by Sharma and Jain (2012).

#### *Effect of organic manure and mineral nutrients on organic carbon content of soil*

The application of organic manures and mineral nutrients significantly increased the carbon content of the soil (Table 1). The significantly maximum organic carbon was recorded in the plots that received FYM @ 10 t ha<sup>-1</sup> and mineral nutrient level M<sub>4</sub>. The significant increase in organic carbon content in the manurial treatment could be attributed to direct incorporation of the organic matter in the soil (Swarup, 1991). Addition of organic

Table 3. Effect of organic and inorganic nutrition on green pod yield of vegetable cowpea

Treatments	Green pod yield (t ha <sup>-1</sup> )
Organic	
C <sub>0</sub> (control)	15.53
F <sub>10</sub> (FYM 10 t ha <sup>-1</sup> )	17.25
V <sub>5</sub> (vermicompost 5 t ha <sup>-1</sup> )	18.57
P <sub>5</sub> (poultry 5 t ha <sup>-1</sup> )	19.01
SEm±	0.40
CD (P=0.05)	1.14
Inorganic nutrients	
M <sub>0</sub> (control)	14.61
M <sub>1</sub> (S)	16.40
M <sub>2</sub> (S+Mo)	17.97
M <sub>3</sub> (S+Mo+Fe)	19.38
M <sub>4</sub> (S+Mo+Fe+Zn)	19.59
SEm±	0.45
CD (P=0.05)	1.29

manure might have also stimulated growth and activity of microbes (Babulkar *et al.*, 2000) and due to better root growth (Varalakshmi *et al.*, 2005). The results are in line with the findings of Bhandari *et al.* (2000).

The level of built up of microbial biomass depends on available organic carbon, nutrients and other soil characteristics. Addition of nutrients in a nutrient deficient soil, as in present experimental soil, is likely to respond to applied nutrients to have high biomass as well as biomass C, N, P and S. Higher available nutrients in biomass and nutrient availability in soil are positively correlated. The addition of mineral nutrients might have resulted in the proliferation of root and shoot growth (Babulkar *et al.*, 2000) and might be due to the fact that organic and inorganic fertilizers contribute to increased biomass of the root system (Bhriuvanshi, 1988), which might have in turn increased organic carbon content of the soil. Similar findings were also reported by Thakur *et al.* (2011).

#### *Effect of organic manure on yield*

The sole application of organic manures (FYM @ 10 t ha<sup>-1</sup>, vermicompost @ 5 t ha<sup>-1</sup>, poultry manure @ 5 t ha<sup>-1</sup>) and of inorganic sources (S, Mo, Fe and Zn) significantly increased the green pod yield (Table 3). The significantly high green pod yield 19.01 and 19.59 t ha<sup>-1</sup> were recorded under treatments P<sub>5</sub> (poultry manure @ 5 t ha<sup>-1</sup>) and conjoint application of sulphur, molybdenum, iron and zinc (M<sub>4</sub>). The yields in treatments P<sub>5</sub>, V<sub>5</sub> and M<sub>4</sub>, M<sub>3</sub> were statistically at par. The higher increase in the yield has been reported to be associated with the release of macro and micronutrients during microbial decomposition (Singh and Ram, 1992). The beneficial effects of FYM/vermicompost addition also improved the soil physical properties (Kofoed, 1987). Organic matter was also a source of energy for soil microflora, which brings about the transformation of inorganic nutrients in soil in available form or applied in the form of fertilizers, which are utilized by growing plants (Sharma *et al.*, 2002). These findings corroborate with the results of several other workers (Ghanshyam *et al.*, 2010; and Singh *et al.*, 2008).

#### *Effect of mineral nutrients on yield*

The application of multinutrients combination significantly increased the yield of green pods

of vegetable cowpea. The improvement in vegetable cowpea was significant under M<sub>3</sub> (S + Mo + Fe), which remained at par with M<sub>4</sub> (Table 2). The application of mineral fertilizers alone might supply one or two nutrients only, but conjoint use of macro and micro-nutrient fertilizers and organic manure would provide all the essential nutrients in proper ratio to plant and soil and also reduce the possibilities of multiple micronutrient deficiencies in particular. It is well established fact that pulse crops require 15-20 kg N, 40-60 kg P<sub>2</sub>O<sub>5</sub> and 20 kg S ha<sup>-1</sup> for successful production (Hand Book of Agriculture, 2011). The responses of some of the micronutrients viz., Mo, Fe and Zn have also been found to be promising in increasing the productivity of the soils (Masood Ali and Mishra, 2000). Significant response of pulses to mineral nutrients has also been reported by Chavan *et al.* (2012).

## References

- Babulkar, P.S., Wandile, R.M., Badole, W.P. and Balpande, S.S. 2000. Residual effect of long term application of FYM and fertilizers on soil properties (vertisols) and yield of soyabean. *Journal of the Indian Society of Soil Science* 48: 89-92.
- Bhandari, A.L., Walia, S.S. and Singh, T. 2000. Production sustainability of maize-wheat system in a Typic Ustipsamment soil as influenced by integrated nutrient sources. *Proceedings of International Conference on Managing Natural Resources for Sustainable Agriculture Production in the 21<sup>st</sup> Century* 3: 889-890, New Delhi.
- Bhargavanshi, S.R. 1998. Long term effect of high dose of farm yard manure on soil properties and crop yield. *Journal of the Indian Society of Soil Science* 36: 784-786.
- Chavan, A.S., Khafi, H.R., Raj, A.D., Parmar, R.M. and Shekh, M.A. 2012. Effect of potassium and zinc on growth and yield of cowpea [*Vigna unguiculata* (L.) Walp.]. *Research on Crops* 6(2): 432-434.
- Ghanshyam, Kumar, R. and Jat, R.K. 2010. Productivity and soil fertility as affected by organic manure and inorganic fertilizer in green gram (*Vigna radiata*) wheat (*Triticum aestivum*) system. *Indian Journal of Agronomy* 55: 16-21.
- Hand Book of Agriculture 2011. Published by Directorate of Knowledge Management in Agriculture. ICAR, New Delhi. p. 1096.
- Hegde, D.M. and Saraf, C.S. 1978. Effect of intercropping and phosphorus fertilization of pigeonpea on fertility status of soil. *Indian Journal of Agronomy* 23: 372-373.
- Kofoed, A.D. 1987. The significance of FYM. *Kodemines Tidsskrift* 19: 37-63.
- Masood Ali and Mishra, J.P. 2000. Nutrient management in pulses and pulses based cropping system. *Fertilizer News* 45(4): 59-69.
- Mann, M.S., Takkar, P.N., Bansal, R.L. and Randhawa, N.S. 1978. Micronutrient status of soil and yield of maize and wheat as influenced by micronutrient and farm yard manure application. *Journal of the Indian Society of Soil Science* 26: 208-214.
- Reddy, Gopal, B. and Reddy, Suryanarayana 1998. Effect of organic manures and nitrogen levels on soil available nutrient status in maize-soybean cropping system. *Journal of the Indian Society of Soil Science* 46: 474-476.
- Singh, R.P., Bisen, J.S., Yadav, P.K., Singh, S.N., Singh, R.K. and Singh, J. 2008. Integrated use of sulphur and molybdenum on growth, yield and quality of blackgram (*Vigna mungo* L.). *Legume Research* 31(1): 214-217.
- Singh, P.N. and Ram, H. 1992. Effect of phosphorus and sulphur on concentration and uptake of micronutrients in chickpea. *Journal of the Indian Society of Soil Science* 40: 307-312.
- Sharma, S.R., Bhandri, S.C. and Purohit, H.S. 2002. Effect of organic manure and mineral nutrients on nutrient uptake and yield of cowpea. *Journal of the Indian Society of Soil Science* 50: 475-480.
- Sharma, S.K. and Jain, N.K. 2012. Effect of balanced fertilization on productivity and soil fertility status of clusterbean. *Legume Research* 35(1): 32-35.
- Swarup, A. 1991. Longterm effect of green manuring (*Sesbania aculeate*) on soil properties and sustainability of rice and wheat yield on a sodic soil. *Journal of the Indian Society of Soil Science* 39: 777-780.
- Thakur, R., Sawarker, S.D., Vaishya, U.K. and Singh, M. 2011. Impact of continuous use of inorganic fertilizer and organic manure on soil properties and productivity under soybean-wheat intensive cropping of a vertisol. *Journal of the Indian Society of Soil Science* 59(1): 74-81.
- Varalakshmi, L.R., Srinivasamurthy, C.A. and Bhaskar, S. 2005. Effect of integrated use of organic manures and inorganic fertilizers on organic carbon, available N, P and K in sustaining productivity of groundnut-finger millet cropping system. *Journal of the Indian Society of Soil Science* 53: 315-318.
- Yadav, D.S. and Kumar, A. 1998. Integrated use of organic and inorganic in rice wheat cropping system for sustained production. In *Long Term Soil Fertility Management Through Integrated Plant Nutrient Supply* (Eds. A. Swarup, D. Damodar Reddy and R.N. Prasad), pp. 247-255. Indian Institute of Soil Science, Bhopal.