

Incidence of *Listeria* in fish and shellfish

M.M. Prasad¹ and Nirmala Thampuran¹

Central Institute of Fisheries Technology, Burla Research Centre,
Burla - 768 017, Sambalpur, Orissa.

¹ Microbiology, Fermentation & Biotechnology Division,
Central Institute of Fisheries Technology, Cochin - 682 029, India

Listeria monocytogenes has been recognized as an important food-borne pathogen since 1981 and seafoods have been implicated in several sporadic cases of Listeriosis. In the present study fresh (57), iced (49) and frozen (84) fish and shellfish from cultured ponds and commercial outlets were screened for *Listeriae*. Among the samples of fresh cultured fish, 1.8% contained *Listeria innocua*. The iced *Penaeus indicus* (PUD) and *Penaeus monodon* contained 20% *Listeria* spp., comprising *L. innocua* and *L. ivanovii*. *Listeria* spp. were detected in 14.28% of the frozen fish samples also. *Listeria monocytogenes* could not be detected in any of the samples.

Key words : *Listeria* spp, Fish, Shellfish, marine, cultured, Andhrapradesh, cold enrichment

The isolation of *Listeria monocytogenes* from frozen cooked crab meat from a Mexican supplier by a commercial laboratory in USA in 1987 led to enhanced concern about the potential danger of *Listeria* contaminated seafood to public health (Anon, 1987). The first case of Listeriosis linked to consumption of fish and seafood was reported in 1989 (Facinelli *et al*, 1989). Seafood have also been implicated in several sporadic cases of Listeriosis and USDA requires zero tolerance of *L. monocytogenes* in ready to-eat foods including seafoods (Fredikssen, 1991; Fuchs & Reilly, 1992; Baker *et al*, 1993).

The psychrotrophic nature of *L. monocytogenes* allows survival or even multiplication of this potential pathogen during refrigeration or temperature-abuse conditions (Jinneman *et al*, 1999). Studies carried out on peelers and processors of seafood processing plants in Mangalore, India, revealed that nearly 4% of the seafood handlers were found to carry *L. monocytogenes* (Jeyasekaran *et al*, 2002a). In the same study floors, drains, tables, containers, and equipments of the seafood processing plants were

found to harbour *Listeria* spp. Investigations on prevalence of *Listeria* spp. in processed fishery products namely, frozen shrimps, squids and fish from the seafood processing plants in Mangalore area have revealed that 11.1% samples were found to contain *Listeria* spp. (Jeyasekaran *et al*, 2002b). These studies indicated that the process of chilling and freezing did not completely eliminate *L. monocytogenes* in shrimps (Jeyasekaran *et al*, 2002c). In view of the potential pathogenic nature and public health significance, the present study was undertaken to screen fish and shellfish for *Listeria* spp. from marine and cultured sources of Andhra Pradesh region.

Materials and Methods

Fish and shellfish samples were collected from commercial outlets and culture ponds in sterile polythene bags and were brought to the laboratory immediately. In case of frozen shellfish, the sample was drawn from all four sides and from centre of the block with a sterile poker in aseptic conditions and after blending, 30 g sample was weighed out aseptically for further analyses.

¹ Corresponding author E-mail: prasadmm@hotmail.com or prasadmothadaka@yahoo.com

Two methods were employed simultaneously for screening of fish and shellfish for *Listeriae*. In the first procedure of USDA method as outlined by Fuchs, (1991), 30 g sample was taken into 270 ml of pre-enrichment broth (tryticase soy broth with 0.6% yeast extract). Sample was triturated with small amounts of TSB employing sterile pestle and mortar in aseptic conditions and in the end whole of TSB was mixed. It was incubated at 30°C for 24h. From this, 10ml was taken and mixed in 90 ml of UVM1 and it was incubated at 30°C for 24h. After the incubation, 0.1ml of UVM1 was added to 10ml of UVM II and it was incubated further at 30°C for 24h. The blackened tubes were streaked on MOX (Modified Oxford Medium) and the plates were incubated at 35°C for 24 & 48 h. Colonies surrounded by black halos were picked for further tests.

In the second method, Cold Enrichment procedure was employed for screening the fish and shellfish samples for *Listeriae*. 30 g sample was weighed aseptically and homogenized with 270 ml sterile enrichment broth and was incubated at 4°C. Tryptose phosphate broth (Hi-media, India) was employed as enrichment broth. At weekly intervals inoculum was taken with sterile loop for subculture on Modified McBrides Agar (MMA) plates. The MMA plates were incubated at 35°C for 48h. Plates were examined using Henry's illumination technique. The composition of MMA was Phenylethanol agar (Difco): 35.5g; Glycine anhydride (Sigma, USA) 10g; Lithium

chloride (Sigma, USA) Distilled water 1 l, pH was adjusted to 7.3. The ingredients were heated, dissolved, dispensed and autoclaved at 121°C for 15 min. Prepared bottles (100ml) each were stored at 4°C. After melting the agar, filter sterilized cyclohexamide 20mg/100ml was added. The plates were poured thin and were stored at 4°C and were used within week's time whenever necessary.

Suspected colonies were picked up and were purified on Trypticase Soy Agar (TSA) with 0.6% yeast extract (YE) plates. The isolates were further tested for morphological and biochemical characters by standard methods (Jones & Seeliger, 1992).

L. monocytogenes NCTC (11994), *Staphylococcus aureus* (NCTC 1803) and *Rhodococcus equi* NCTC (1621) were the test cultures used in this study.

Results and Discussion

Incidence of *Listeria* in fresh fish is shown in Table 1. Out of 57 samples screened from culture ponds and local outlets, only *Labeo rohita* harboured *Listeria spp* and it was identified as *L. innocua*. *L. innocua* is not considered to cause disease in humans (Rocourt & Seeliger, 1985) and has been reported in 33% freshly caught marine and brackish fish samples (Fuchs & Surendran, 1989).

Occurrence of *Listeria* in iced fish and shellfish in the present study are shown in Table 2. *Listeria monocytogenes* was not detected in iced fish, *Penaeus indicus* (Indian

Table 1. Incidence of *Listeria* in fresh fish (n= 57)

No. of samples screened	Description of the sample	Source	Samples Positive for <i>Listeria spp</i> .
10	<i>Hypophthalmichthys molitrix</i>	Local outlet	Negative
10	<i>Cirrhinus mrigala</i>	Culture Ponds	Negative
10	<i>Labeo rohita</i>	Culture Ponds	Positive (1) <i>L. innocua</i>
5	<i>Johnius dussumieri</i>	Local outlet	Negative
17	<i>Mugil cephalus</i>	Local outlet	Negative
5	<i>Megalops cundinga</i>	Local outlet	Negative

Table 2. Incidence of *Listeria* in iced fish and shellfish (n=49)

No.	Description of the sample	Source	Samples Positive for <i>Listeria</i> spp.	<i>Listeria</i> spp isolated / No. samples
4	<i>Cirrhinus mrigala</i> (iced for 5 days)	Culture Ponds	Nil	
3	<i>Cirrhinus mrigala</i> (kept on ice for 13 days)	Culture Ponds	Nil	
16	<i>Penaeus indicus</i> (PUD)	Local outlets	3	2 <i>Listeria innocua</i> (2) <i>Listeria ivanovii</i> (1)
9	<i>Penaeus monodon</i>	Local outlets	2	<i>L.innocua</i> (1) <i>L.ivanovii</i> (1)
11	<i>Penaeus indicus</i> (P.D)	Local outlets	Nil	
6	<i>Parapenaeopsis styliifera</i> (8 days in ice)	Local outlets	Nil	

white prawn) (PD) or in deep-sea shrimp. Other *Listeria* spp. were detected in 10.2% of the iced fish and shellfish samples screened. Among the *Penaeus indicus* (PUD) and *Penaeus monodon*, 20% were positive for *Listeria* spp., and the predominant species were *Listeria innocua* followed by *L. ivanovii*.

The occurrence of *Listeria* in frozen fish and shellfish samples is depicted in Table 3. Out of 84 samples of frozen fish and shellfish screened 14.28% were positive for *Listeria* spp. *Listeriae* were not detected in any of the deep-sea prawns and fish brought for analysis. Both PUD and PD *P. indicus*

harboured *Listeria* spp., at 18.75 and 16.66%, respectively. Interestingly *L. innocua* is dominant flora in PUD prawns while PD prawns harboured mainly *L. ivanovii*. In this study occurrence of *L. monocytogenes* was not seen in any of the samples screened. Earlier reports also did not show occurrence of *L. monocytogenes* in seafoods. However, *L. innocua* and *L. grayi* were detected (Fuchs & Surendran 1989; Kamat & Nair 1994).

In the present study the incidence of *Listeria* spp is more in shrimps than in fishes. This is in agreement with the report of Jeyasekaran *et al* (2002b) in which the

Table 3. Incidence of *Listeria* in frozen fish and shellfish (n=84)

No. of samples screened	Sample	Source	Samples Positive for <i>Listeria</i> spp.	<i>Listeria</i> spp/ No. positive
3	<i>Nemipterus japonicus</i>	Market	Nil	—
3	<i>Upeneus vittatus</i>	Market	Nil	—
6	<i>Parapenaeopsis styliifera</i>	Deep sea	Nil	—
16	<i>P. indicus</i> (PUD)	Local outlets	3	<i>L. innocua</i> (2) <i>L. ivanovii</i> (1)
8	<i>P. monodon</i> (Headless)	Local outlets	1	Positive (1) <i>L. innocua</i>
18	<i>P. indicus</i> (PD)	Local outlets	3	Positive (3) <i>L. innocua</i> (1) <i>L. ivanovii</i> (2)
10	<i>P. monodon</i> (PUD)	Local outlets	2	<i>L. ivanovii</i> (2)
20	<i>Metapenaeus dobsoni</i>	Local outlets	3	<i>L. innocua</i> (1) <i>L. ivanovii</i> (2)

Table 4. Comparison of detection of *Listeria* by cold enrichment and USDA methods

Sample	No samples positive for <i>Listeria</i>		No. of isolates positive for <i>Listeria</i> by	
	Cold enrichment	USDA	Cold enrichment	USDA
			No. positive / total isolates	No. positive / total isolates
Fresh fish	1	1	4/4	4/4
Fresh shrimp	None	None	NA	NA
Iced fish	Not detected	Not detected	Not detected	Not detected
Iced shrimp	5	5	25/25	22/25
Frozen fish				
Frozen shrimp	12	11	77/77	61/72
Total	18	17	106	87

occurrence of *Listeria* spp. in shrimps was very high (14.3%), when compared to its incidence in finfishes (4.3%). The positive isolates were identified as *L. monocytogenes* and *L. innocua* (Jeyasekaran *et al*, 2002b), whereas, in the present study the distribution in occurrence of *L. ivanovii* and *L. innocua* was equal. The occurrence of *Listeria innocua* in sea food samples is also reported by Weagent *et al*, (1988), Fuchs & Sirvas, (1991) and Jeyasekaran *et al*, (2002b).

A comparison of detection of *Listeria* from different fish and shellfish samples by cold enrichment method and *Listeria* isolation method is shown in Table 4. A perusal of data in the Table 4 reveals that wherever *Listeria* presence is detected by cold enrichment method in the fresh, iced and frozen fish and shell fish samples (18), all the suspected *Listeria* isolates were confirmed as *Listeria* spp while in the USDA *Listeria* isolation method confirmation of the suspected isolates were slightly lower (17/18) in comparison with cold enrichment method. Hayes *et al* (1991) made similar observations in the study on comparison of cold enrichment methods and USDA-FSIS method. Biester & Schwarte (1939) reported for the first time detection of *Listeria* by cold enrichment method. Fatty acid (anteiso- $C_{15:0}$) is reported to play an important role in adaptation of *Listeria* to cold temperatures, with mutants deficient in this fatty acid being shown to be cold sensitive (Annous *et al*, 1997).

During the process of isolation, some colonies were seen to have morphological characters similar to *Listeriae*. However, the major hindrance was subculture on TSA plates. In this study 14 of the 120 suspected *Listeria* isolates belonged to this category. Since these isolates were from iced or processed samples, there is a possibility for viable but non-culturable (VBNC) state. Besnard *et al*, (2000) examined factors inducing *L. monocytogenes* into the VBNC state and concluded that inoculum size, temperature (4°C and 20°C), NaCl concentration (0-7%), pH (5 and 6), and presence of sunlight were contributory factors for development of VBNC state. In the present study the samples were exposed to all the conditions mentioned at different stages of post harvest handling. The presence of VBNC cells of this pathogen if present, could pose a major public health hazard since the cells cannot be detected by routine culture methods (Oliver, 2005). The study of Cappelletti *et al*, (2005) suggested that *Listeria monocytogenes* might remain in the aquatic environment for prolonged period in the VBNC state; but did not exhibit any pathogenicity in the conditions tested and stressed the role of VBNC cells in environmental transmission and virulence of *L. monocytogenes*.

This study revealed that out of 190 fresh, iced and frozen shellfish samples screened 9.5% of them harboured *Listeria* spp

with equal distribution of *L. innocua* and *L. ivanovii*. None of the samples screened were positive for *Listeria monocytogenes*. Cold enrichment facilitated higher number of confirmed isolates for *Listeria spp* than the USDA method.

Authors wish to thank Dr. K. Devadasan, Director, CIFT, Kochi, for encouraging carrying out the work and permission to publish the work. Thanks are due to technical staff at Visakhapatnam Research Centre of CIFT for the support during the course of this study.

References

- Annous, B.A., Becker, L.A., Bayles, D.O., Labeled, D.P and Wilkinson, B.J. (1997) Critical role of anteiso-C_{15:0} fatty acid in the growth of *Listeria monocytogenes* at low temperatures. *Appl. Environ. Microbiol.* **63**, pp 3887-3894.
- Anonymous, (1987). *First Listeria finding in crabmeat confirmed by FDA*. Food Chem. News. **29**, 38.
- Baker, M., Brett, M., Short, P., Calder, L. and Thointon, T. (1993) *Listeriosis* and mussels. Communicable Diseases in New Zealand. **93**, pp 13-14.
- Besnard, V., Federighi, M., Cappelier, J.M. (2000) Development of direct viable count procedure for the investigation of VBNC state in *Listeria monocytogenes*. *Lett. Appl. Microbiol* **30**, pp 1-6.
- Biester, H.E and Schwarte, L.H (1939) Studies on *Listerella* infection in sheep. *J. Infect. Dis.* **64**, pp 135-144.
- Cappelier, J.M., Besnard, V., Roche, S., Garrec, N., Zundel, E., Velge, P., Federighi, M. (2005) Avirulence of Viable But Non-Culturable *Listeria monocytogenes* cells demonstrated by in vitro and in vivo models. *Veterinary Research*, **36**, pp 589-599.
- Facinelli, B., Varaldo, P.E., Toni, M., Casolari, C and Fabio, V. (1989) Ignorance about *Listeria*. *Brit Med J.* **199**, p738.
- Frediksen, W. (1991) *Listeria* epidemiology in Denmark 1981-1990. In: Proceedings of International Conference on Listeria and Food Safety, Laval, France. ASEPT (eds) pp 48-49.
- Fuchs, R.S. (1991) *Listeria monocytogenes*. *ASEAN Food J.* **6**, pp 3-13.
- Fuchs, RS. and Reilly, P.J.A. (1992) The incidence and significance of *Listeria monocytogenes* in seafoods. In: Proceedings of an International Conference on "Quality Assurance in the Fish Industry", (Huss, H.H., Jakobsen, M & Liston, J., Eds) pp 217-230. Copenhagen, Denmark, Elsevier Science Publishers B.V., The Netherlands.
- Fuchs, R.S. and Sirvas, S. (1991) Incidence of *Listeria monocytogenes* in acidified fish product, Ceviche. *Lett Appl Microbiol.* **12**, pp 88-90.
- Fuchs, R.S. and Surendran, P.K. (1989) Incidence of *Listeria* in tropical fish and fishery products. *Lett Appl Microbiol.* **9**, pp 49-51.
- Hayes, P.S., Graves, L.M. Ajello, G.W., Swaminathan, B., Weaver, R.E., Wenger, J.D., Schuhat, A., Broome, C.V. and the listeria study group (1991). Comparison of cold enrichment and the US Department of Agriculture methods for isolating *Listeria monocytogenes* from naturally contaminated foods. *Appl. Environ. Microbiol.* **57**, pp 2109-2113.
- Jeyasekaran, G., Indrani Kaarunasagar. and I. Karunasagar. (2002,a) Prevalence of *Listeria spp.* in Seafood Handlers. *J Food Sci. Technol.* **39**, pp 173-175.
- Jeyasekaran, G., Indrani Kaarunasagar. and I. Karunasagar. (2002,b) Occurrence of *Listeria spp.* in Processed Fishery Products. *J Food Sci. Technol.* **39**, pp 188-191.
- Jeyasekaran, G., Indrani Kaarunasagar and I. Karunasagar. (2002,c) Effect of Chilling and Freezing on the Survival of *Listeria monocytogenes* in Shrimps. *J Food Sci. Technol.* **39**, pp191-193.
- Jinneman, K.C., Wekell, M.M. and Eklund, M.W. (1999). Incidence and behaviour of

- Listeria monocytogenes* in fish and seafood. In: *Listeria, Listeriosis and food safety*. 2nd Edn, Revised and Expanded. (Ryser, E.T and Marth. E.H., Eds), pp601-630, Marcel Dekker Inc., New York. Basel.
- Jones, D. and Seeliger, H.P.R. (1992) The Genus *Listeria* In: *The Prokaryotes: a handbook on the biology of bacteria: Eco-physiology, isolation, identification, applications* (Balows, A., Truper, H.G., Dworkin, M., Harder, W. and Schleifer, K.H. Eds), pp 1595-1616. Springer-Verlag, New York.
- Kamat, A.A. and Nair, P.M. (1994) Incidence of *Listeria species* in Indian seafoods and meat. *J Food Safety*. **14**, pp 117-130.
- Oliver, J.D. (2005) Viable But Nonculturable Bacteria in Food Environments. In: *Foodborne pathogens: Microbiology and Molecularbiology* Fratamico, P.M., Bhunia, A.K and Smith, J.L. (Eds), Horizon Scientific Press. pp 99-112.
- Rocourt, J. and Seeliger, H.P.R. (1985). Distribution des especes du genre *Listeria*. *Zentralblatt fur Bakteriologie, Mikrobiologie, Infektionskrankheiten und Parasitologie (Stuttgart)*. **259**, pp 317-330.
- Weagant, D., Sado, P.N., Colbum, K.G., Torkelson, J.D., Stanley, F.A., Krane, M.H., Shields, S.C. and Thayer, C.F. (1988) The incidence of *Listeria species* in frozen seafood products. *J Food Prot*. **51**, pp 655-657.