

Modified Atmosphere Packaging of Fish – A Review

T.K. Srinivasa Gopal

Principal Scientist

and

C.N. Ravi Shankar

Senior Scientist

*Central Institute of Fisheries Technology,
Matsyapuri P.O., Cochin - 682 029*

Shelf life and quality of fresh fish can be extended by the application of modified atmosphere packaging using high barrier packaging film and refrigerated storage. Modified atmosphere with high concentration of CO₂ inhibits or slows down the growth of various aerobic spoilage bacteria of fish products by extending the lag phase. It provides conditions for the growth of gram positive bacteria and food pathogens within the package. Major quality hazard is the risk of foodborne botulism. This paper reviews the literature relating to the scientific basis of these claims.

Key words : Modified atmosphere packaging, vacuum packaging, Seafood, Microbiological safety, Sensory quality

Fish and other sea products are susceptible to spoilage from postmortem microbial growth, biochemical end products resulting from the microbial growth or combinations of both. Their low body temperature provides a natural environment for psychrotrophic spoilage microflora (Brody, 1989). Presently ice and mechanical refrigeration are the most common means of retarding microbial and biochemical spoilage in freshly caught seafood during distribution and marketing. However as the ice melts it tends to contaminate fresh products, accelerating spoilage and reducing shelf life (Brody, 1989).

Modified atmosphere packaging (MAP), a technologically viable method has been developed as a supplement to ice or mechanical refrigeration to reduce the losses and extend the storage life. According to the definition of Sacks and Gore (1987), MAP is replacement of air in a pack by a different mixture of gases, where the proportion of each component is fixed when the mixture is introduced, but no further control is exercised during storage. Two forms exist, namely vacuum packaging and gas, gas flush or gas exchange packaging. Vacuum packaging (VP) involves placing a product in a film of low oxygen permeability, the removal of air from the package and the application of a hermetic seal (Smith *et al.*, 1990). Gas packaging is an extension of this process involving removal of air from the pack and its replacement with specific

gases, either singly or in combination. In vacuum packaging the two major spoilage agents namely aerobic bacteria and oxidative reactions require oxygen. Therefore its unavailability will inhibit spoilage and thus maximize quality and /or storage life. Some deterioration, however will occur due to anaerobic / microaerophilic organisms and nonoxidative reactions. This is usually minimized by chilled storage. Product compression is unavoidable and makes vacuum packaging unsuitable for many products. Controlled atmosphere is packaging in an atmosphere where the composition of gases is continuously controlled throughout storage. The gaseous atmosphere of the vacuum packaging is altered during storage, thus considered modified due to a 10 to 20% increase in the carbon dioxide content produced by microbial activity. This carbon dioxide may inhibit the growth of undesirable organisms (Silliker & Wolfe, 1980).

Development of modified atmosphere packaging for fish

Development started in the 1930s with several workers observing that CO₂ atmosphere prolonged the storage life of fish (Coyne, 1933; Stansby & Griffiths, 1935; Hjorth-Hansen, 1933). Subsequent interest in the use of CO₂ turned to refrigerated seawater systems as an alternative to the use of iced storage for small fish on board or for road transport (Nelson & Barnet, 1971;

Barnett *et al.*, 1978; Hiltz *et al.*, 1976; Bullard & Collins, 1978). Some interest in the technically more difficult task of transporting chilled, whole fish in carbon dioxide atmospheres then appeared. This was a period when all variables associated with manipulation of storage temperatures were keenly examined. The earliest use of CO₂ in retail products exploited the solubility of CO₂ to produce a snugging down effect (Douglas, 1970), regarded as characteristic of CO₂ packs. The vacuum appearance and prevention of movement of the product was seen, ironically in view of the later developments in MAP, as being advantageous but there was only a small amount of kipper fillet packed in this way, with no application to white fish or other fish products (Abbey, 1970). Reductions in rates of deterioration of several species of fish have been recorded in tests using pressure chambers under both hyperbaric (Charm *et al.*, 1977) and hypobaric (Varga *et al.*, 1980; Haard & Lee, 1982) conditions. With unfrozen material, nitrogen flushing for oxygen sensitive products was recommended and applied to shrimps and prawns, but was not recommended for meat and fish which are susceptible to spoilage by anaerobic bacteria (Anon, 1967). As had occurred earlier with CO₂ preservation of whole fish, development and application of controlled gas mixtures for consumer packaging was concerned first with foods other than fish (Schweisfurth & Kalle, 1970; Anon, 1977). According to Kimber (1984), the technology of gas packaging was first perfected and patented in 1963 by Bohme and Kalle Films but it took until 1977 to produce gas flushed packs successfully. The first UK application to fish products was in 1979 in Northern Ireland (Kimber, 1984), with a few specialty products. The technique became more widespread as manufacturers of vacuum packaging equipment adopted their products, and super markets sought alternative ways of presenting fish. With equipment capable of making more precisely designed and practicable products, research workers turned their attention to the demands of prepackaging fish products and the use of gases – mainly CO₂. Trials were conducted with a large range of species and products and many of the results have been published (Martin, 1981; 1982; Cann, 1984; Anon., 1985). There has been an increase in gas packaged food products in the market during the last three decades. This increase has brought improvements to the packaging industry, which has led to the

development of high barrier polymers and thermo mold packaging equipment.

The three main commercially used gases in modified atmosphere packaging are carbon dioxide, nitrogen and oxygen.

Function of carbon dioxide

CO₂ is soluble in water and also in lipids. Its general effects on microorganisms are an intensification of their growth stage and a decrease in the growth rate during the logarithmic stage (Farber, 1991). CO₂ is soluble in water, forming carbonic acid that may lower the pH (Smith *et al.*, 1990). The anti microbial efficacy of CO₂ is greatly enhanced as the storage temperature of the product is reduced due to its increasing solubility with decreasing temperature. Thus to guarantee maximum anti microbial effect, the storage temperature is kept as low as possible (Finne, 1982; Farber, 1991).

The concentration of CO₂ in the food is dependent on water and fat content of the products, and of the partial pressure of CO₂ in the atmosphere, according to Henry's law (Ho *et al.*, 1987). Devlieghere *et al.* (1998a, 1998b) have demonstrated that the growth inhibition of microorganisms in MAP is determined by the concentration of dissolved CO₂ in the product. After the packaging is opened, CO₂ is slowly released by the product and it continues to exert a preservative effect for a certain period of time, referred to as CO₂'s residual effect (Stammen *et al.*, 1990).

The action of CO₂ on the preservation of foods was originally thought to be caused by displacement of some or all of the O₂ available for bacterial metabolism, thus slowing growth (Daniels *et al.*, 1985). However, experiments with storage of bacon and pork showed a considerable increase in shelf life under 100% CO₂ compared with storage in normal air atmospheres (Callow, 1932), but the preservative effect was not because of the exclusion of O₂, as storage in 100% nitrogen (N₂) offered no advantage over normal air storage.

A drop in surface pH is observed in MAP products because of the acidic effect of dissolved CO₂, but this could not entirely explain the bacteriostatic effect (Coyne, 1933). It was shown that CO₂ was more effective at lower temperatures and that the change in pH caused by CO₂ did not account for the retardation of growth. In a study on several pure cultures of bacteria isolated from fish products, CO₂ atmospheres

were found to inhibit the growth of bacteria markedly, whereas normal growth patterns were observed under air or N_2 atmospheres (Coyne, 1932). It was also observed that bacterial growth was inhibited even after the cultures were removed from the CO_2 atmosphere and transferred to an air environment, interpreted as residual effect of CO_2 treatment. Bacterial growth was distinctly inhibited under 25% CO_2 and almost no growth was observed under higher CO_2 concentrations for 4 days at 15°C. The results obtained could neither be explained by the lack of O_2 nor the pH effect. The effect of CO_2 on bacterial growth is complex and four activity mechanisms of CO_2 on microorganisms has been identified (Parkin & Brown, 1982; Daniels *et al.*, 1985; Farber, 1991)

1. Alteration of cell membrane function including effects on nutrient uptake and absorption;
2. Direct inhibition of enzymes or decreases in the rate of enzyme reaction;
3. Penetration of bacterial membranes, leading to intracellular pH changes;
4. Direct changes in physico-chemical properties of proteins

Probably a combination of all these activities account for the bacteriostatic effect. In food having high moisture and/ or fat amounts, the excessive absorption of CO_2 may lead to a phenomenon known as packaging collapse (Parry, 1993). A certain amount of CO_2 will dissolve into the product to inhibit bacterial growth (Gill & Penney, 1988). The ratio between this volume of gas and volume of food product (G/P ratio) should usually be 2:1 or 3:1. This high G/P ratio is also necessary to prevent package collapse because of the CO_2 solubility in wet products. Dissolved CO_2 takes up much less volume compared with CO_2 gas and after packaging a product in CO_2 atmosphere, a drop in pressure develops within the package and package collapse occurs. Increase in dripping is caused by the dissolution of gases on the muscle surface in atmospheres containing high CO_2 levels (>60%), reduced pH and, consequently, low protein water retention ability (Parry, 1993; Randell *et al.*, 1995). As a consequence, high CO_2 concentrations promote organoleptic changes as, for example, texture alterations in meat.

Function of nitrogen

N_2 is an insipid and inert gas, showing low solubility in water and lipids. It is used for

displacing the oxygen from the packaging, decreasing oxidative rancidness and inhibiting the growth of aerobic microorganisms (Farber, 1991). Due to its solubility, it is used as a filling gas, preventing the possible packaging collapse caused by the accumulation of CO_2 .

Function of oxygen

O_2 generally stimulates the growth of aerobic bacteria and may inhibit the growth of exclusively anaerobic bacteria, although anaerobic microorganisms show different sensitivity levels to oxygen (Farber, 1991). The presence of oxygen may cause oxidative rancidity problems in fish presenting high lipid amounts, promoting the formation of low molecular weight aldehydes, ketones, alcohols and carboxylic acids. The use of oxygen in modified atmospheres is generally avoided with this kind of fish, in order to minimize such effects. The use of O_2 in modified atmosphere packaging for fish is supported by Davis and Slade (1995), who states that there are evidences showing that the use of O_2 reduces the exudation in fish during storage, and suggested that O_2 can be used in low fat fishes. Reddy *et al.* (1992) claim that the use of oxygen associated with N_2 or CO_2 gives a false idea of reducing botulinum risks in fresh packed fish and may lead to illusory safety. However for some products oxygen could or should be used. High levels of oxygen are used in red meat and red fish meat to maintain the red colour of the meat, to reduce and retard the browning caused by formation of metmyoglobin. Oxygen in MAP packages of fresh fish will also inhibit reduction of TMAO to TMA (Boskou & Debevere, 1997).

Examples of some meat and fish products currently gas packaged as well as the composition of gas mixtures used to extend the shelf life of each product are shown in Table 1.

Table 1. Examples of gas mixtures for selected food products

Product	Temp°C	% O_2	% CO_2	% N_2
Fresh meat	0-2	70	20	10
Cured meat	1-3	0	50	50
Poultry	0-2	60-80	20-40	0
Fatty fish	0-2	0	60	40
White fish	0-2	30	40	30
Cheese	1-3	0	60	40
Baked product	R.T	0	60	40
Pizza	R.T	0	60	40
Dry snacks	R.T	0	20-30	70-80

R.T - Room temperature; (Smith *et al.*, 1990)

Factors influencing shelf life of fish packaged under modified atmosphere

Several interrelated factors influence the shelf life and keeping quality of fish packaged under CO₂ enriched atmospheres. These include concentration of gas mixture, packaging film permeability, storage temperature and microbial contamination.

i) Concentration of gases

Concentration of gases is a very important factor for shelf life of fish packaged under MAP conditions. The bacteriostatic effect of MAP storage increases with increasing CO₂ concentrations (King & Nagel, 1967; Enfors & Molin, 1980; Gill & Tan, 1980). With cod fillets stored in different concentrations of CO₂ and O₂/N₂, Stenstrom (1985) showed that the shelf life increased as the CO₂ concentration increased. Haines (1932) reported that concentrations as low as 10 to 20% CO₂ were sufficient to inhibit the growth of *Pseudomonas* and *Achromobacter* species. High CO₂ concentrations might be used for many types of seafood with little risk of heme protein discoloration because of the low pigment concentration (Parkin & Brown, 1982). Carbon dioxide in sufficient concentrations can inhibit the growth of spoilage molds (Brown, 1922; Moran *et al.*, 1932; Tomkins, 1932; Hintlian & Hotchkiss, 1986). Coyne (1933), in one of the original studies on use of carbon dioxide atmospheres to prolong fish quality, recommended concentrations of 40 and 60% CO₂. Shewan (1949) recommended concentrations between 30 and 40% CO₂ atmospheres and reported superior quality for fish stored in the higher concentration of CO₂. Tarr (1954) recommended a minimum of 40 to 50% CO₂ in the package headspace to derive maximum benefit of CO₂ storage of fresh fish. Based on these studies, approximately 50-60% CO₂ in the package headspace is commonly used, with the remainder comprising of a mixture of O₂ and N₂.

ii) Packaging film permeability

MAP can be effective if used in conjunction with packaging materials of correct O₂/CO₂ permeability characteristics. It is no use having the correct atmosphere if the film allows the atmosphere to change too rapidly. The properties required for a suitable packaging film may not be found in one polymer and individual polymers are laminated to produce films of superior barrier properties. Examples of

polymers used in construction of barrier films are: Polyethylene, Polypropylene, Polyvinylidene chloride (PVDC), Ethylene vinylacetate (EVA), Ethylene vinyl alcohol (EVOH), and Metallised polyesters. Examples of high, medium and low barrier films are shown in Table 2.

Table 2. Oxygen permeation rates for packaging films expressed in different units*

Film	cc per 100 in ² /day	cc per m ² /day mm Hg	cc per m ² /day
PP/EVOH/PP	<0.001	<0.00001	0.01-0.02
Foil laminate (mylar/AL/poly)	<0.01	<0.00001	<0.01-0.1
PVDC	1	0.02	15
Acrylonitrile Polymer	1	0.02	15
PET (polyester)	4-6	0.08-0.13	60-100
PVC	10	0.2	150
HDPE	130	2.6	1980
PP	150	3.0	2280
LDPE	400-500	8-10	6000-7000

*For 1 mil flat (unless composite laminate) at 30°C and 50% RH

iii) Storage temperature

The effectiveness of MAP decreases as the storage temperature increases due to the fact that the solubility of CO₂ also decreases at higher temperatures. For respiring products, increasing the temperature also increases the rate of respiration, resulting in a decrease in shelf life. The effects of temperature abuse are particularly important from the standpoint of safety. Temperature abuse of MAP muscle foods may result in the rapid growth of both spoilage and pathogenic bacteria. The minimum reported temperature for *Salmonella* and *Escherichia coli* inoculated in ground meat grew equally well at 12.5 °C when the meat was packaged in low and high permeability film (Goepfert & Kim, 1975). *Staphylococcus aureus* can grow and produce enterotoxin under anaerobic conditions at 10°C or less while *Yersinia enterocolitica* has been reported to grow at temperatures as low as -2°C. Of major concern, with respect to safety of MAP fish, is the growth and toxin production by *Clostridium botulinum* type E (Palumbo, 1986). Proper refrigeration is therefore essential in order to assure the effectiveness of CO₂ as an anti microbial agent and to prevent potential growth of pathogenic organisms.

iv) Microbial contamination

The shelf life extension of fish under modified atmospheres is dependent on the initial

microbial load and the types of organisms. Higher initial counts will accelerate spoilage and shorten the shelf life of fish products. Furthermore, high numbers of aerobic bacteria will also consume rapidly headspace O_2 and may change electronegative potential of product and enhance the growth of anaerobic organism such as *Clostridium botulinum* type E.

Effect of MAP on the shelf life of fish

In vacuum-packed ice stored fish from temperate marine waters an increased development of TMA is seen while the shelf life is unaffected compared to aerobically stored fish. The number of *Pseudomonas* is reduced, but *S. putrefaciens*, which is capable of anaerobic respiration using TMAO, grows to 10^6 - 10^8 cfu/g (Gram *et al.*, 1987; Jorgensen *et al.*, 1988; Dalgaard *et al.*, 1993). Jorgensen *et al.* (1988) observed that vacuum-packed cod consisted some very large, almost yeast like cells and suggested that these were involved in spoilage. It was recently shown that these cells are heat sensitive *Photobacterium phosphoreum* (Dalgaard *et al.*, 1993).

Banks *et al.* (1980); Brown *et al.* (1980); Finne (1982); Layrisse and Matches (1984); Lannelongue *et al.* (1982) and Stenstrom (1985) reported that CO_2 has been shown to delay spoilage of fresh seafood by inhibiting psychrotrophic aerobic Gram negative bacteria. Gill and Molin (1991) attributed the high growth rate of lactic acid bacteria and *S. putrefaciens* under 40% CO_2 +30% O_2 +30% N_2 to the tolerance of these microorganism to CO_2 . Brown *et al.* (1980) noted an atmosphere of CO_2 inhibited microbial growth and extended sensory acceptance of salmon fillets. Ozogul *et al.* (2000a) reported lowest counts in MAP stored herring compared to those in air-stored herring. Ozogul *et al.* (2004) observed significant differences ($P<0.05$) between samples kept in air and in MAP. Dalgaard (1995); Gram and Huss (1996); Drosinos and Nychas (1996) suggested that *P. phosphoreum*, *S. putrefaciens*, lactic acid bacteria and *Brochothrix thermosphacta*, bacteria resistant to carbon dioxide are important for the spoilage of fresh fish stored under either VP or MAP. In similar studies Oberlender *et al.* (1983); Satham *et al.* (1985); Wang and Orgydzia (1986) reported that lactic acid bacteria and *Alteromonas* are the dominant organism in fish stored under MAP conditions. Satham *et al.* (1985); and Fletcher *et al.* (1988) reported that *B. thermosphacta* were the dominant organism in morwong fish and scallops when

these were treated with polyphosphates or potassium sorbate and then stored under 100% CO_2 . *S. putrefaciens* and some Vibrionaceae produce H_2S from the sulphur containing amino acid L- cysteine (Gram *et al.*, 1987; Stenstrom & Molin, 1990). In contrast, neither *Pseudomonas* nor *P. phosphoreum* produce significant amounts of H_2S . Thus, hydrogen sulphide, which is typical of spoiling iced cod stored aerobically, is not detected in spoiling CO_2 packed cod (Dalgaard *et al.*, 1993). Methylmercaptan (CH_3SH) and dimethylsulphide [$(CH_3)_2S$] are both formed from methionine (Herbert & Shewan, 1975).

Quality Changes of fish in MAP storage

i) Physical Changes : Composition of the headspace gas mixtures

When compared to some packs containing meat products with similar concentrations of CO_2 (McMullen & Stiles, 1991), changes in the composition of gas mixtures within MAP fish packs appear to be rapid. The proportion of CO_2 in the headspace falls as CO_2 dissolves in fish tissues (Strasline *et al.*, 1982 and Davis, 1990). As a consequence, proportions of the diluent components increase and concentrations of oxygen, often above the atmospheric levels to start with, increase further. Later, as the CO_2 solubilisation rate is overtaken by the rate of release caused by bacterial respiration, the curve reverses. Any chemical effects on the fish tissues will be affected by the amount of CO_2 , which dissolves, but as the contaminant flora is limited to the fish surfaces, much of the bacteriostatic effect is likely to be influenced by the residual atmosphere. Davis (1990), working with cod fillets packed in a gas mixture containing 40% CO_2 found that the G/P ratio had to be at least 2:1 for maximal benefit. For a given gas mixture, high gas/ fish ratio will present a very different chemical balance from a low ratio. Similarly, any given mass ratio of CO_2 to product will not necessarily confer the same benefits (or harm) when applied via different gas mixtures. Silva and White (1994) in modified atmosphere packed refrigerated channel cat fish observed CO_2 concentration fluctuated between 63% and 87% for higher CO_2 concentration packs and 16 to 33% in low CO_2 concentration packs. Samples stored under air had increased CO_2 . The increase in CO_2 in air packed fillets was the result of microbial respiration. Debevere and Boskou (1996) observed in MAP cod fillets a decrease in CO_2 because of diffusion in to the fish muscle.

Table 3. Shelf life of fish products packaged under MAP, vacuum and air

Type of fishery products	Storage temperature (°C)	Atmosphere CO ₂ :N ₂ :O ₂	Shelf life (days)	Reference
Cod (<i>G.morhua</i>) whole	2	100:0:0	10	Jensen <i>et al</i> (1980)
	2	60:40:0	10	
	2	40:60:0	9-10	
	2	Vacuum	8-9	
	2	Air	7	
Rockfish (<i>Sebastes</i> spp.) fillets	1.7	CA 80:20 air	13	Parkin <i>et al</i> (1981)
		Air	6	
Salmon, king (<i>Oncorhynchus tshawytsche</i>) fillets	2	Air	8	Stier <i>et al</i> (1981)
	4.4	60:15:25	12	
	4.4	Air	6	
	22.2	60:15:25	2	
	22.2	Air	1	
Cod fillets	0	40:30:30	12.5	Cann <i>et al</i> (1983)
	5	40:30:30	<7	
	10	40:30:30	3	
	0	Vacuum	9	
	5	Vacuum	<4	
	10	Vacuum	2	
	10	Vacuum	2	
Crayfish (<i>Pacifastacus leniusculus</i>) whole cooked	4	80:20air	21	Wang & Brown (1983)
	4	Air	14	
Swordfish (<i>Xiphias gladius</i>) steaks	2	Air	6*	Oberiender <i>et al</i> (1983) *Shelf life based on microbial data only
	2	CA 100:0:0	>22*	
	2	CA 70:0:30	>22*	
	2	CA 40:0:60	14*	
	2	CA 70:30:0	>22*	
	2	CA 40:60:0	20*	
	2	CA 40:60:0	20*	
Trout whole	0	60:40:0	8	Cann <i>et al</i> (1984)
	5	60:40:0	8	
	10	60:40:0	3.8	
	0	Vacuum	9	
	5	Vacuum	6.5	
	10	Vacuum	3.7	
	10	Vacuum	3.7	
Snapper (<i>Chrysophrys auratus</i>) fillets	3	100:0:0	6-8	Scott <i>et al</i> (1984) *No/medium O ₂ barrier mat **High oxygen barrier material
	3	Vacuum	3*	
	3	Vacuum	6**	
Cod (<i>Gadus morhua</i>) fillets	3	Air	3	Woyewods <i>et al</i> (1984)
	1	CA 60:40:0	12	
	1	60:40:0	12	
Salmon steaks	1	Air	9	Cann <i>et al</i> (1984)
	0	60:40:0	12.9	
	5	60:40:0	7.1	
	10	60:40:0	3.4	
	0	Vacuum	11.8	
	5	Vacuum	8	
	5	Vacuum	8	
Flounder (<i>Limanda ferrugina</i>) fillets	26	Air	2	Post <i>et al</i> (1985)
	26	Vacuum	2	
	26	0:100:0	4	
	26	100:0:0	1	
	12	Air	5	
	12	Vacuum	8	
	12	0:100:0	7	
	12	100:0:0	8	
	8	Air	5	
	8	Vacuum	7	
	8	0:100:0	4	
	8	100:0:0	10	
	8	100:0:0	10	
Shrimp, spotted (<i>Pandalus platyceros</i>) whole head on/off	0	CA100:0:0	>14	Matches & Layrisse (1985)
	0	Air	7	
	0	Air	7	
Cod (<i>G.morhua</i>) fillets	26	100:0:0	2-3	Post <i>et al.</i> (1985) *All packaging types at
	26	Other atmosphere*	2	

Table 3 contd.

	12	Air	6	26°C (air, vacuum, 100% Nitrogen, 90:8:2 and 65:31:4) except 100% CO ₂ had a shelf life of 2 days
	12	Vacuum	10	
	12	0:100:0	13	
	12	100:0:0	11	
	8	Air	06	
	8	Vacuum	16	
	8	0:100:0	17	
	8	100:0:0	23	
	8	90:8:2	17	
	8	65:31:4	16	
	8	100:0:0	40-53	
Cod fillets	4		11	
	0-2	Air	19	Gopal, 1985
		CAP(50% O ₂ +50%CO ₂)	5	
		Air	11	
		CAP(50% O ₂ +50%CO ₂)	2*	
Whiting (<i>Merluccius bilinearis</i>) fillets	5-7	All atmosph*	5	Post <i>et al</i> (1985)
	26	Air	9	*All packaging types at 26°C (air, vacuum, 100%N ₂ , 100%CO ₂ , 90:8:2 and 65:31:4) had a shelf life of 2 days
	12	Vacuum	9	
	12	0:100:0	12	
	12	100:0:0	4	
	3	Air	10	
	8	Vacuum	10	
	8	0:100:0	15	
	8	100:0:0	13	
	8	90:8:2	7	
	8	65:31:4	15	
	4	100:0:0	9	
Snapper (<i>C.auratus</i>) fillets	-1	40:60:0	9	Scott <i>et al.</i> (1986)
	-1	Air	18	
	-1	100:0:0	12-13	
Cod (<i>G.morhua</i>) whole/fillets	0	Air	20	Villemure <i>et al</i> (1986)
Trout (<i>Salmo gairdneri</i>) fillets	0	25:75:0	20	Barnett <i>et al</i> (1987)
	1.7	80:20:0	20*	*Treated with 2% potassium sorbate dip
	1.7	80:20:0	10	Fujii <i>et al</i> (1989)
	1.7	Air	4	
Sardines (<i>Sardinops melanostictus</i>)	5	80:20:0	4	
	5	20:80:0	4	
	5	Air	2	
Catla (<i>Catla catla</i>), fillets	0-4	Air	12	Gopal <i>et al</i> (1990)
	50%	20		
	O ₂ +50%CO ₂			
	20%	28		
	O ₂ +80%CO ₂			
Cod (<i>G.morhua</i>) fillets	0 ²	2:98:0	14	Dalgaard <i>et al</i> (1993)
	0	3:97:0	13	
	0	29:71:0	16	
	0	48:52:0	20	
	0	97:3:0	15-16	
Haddock (<i>Melanogrammus aeglefinus</i>) whole	0	40:30:30	10	Dhananjaya & Stroud (1994)
	0	Air	8	
	5	40:30:30	7	
	5	Air	7	
	10	40:30:30	4	
	10	Air	4	
Catfish, channel (<i>Catalurus punctatus</i>) fillet strips	2	80:20 air	28	Silva & White (1994)
Cod, blue (<i>Arapercis colias</i>) fillets	3	100:0:0	49	Penney <i>et al</i> (1994)
commercially smoked	3	Vacuum	14	
	3	Air	14	
	-1.5	100:0:0	113	
	-1.5	Vacuum	35	
	-1.5	Air	28	

Table 3 contd.

Herring (<i>Clupea harengus</i>) fillets	0	60:40:0	14	Dhananjaya & Stroud (1994)
Herring (<i>C.harengus</i>) whole	0	Air	12	
Hybrid striped bass (<i>Morone saxatilis</i> , <i>M.chrysops</i>) strips	0	60:40:0	14	Dhananjaya & Stroud (1994)
Tilapia (<i>Tilapia</i> spp.) fillets	0	Air	12	Handumrongkul & Silva (1994)
	2	60:34:6	13	
	2	Air	7	
	4	75:25:0	>25	Reddy <i>et al</i> (1995)
	8	75:25:0	13-16	
	16	75:25:0	9-13	
	4	Air	9-13	
	8	Air	6-9	
	16	Air	3-6	
Trout, rainbow (<i>Oncorhynchus mykiss</i>) fillets	2	20:80:0	6*	Randell <i>et al</i> (1995)
	2	20:80:0	9**	
	2	40:60:0	6*	*G/P ratio 0.4
	2	40:60:0	9**	**G/P ratio 1.0
	2	Vacuum	6	
Herring, Baltic fillets	2	20:80:0	3*	Randell <i>et al</i> (1995)
	2	20:80:0	3**	*volume of gas to product ratio 0.4
	2	40:60:0	6*	**volume of gas to product ratio 1
	2	40:60:0	8**	
	2	Vacuum	3	
Mackerel (<i>Scombrus scombrus</i> L) fillets	-2	100:0:0	>21	Hong <i>et al</i> (1996)
Sea bream (<i>Argyrozona argyrozona</i>)	0-4	Air	22	Gopal <i>et al</i> (1996)
		20%	40	
		O ₂ +80%CO ₂		
Salmon, atlantic (<i>Salmo salar</i>) slices	2	100:0:0	18	Pastoriza <i>et al</i> (1996)
Catfish fillets	16	Air	3	Reddy <i>et al</i> (1997a)
	16	75:25:0	4	
	16	Vacuum	3	
	8	Air	6	
	8	75:25:0	13	
	8	Vacuum	6	
	4	Air	13	
	4	75:25:0	38-40	
	4	Vacuum	20-24	
Salmon fillets	16	Air	4	Reddy <i>et al</i> (1997b)
	16	75:25:0	5-6	
	16	Vacuum	3	
	8	Air	13-17	
	8	75:25:0	20-24	
	8	Vacuum	>6<10	
	4	Air	24-27	
	4	75:25:0	55-62	
	4	Vacuum	34-38	
Hake (<i>Merluccius merluccius</i>) slices	2	50:45:5	14	Pastoriza <i>et al</i> (1998)
	2	50:45:5	16*	*5 min 5%NaCl dip
	2	Air	7-8	
Cod fillets	16	Air	3-4	Reddy <i>et al</i> (1999)
	16	75:25:0	6	
	16	Vacuum	3-4	
	8	Air	13-17	
	8	75:25:0	24-27	
	8	Vacuum	13	
	4	Air	20-24	
	4	75:25:0	55-60	
	4	Vacuum	24-27	
Salmon (<i>S.salar</i>) fillets	10	Vacuum	3	Randell <i>et al</i> (1999)
	2	60:40:0	17	
	2	40:60:0	17	
	2	Vacuum	17	
	2	Air	11	

Note: Initial atmosphere in percent CO₂:N₂:O₂. If mixtures of carbon dioxide and air is used this is reported as % CO₂: % air (e.g. 80:20 air), if initial atmosphere is maintained during storage either in controlled atmosphere tank or by reflushing, this is indicated with CA.

Reciprocally to the CO_2 content the proportion of O_2 increased due to respiration of bacteria and the proportion of O_2 decreased later on. Amanatidou *et al.* (2000), in fresh Atlantic salmon using 50% CO_2 + 50% O_2 observed increased O_2 concentration in the packs.

ii) Changes in thaw drip during storage

Exudate or drip in increased amounts is another consequence of treating fish with CO_2 . Normally, the small amount released from raw fillets is not a great problem but it becomes a limiting factor for some products in MAP. The problem may be solved by limiting the amount of CO_2 and by placing the fish on absorbent pads within the packs (Tiffiney & Mills, 1982; Cann, 1984). The lower water content of smoked products and fatty fish leaves them less vulnerable to further losses under 60% CO_2 than with raw white fish. The problems are aggravated at the lower storage temperature, perhaps a reflection of the effect on the solubility of CO_2 . Pastoriza *et al.* (1996) observed no marked effect of CO_2 on exudation of salmon slices stored in ice under CO_2 atmospheres. Dalgaard *et al.* (1993) observed an increase in drip loss during MAP storage of cod fillets, especially at high CO_2 concentrations. Pastoriza *et al.* (1998) observed increased exudation in air and MAP stored fish showed exudates values higher than control after 7 days of storage. Fey and Regenstein (1982) found increased drip losses for red hake, Chinook salmon and to a lesser extent, Sockeye salmon stored in 60% CO_2 , 21% O_2 and 19% N_2 compared to air packed product. Tiffiney and Mills (1982) found that packing in 100% CO_2 increased the rate and quantity of drip loss. In all cases the amount of drip loss of fish stored in high CO_2 concentrations was higher at 0°C than at 5°C.

iii) Changes in pH during storage

Apparent contradictions arise in considering almost every aspect of MAP fish including muscle pH. Fey and Regenstein (1982) reported little or no change whilst others have seen a decrease proportional to CO_2 concentration (Lannelongue *et al.*, 1982a, 1982b; Belleau & Simard, 1987). Fish muscle of relatively high post mortem pH can be expected to be more affected by a given amount of CO_2 than muscle of lower pH but this will be complicated by other variations in the chemical composition affecting the buffering capacity of the tissue. Additional

variation in reported measurement may also occur because of differences in method of measurements, mainly because of the gradients, which occur between the product surface and deeper tissue (Tiffiney & Mills, 1982). Initial dissolution of CO_2 (preceding formation of carbonic acid) and bacterial activity (which produces high pH waste products) are surface phenomena, and, therefore, the most rapid and extreme fluctuations in pH occur on the surface. The general pattern seems to be that after any initial fall in pH, surface pH rises whilst internal pH lags behind.

Pastoriza *et al.* (1996) observed an increase in pH in air stored samples throughout the storage period, with a value of 8.03 after 21 days; MAP stored samples showed lower pH values throughout storage. Dalgaard *et al.* (1993) reported a pH increase from 6.6 to 6.8 in low CO_2 concentrations, but it remained almost constant at 6.6 in high CO_2 concentrations. Debevere and Boskou (1996) observed counter effects of CO_2 on the pH increase by TVB production, resulting in a stabilization of the pH around 6.7. Pastoriza *et al.* (1996) observed slight increase of pH in the control during the low temperature storage of salmon slices whereas in MAP, an initial decrease of pH was observed followed by an increase after 6 days of storage. Layrisse and Matches (1984) indicated that CO_2 is rapidly dissolved or absorbed in MAP stored muscle and consequently, pH decreased. Parkin *et al.* (1981) and Lannelongue *et al.* (1982) observed that the decrease of pH could be proportional to the CO_2 content of the package. Banks *et al.* (1980) and Parkin *et al.* (1981) attributed the pH change due to conversion of CO_2 to carbonic acid at the fish muscle surface. Pastoriza *et al.* (1998) observed a pH value of 7.74 in control samples of hake slices after 2 weeks iced storage; fish stored under MAP conditions showed pH values lower than the control after ten days storage. Lopez- Galvez *et al.* (1998) observed a pH increase in refrigerated storage of air packaged sole and only a slight increase was observed in samples stored in 20% CO_2 and no change in pH were observed when samples were stored in CO_2 /air (40/60) (v/v). Therefore in general, the dissociation of carbonic acid in fish flesh results in a slight drop in pH. Both the buffering capacity of the fish proteins and the composition of the spoilage flora determine the magnitude of pH change (Cutting, 1953). However Barnett *et al.*

(1978) found no significant change in pH of salmon flesh stored in 90% CO₂. The extent to which pH decreases is proportional to the concentration of CO₂ in the atmosphere (Lannelongue *et al.*, 1982a; Tiffiney & Mills, 1982). Parkin *et al.* (1981) noted a drop in pH of rockfish from pH 6.7 to 6.3, which was maintained throughout the storage period. However Fey and Regenstein (1982) found that after an initial decrease, pH increased and after 27 days storage in a CO₂ enriched environment had reached a level similar to its initial pH.

iv) Changes in Biochemical parameters

a) Changes in Trimethylamine Nitrogen (TMA-N) content

Dalgaard *et al.* (1993) observed that the level of TMA was typically around 10-15 mg TMA-N/100g in aerobically stored fresh fish when rejected by sensory panels. Ababouch *et al.* (1996) observed that the limit of acceptability for sardines was found to be 5-10 mg TMA-N/100g of samples according to comparison of sensory and chemical data. Pastoriza *et al.* (1996) observed very low TMA values for salmon slices. Such low values can be consequences of a very low bacterial load leading to a very much-reduced enzymic deterioration of fish muscle (Gerdes *et al.*, 1989). Ozogul *et al.* (2000b) reported a value of 13.5 mg in herring kept in MAP (CO₂:N₂ 60/40) for 10 days at 2±2°C. Cann *et al.* (1983) reported a TMA value of 10.4 mg/100g in herring fillets stored in MAP with the same gas mixture for 11 days at 0°C. Ozogul *et al.* (2004) reported that significant differences were discovered between sardines held under air and vacuum pack and MAP after 4 days of storage. The concentration of TMA in numerous fatty fishes never reached the limit of 5 mg / 100g although the rejection limit in flesh is usually 5-10 mg /100g.

b) Changes in Total Volatile Base Nitrogen (TVB-N)

The more rapid increase of TVB-N at high microbial numbers indicated the stage of substantial spoilage of the fish. Ababouch *et al.* (1996) reported that the limit of acceptability for sardines was 25-35 mg /100g of flesh. Ozogul *et al.* (2004) reported TVB-N content of sardines stored in air, vacuum pack, and MAP at 4°C. Initial TVB-N value was 5 mg/100g for sardines stored under air. The release of total volatile bases increased up to 15 mg/100g for sardines in

VP and 17 mg/100g in MAP at the last day of sensory acceptability for each storage conditions. Pastoriza *et al.* (1996) observed salmon slices stored under CO₂ showed a much slower increase, with values much lower than those considered as limit of acceptability after 20 days storage, and similar to those of control after 10 days storage. They considered 30 mg TVB-N/100g as limit of acceptability. Pastoriza *et al.* (1998) observed a reduction in TVB-N values during MAP storage of hake slices, assuming 35 mg TVB-N/100g as the limit of acceptability for consumption of fish.

c) Changes in Thio Barbituric Acid Value (TBA)

Boyd *et al.* (1992) reported very low TBA values in pond raised striped bass and Kyrana and Longovis, (2002) and Papadopoulos *et al.* (2003) also reported very low TBA values for farmed sea bream stored under air in chilled conditions. Pastoriza *et al.* (1996) found increased TBA values in CO₂ modified atmospheres compared to air stored samples in salmon slices and hake. Amanatidou *et al.* (2000) found that salmon samples stored under modified atmospheres and kept under chilled conditions, with TBA value above 1.9 were characterized by unacceptable organoleptic characteristic

d) Changes in K value during storage:

The initial quality loss in fish is primarily caused by autolytic changes. Of particular importance in this respect is the degradation of nucleotides (ATP- related compounds), which is caused by autolytic enzymes.

Boyle *et al.* (1991) studied the adenine nucleotide degradation in modified atmosphere chill stored whitefish and rainbow trout. The results indicated CO₂ atmospheres did not alter the K values compared to those observed for aerobically held fish. However CO₂ atmospheres caused decrease in hypoxanthine concentrations compared to aerobically held samples. Huynh *et al.* (1992) found the same results with sockeye salmon and herring. Randell *et al.* (1995) studied the effect of G/P ratio and CO₂ concentration on the shelf life of MAP fish. The results indicated CO₂ concentration did not affect the K values of Rainbow trout. Reddy *et al.* (1997) also found that at 4, 8, and 16°C storage K values of MAP fillets of cat fish increased gradually during early storage time and decreased towards end of

storage period indicating no relationship between sensory spoilage and K value. Lopez-Galvez *et al.* (1998) with sole fillets reported no effect of the atmosphere on K values. Ozogul *et al.* (2000a) studied the effects of modified atmospheres on K values of herring stored at 2°C. They observed 60% CO₂ atmospheres showed lower K values compared to those observed for aerobically stored fish, in addition CO₂ decreased the formation of Hypoxanthine compared to aerobically and vacuum held fish. Ozogul *et al.* (2004) studied the effects of modified atmospheres on K values of sardines at 4°C. The lowest increase in K value was observed for sardines stored in MAP, which was possibly influenced by the presence of CO₂.

v) Changes in colour during storage

Discoloration can occur via bleaching action of cut surfaces (Cann, 1984) probably due to low pH precipitation of sarcoplasmic proteins (Statham & Bremner, 1989). There is also risk of discoloration of haem pigments (Przybylski *et al.*, 1989). Brown *et al.* (1980) demonstrated that inclusion of 1% CO in the gas mix could help retain a red color, unlike Fey and Regenstein (1982) who observed a negative effect. Inclusion of O₂ as a means of maintaining more attractive red color of haem pigments is recommended for red meats but high concentrations are necessary. With fatty fish, Tiffiney and Mills (1982) found that the fresh appearance was retained for longer in O₂- free packs. High concentrations of CO₂ cause discoloration of meats, especially red meats. The degree of redness depends upon the amount of pigment present in the meat and the availability of O₂. Myoglobin can be oxidized in an atmosphere rich in CO₂ to form metmyoglobin, thus reducing the amount of redness to a light brown discoloration. Since fish are low in myoglobin, the discoloration induced by elevated concentrations of CO₂ is not as noticeable as in red meat. Colour changes in fish include graying of the cornea (Coyne, 1933), bleaching of the skin and damage to bloom at high CO₂ concentrations (>60%) (Stansby & Griffiths, 1935). Goodfellow (1982) found it necessary to restrict CO₂ levels to 25% or less to prevent discoloration problems.

Silva and White (1994) in modified atmosphere packaged refrigerated channel cat fish observed increased Hunter L* values in all packaging environments over time. Hunter a*

value had an initial decrease in fish held in high CO₂ concentration but increased as time progressed. Hunter b* values showed no differences among any treatments. Pastoriza *et al.* (1996) observed increased Hunter L* and b* values and decreased Hunter a* value in both air and MAP stored samples during the storage period. Amanatidou *et al.* (2000) in modified atmosphere packaging of Atlantic salmon observed a product with lightness above the threshold value of 70 or red colour below 13 is unacceptable.

vi) Changes in texture during storage

Texture of the fish is very important, since a poor texture can result in bad visual appearance of the product. Springiness can be used to simulate finger feel of the raw fillets/ portions, and shear value as an index of tenderness/firmness of the raw fish. The most common types of measurements are based upon rheological principles of shear strength, puncture and compression.

Amanatidou *et al.* (2000) observed a lower cutting strength in MA packed fresh Atlantic salmon, indicating an increase in the softness upon storage. Textural characteristics were retarded in MAP compared to control. Fagan *et al.* (2004) tested effects of MAP with freeze chilling on some quality parameters of raw whiting, mackerel and salmon portions. The results indicated that storage time had no effect on the springiness of whiting, mackerel or salmon portions. MAP had no effect on the shear values of whiting fillets or salmon portions. However mackerel fillets stored in 100% CO₂ had higher shear values than samples from other treatments.

Effect of MAP on microbiological hazards

Fish stored under MAP may not increase the risks from *Salmonella*, *Staphylococcus*, *C.perfringens*, *Yersinia*, *Campylobacter*, *Vibrio parahaemolyticus* and *Enterococcus* above those expected for air stored products (Silliker & Wolfe, 1980; Reddy *et al.*, 1992). Growth of *L.monocytogenes* is apparently not significantly affected by MAP (Rocount & Cossart, 1997). Pothuri *et al.* (1996) observed an 8 days extension of the lag phase of *L.monocytogenes* in samples of crayfish tail meat treated with 1% lactic acid and packaged under MA conditions (75% CO₂:10% O₂, 15% N₂) as compared with that in air or vacuum packaging. 100% CO₂ delayed, but did not inhibit the growth of *L.monocytogenes* as

compared with air in inoculated raw and cooked seafood nuggets (Lyver *et al.*, 1998) and was able to reach 10^6 cfu/g before spoilage occurred in all packaging methods. Packaging under enhanced CO_2 atmospheres will delay the growth of *L.monocytogenes*, but the MAP and chilled storage alone are not sufficient to control the growth of the pathogen in some products. To ensure the safety, especially for value added sea food products with extended shelf life, additional hurdles must be applied. Modified atmosphere packaging conditions create an environment, which supports the growth of psychrotrophic, anaerobic bacterial pathogens, including non-proteolytic types of *C. botulinum*.

Genigeorgis (1985); Baker and Genigeorgis (1990); Stammen *et al.* (1990); Reddy *et al.* (1992) have reviewed the risks from botulism in MAP fish. Post *et al.* (1985); Garcia and Genigeorgis (1987); Taylor *et al.* (1990) reported that toxin has been detected in MAP fish prior to the product being considered spoiled. Cann *et al.* (1984); Reddy *et al.* (1996; 1997a; b; 1999) and Cai *et al.* (1997) reported that MA or vacuum packed fish spoiled prior to or in coincidence with toxin production. Gibson *et al.* (2000) demonstrated that 100% CO_2 could have an inhibitory effect on the growth of *C.botulinum* at chill temperatures, and an inhibitory effect was observed when combining 100% CO_2 with increased NaCl level and decreased pH level. Reddy *et al.* (1999) reported the fat content might influence the margin of safety of MA packaged aqua cultured fresh fillets during storage. The safety margin being less for fatty fish when compared with lean fish. Lilly and Kautter (1990) reported in a study of 1074 test samples of commercial, vacuum packaged fresh fish, none of the marginally organoleptic acceptable samples was positive for *C. botulinum* after 12 days at 12°C . The authors concluded that either the fish did not contain *C.botulinum* spores, or the spores were unable to grow out and produce toxin before the spoilage made the product marginally unacceptable. However, an increased use of modified atmosphere and other minimal processing technologies combined with improper cold chains and abuse of temperature may represent an increased risk of botulism.

References

- Ababouch, L.H., Souibri, L., Rhaliby, K., Ouadhi, O., Battal, M., Busta, F.F. (1996) Quality changes in sardines (*Sardina pilchardus*) stored in ice and at ambient and temperatures. *Food Microbiol.* **13**, pp 123-132.
- Abbey, J.R. (1970) Gas packaging with carbon dioxide. *Food Manuf.* **45**, pp 37-40.
- Amanatidou, A., Schluter, O., Lemkau, K., Gorris, L.G.M., Smid E.J., Knorr, D. (2000) Effect of combined application of high pressure treatment and modified atmosphere on the shelf life of fresh Atlantic salmon. *Innovative Food Science and Emerging Technologies* **1**, pp 87-98.
- Anon (1977) Controlled atmosphere pack for meat products. *Food Manuf.* **52**, pp 90-101.
- Anon (1985) Guidelines for the Handling of Fish Packed in a Controlled atmosphere Sea Fish Industry Authority, Edinburgh.
- Anon (1967) Eskimo double – new pea pack, cod fillets. *Food Manuf.* **41**, pp 35-36.
- Baker, D.A. and Genigeorgis, C. (1990) Predicting the safe storage of fresh fish under modified atmospheres with respect to *Clostridium botulinum* toxigenesis by modeling length of the lag phase growth. *J. Food Prot.* **53**, pp 131-140.
- Banks, H., Nickelson, II, R. and Finne, G. (1980) Shelf life studies on carbon dioxide packed finfish from the Gulf of Mexico. *J. Food Sci.* **45**, pp 157-162.
- Barnett, H.J. Stone, F.E., Roberts, G.C., Hunter, P.J., Nelson, R.W. and Kwok, J.A. (1978) A study in the use of a high concentration of CO_2 in a modified atmosphere to preserve fresh salmon. *Mar. Fish. Rev.* **44**: pp 7-9.
- Barnett, H.J., Conrad, J.W. and Nelson, R.W. (1987). Use of laminated high-and low density polythene flexible packaging to store trout (*Salmo gairdneri*) in modified atmosphere. *J. Food Prot.* **50** pp 645-651.
- Barnett, H.J., Stone, F.E., Roberts, G.C., Hunter, P.J., Nelson, R.W. and Kwok, J. (1982). A study in the use of a high concentration of CO_2 in a modified atmosphere to preserve fresh salmon. *Mar. Fish. Rev.* **44** pp 7-11.
- Belleau, L. and Simard, R.E. (1987) Effets d'Atmosphere de dioxyde de carbone et d'azote sur des filets de poisson. *Sciences des Aliments*, **7**, pp 433-446.

- Boskou, G. and Debevere, J.J. (1997) Reduction of trimethyl amine oxide by *Shewanella* spp. under modified atmospheres in vitro. *Food Microbiol.* **14**, pp 543-553.
- Boyd, A.C., Green, D.P., and Le Pors, L.A. (1992) Quality changes of pond raised hybrid bass during chill pack and refrigerated storage. *J. Food Sci.* **57**: pp 59-62
- Boyle, J.L., Linsay, R.C., Stuijber, D.A. (1991) Adenine Nucleotide degradation in Modified Atmosphere chill stored fresh fish. *J. Food Sci.* **56** pp 1267-1269.
- Bremner, A., Olley, J., Statham, J.A. and Vail, A.M.A. (1988) Nucleotide catabolism: Influence on the storage life of tropical species of fish from the north west shelf of Australia. *J. Food Sci.* **53**, pp 6-11.
- Brody, A.L. (1989). Modified atmosphere packaging of seafoods. In: *Controlled/ Modified Atmosphere/Vacuum packaging of foods* (ed. A.L. Brody). pp. 59-65. Trumbull, CT: Food & Nutrition Press.
- Brown, W. (1922) On the germination and growth of fungi at various temperatures and in various concentrations of oxygen and carbon dioxide. *Ann. Bot.* **36**: pp 257-265.
- Brown, W.D., Albright, M., Watts, D.A., Heyer, B., Spruce, B. and Price, R.J. (1980) Modified atmosphere storage of rock fish and silver salmon. *J. Food Sci.* **45**: pp 93-101
- Bullard, F.A. and Collins, J. (1978) Physical and Chemical changes of pink shrimp, (*pandalus borealis*) held in carbon dioxide modified refrigerated seawater compared with pink shrimp held in ice. *Fish. Bull.* **76**: pp 73-81.
- Cai, P., Harrison, M.A., Huang, Y.W. and Silva, J.L. (1997) Toxin production by *Clostridium botulinum* type E in packaged Channel catfish. *J. Food Prot.* **60**, pp 1358-1363.
- Cann, D.C., Smith, G.L. and Houston, N.C. (1983) *Further studies on Marine fish Stored under Modified Atmosphere Packaging*. Ministry of Agriculture, Fisheries and Food, Torry Research Station, Aberdeen, Scotland.
- Cann, D.C. (1984). Packing fish in a modified atmosphere, Torry Advisory Note No. 88, Ministry of Agriculture, Fisheries and Food, Torry Research Station, Aberdeen.
- Cann, D.C., Houston, N.C., Taylor, L.Y., Smith, G.L., Thomson, A.B. and Craig, A. (1984) *Studies of Salmonids Packed and Stored under a Modified Atmosphere*. Aberdeen: Torry Research Station.
- Cann, D.C., Houston, N.C., Taylor, L.Y., Stroud, G., Early, J. and Smith, G.L. (1985). *Studies of shellfish packed and stored under a modified atmosphere*, Ministry of Agriculture, Fisheries and Food, Torry Research Station, Aberdeen.
- Charm, S.E., Longmaid III, H.E. and Carver, J. (1977) A simple system for extending refrigerated, nonfrozen preservation of biological material using pressure. *Cryobiology*, **14**, pp 625-636
- Coyne, F. P. (1932) The effect of carbon dioxide on bacterial growth with special reference to the preservation of fish. Part I, *J. Soc. Chem. Ind.* **51**, pp 119T-121T.
- Coyne, F.P. (1933) The effect of carbon dioxide on bacterial growth with special reference to the preservation of fish. Part II- Gas storage of fresh fish. *J. Soc. Chem. Ind.* **52**: 19T-24T.
- Cutting, C.L. (1953) Changes in pH and buffering capacity of fish during spoilage. *J. Sci. Food Agric.* **12**: pp 597
- Dalgaard, P. (1995) Qualitative and quantitative characterisation of spoilage bacteria from packed fish. *Int. J. Food Microbiol.* **26**, pp 319-388.
- Dalgaard, P. (1995a) Modelling of microbial activity and prediction of shelf life for packed fresh fish. *Int. J. Food Microbiol.* **26**, 305-317.
- Dalgaard, P. (1995b) The effect of anaerobic conditions and carbon dioxide. In: H.H. Huss (editor), *Quality and Quality changes in Fresh Fish*. FAO Fisheries Technical Paper No. 348, FAO, Rome, Italy.
- Dalgaard, P., Gram, L. and Huss, H.H. (1993) Spoilage and shelf life of cod fillets packed in vacuum or modified atmospheres. *Int. J. Food Microbiol.* **19**, pp 283-294.
- Daniels, J.A., Krishnamurth, R. and Rizvi, S.S.H. (1985) A review of effects of CO₂ on microbial growth and food quality. *J. Food Prot.* **48**, pp 532-537.

- Davis, A.R. and Slade, A. (1995) Fate *Aeromonas* and *Yersinia* on modified atmosphere packaged (MAP) cod and trout. *Lett. Appl. Microbiol.* **21**, pp 354-358.
- Davis, H.K. (1990) Some effects of modified atmosphere packaging gases on fish and chemical tests for spoilage, in *Chilling and Freezing of New Fish Products*, Int. Inst. Refrig., Commission C2, Aberdeen 1990-3, pp. 201-207.
- Debevere, J. and Bouskou, G. (1996) Effect of modified atmosphere packaging on the TVB/TMA producing micro flora of cod fillets. *Int. J. Food Microbiol.* **31**, pp 221-229.
- Devlieghere, F., Debevere, J. and Van Impe, J. (1998a) Effect of dissolved carbon dioxide and temperature on the growth of *Lactobacillus sake* in modified atmospheres. *Int. J. Food Microbiol.* **41**, 231-238.
- Devlieghere, F., Debevere, J. and Van Impe, J. (1998b) Concentration of carbon dioxide in the water phase as a parameter to model the effect of a modified atmosphere on micro-organisms. *Int. J. Food Microbiol.* **43**, 105-113.
- Dhananjaya, S., and Stroud, J. (1994) Chemical and sensory changes in haddock and herring stored under modified atmosphere. *Int. J. Food Sci. and Technol.* **29**, pp 575-583.
- Douglas (1970) Laminates for vacuum and gas packaging. *Food Manuf.* **45**(4), pp 46-49.
- Drosinos, E.H. and Nychas, G.-J.E. (1996) *Brochothrix thermosphacta*, a dominant microorganism in Mediterranean fresh fish (*Sparus aurata*), stored under modified atmosphere. *Italian J. Food Sci.* **8**, pp 323-329.
- Enfors, S.O. and Molin, G. (1980) Effect of high concentrations of carbon dioxide on growth rate of *Pseudomonas fragi*, *Bacillus cereus* and *Streptococcus cremoris*. *J. Appl. Bacteriol.* **48**: pp 409-420.
- Fagan, J.D., Gormley, T.R. and Mhuirheartaigh, M.M. Ui. (2004) Effect of modified atmosphere packaging with freeze chilling on some quality parameters of raw whiting, mackerel and salmon portions. *Innovative Food Science and Emerging Technologies*. **2**, pp 205-214.
- Farber, J.M. (1991) Microbiological aspects of modified atmosphere-packaging technology - a review. *J. Food Prot.* **54**, pp 58-70.
- Fey, M.S. and Regenstein, J.M. (1982) Extending shelf life of fresh wet red hake and salmon using CO₂-O₂ modified atmosphere and potassium sorbate ice at 1°C. *J. Food Sci.* **47**: pp 1048-1054.
- Finne, G. (1982) Modified and controlled atmosphere storage of muscle foods. *Food Technol.* **36**, pp 128-133.
- Fletcher, G.C., Murrel, W.G., Statham, J.A., Stewart, B.J. and Bremner, H.A. (1988) Packaging of scallops with sorbate: an assessment of the hazard from *Clostridium botulinum*. *J. Food Sci.* **53**, pp 349-358.
- Fuji, T., Hirayama, M., Okuzumi, M., Yasuda, M., Nishito, H. and Yokotama, M. (1989). Shelf life studies on fresh sardine packaged with carbon dioxide-nitrogen gas mixture. *Nippon Suisan gaikkashi*, **55**: pp 1971-75
- Garcia, G. and Genigeorgis, C. (1987) Quantitative evaluation of *Clostridium botulinum* nonproteolytic types B, E, and F growth risk in fresh salmon tissue homogenates stored under modified atmospheres. *J. Food Prot.*, **50**, pp 390-397.
- Genigeorgis, C.A. (1985) Microbiological and safety implications of the use of modified atmospheres to extend the storage life of fresh meat and fish. *Int. J. Food Microbiol.* **1**, pp 237-251.
- Gerdes, D.L., Hoffstein, J.J., Finerty, M.W. and Grodner, R.M. (1989) The effects of elevated CO₂ atmospheres on the shelf life of freshwater crawfish (*Procambaris clarkii*) tail meat. *Lebensm Wiss Technol.* **22**, 315-318.
- Gill, C.O. and Penny, N. (1988) The effect of the initial gas volume to meat weight on the storage life of chilled beef packed under carbon dioxide. *Meat Sci.* **22**, 53-63.
- Gill, C.O. and Tan, K.H. (1980) Effect of carbon dioxide on growth of meat spoilage bacteria. *Appl. Environ. Microbiol.* **39**:317.
- Gill, C.O., and Molin, G. (1991) Modified atmospheres and vacuum packaging. In: N.J. Russell, and G.W. Gould, *Food preservatives* (pp. 172-199). Glasgow and London: Blackie Scientific.

- Gillespie, N.C. and MacRae, I.C. (1975) The bacterial flora on some Queensland fish and its ability to cause spoilage. *J. Appl. Bacteriol.* 39, 91-100.
- Goepfert, J.M. and Kim, H.V. (1975) Behaviour of selected foodborne pathogens in raw ground beef. *J. Milk Food Technol.* 38: 443.
- Goodfellow, S.J. (1982) An overview of controlled atmosphere packaging of foods. *Matin. R.E.*, (ed.) Proceedings of the First National Conference on modified and controlled atmosphere packaging of seafood products. National Fisheries Institute. 250.
- Gopal, T.K.S., Nambiar, V.N., Battacharya, S.K., Joseph, J. and Prabhu, P.V. (1990). Modified Atmosphere storage of freshwater fish fillets of *Catla catla*. *Proc. International Institute of Refrigeration on Chilling and Freezing of new fish proucts.* Aberdeen, U.K., pp. 143-146.
- Gopal, T.K.S. (1985). Studies on the storage life of cod fillets using controlled atmosphere packaging. *Proc. Symposium on Recent Developments in Food Packaging*, AFSTI, India, pp. 86-89.
- Gopal, T.K.S., Nambiar, V.N., Joseph, J., Lalitha, K.V., Prabhu, P.V. (1996). Modified atmosphere storage of sea bream fillets from sea bream (*Argyroxona argyrozona*). *Proc. Workshop on Scientific results of FORV Sagar Sampada*, DOD, New Delhi. pp. 513-519
- Gorczyca, E. and Pek Poh Len (1985) Mesophilic spoilage of bay trout (*Arripis trutta*), bream (*Acanthopagrus butchri*) and mullet (*Aldrichetta forsteri*). In: A. Reilly (editor), *Spoilage of Tropical Fish and product Development*, FAO Fish. Rep. 317 Suppl. FAO, Rome, Italy, pp. 123-132.
- Gormley, T.R. and Zeuthen, P. (eds.) (1990) *Chilled Foods: the Ongoing Debate*, edited reports on chilled food workshops held as part of the Commission of the European Communities, 1984-1988 COST91 bis SG 3 programme on chilling, chilled storage and distribution of food, Elsevier Science Pub. Ltd.
- Gram, L. and Huss, H.H. (1996) Microbiological spoilage of fish and fish products. *Int. J. Food Microbiol.* 33, pp 121-137.
- Gram, L. and Melchiorson, J. (1996) Interaction of two spoilage bacteria, *Shewanella putrefaciens* and *Pseudomonas* sp. In fish model systems. *J. Appl. Bacteriol.* 80, pp 589-595.
- Gram, L., Trolle, G. and Huss, H.H. (1987) Detection of specific spoilage bacteria from fish stored at low 0°C and high 20°C temperatures. *Int. J. Food Microbiol.* 4, pp 65-72.
- Gram, L., Wedell-Neergaard, C. and Huss, H.H. (1990) The bacteriology of fresh and spoiling Lake Victorian Nile perch (*Lates niloticus*). *Int. J. Food Microbiol.* 10, pp 303-316.
- Haard, N.F. and Lee, Y.Z. (1982) Hypobaric storage of Atlantic salmon in carbon dioxide atmosphere. *Can. Inst. Food Sci. Technol. J.* 15, pp 68-75
- Haines, R.B. (1932) The influence of carbon dioxide on the rate of multiplication of certain bacteria as judged by viable counts. *J. Soc. Chem Ind. (London)* 52, pp 13-21.
- Handumrongkul, C. and Silva, J.L. (1994). Aerobic counts, colour, and adenine nucleotide changes in CO₂ packed refrigerated striped bass strips. *J. Food Sci.*, 59, pp 67-69.
- Herbert, R.A. and Shewan, J.M. (1975) Precursors of volatile sulphides in spoiling North Sea cod (*Gadus morhua*). *J. Sci. Food Agric.* 26, pp 1195-1202
- Hiltz, D.F., Lall, B.S., Lemon, D.W. and Dyer, W.J. (1976) Deteriorative changes during frozen storage in fillets and minced flesh of silver hake processed from round fish held in ice and refrigerated sea water. *J. Fish. Res. Board Can.* 33, pp 2560-2565.
- Hintlian, C.B. and Hotchkiss, J.H. (1986) The safety of modified atmosphere packaging: A review. *Food Technol.* 50, pp 70-80
- Hjorth - Hansen, S. (1933) Investigations relating to the cooking and handling of fresh shrimp. *Annu. Rep. Norw. Fish.* 3, pp 1-21
- Ho, C.S., Smith, M.D. and Shanahan, J.F. (1987) Carbon dioxide transfer in biochemical reactors. *Adv. Biochem. Engng.* 35, pp 83-125.
- Hong, L., Leblanc, E.L., Hawrysh, Z.J. and Hardin, R.T. (1996) Quality of Atlantic

- mackerel (*Scomber scombrus* L.) fillets during modified atmosphere storage. *J. Food Sci.* **61**, pp 646-651.
- Huynh, M.D., Mackey, R. and Gawley R. (1992) Freshness assessment of Pacific fish species using K value. In: *Seafood Science and Technology* (edited by E.Graham Bligh) pp. 258-268. Oxford: Fishing News (Books) Limited.
- Jensen, M.H., Petersen, A.R. and Jepsen, A. (1980) Storage of chilled cod under vacuum at various concentrations of carbon dioxide. Connell, J.J., ed. *Advances in Fish Science and Technology*. Torry Research Station Jubilee conference. Farnham : Fishing News p.294.
- Jorgensen, B.R. and Huss, H.H. (1989) Growth and activity of *Shewanella putrefaciens* isolated from spoiling fish. *Int. J. Food Microbiol.* **9**, pp 51-62.
- Jorgensen, B.R., Gibson, D.M. and Huss H.H. (1988) Microbiological quality and shelflife prediction of chilled fish. *Int. J. Food Microbiol.* **6**, pp 295-307.
- Kiesvaara, M., Hattula, T. and Karppinen, S. (1990) Development of methods used in the quality classification of fish. Technical Research Centre of Finland Research Notes, 1193.
- Killefer, D.H. (1930) Carbon dioxide preservation of meat and fish. *Ind. Eng. Chem.* **22**, pp 140-150
- Kimber, A. (1984) Update on the gas flush pack. *Food Manuf.* **59**, pp 22-25.
- King, A.D. Jr. and Nagel, C.W. (1967) Growth inhibition of a *Pseudomonas* by carbon dioxide. *J. Food Sci.* **32**, pp 575-585
- Kyrana, V.R. and Lougovois, V.P. (2002) Sensory, chemical, microbiological assessment of farm raised European seabass (*Dicentrarchus labrax*) stored in melting ice. *Int. J. Food Sci. Technol.* **37**, pp 319-328.
- Lannelongue, M., Finne, G., Hanna, M.O., Nickelson II, R. and Vanderzant, C. (1982a) Microbiological and chemical changes during storage of swordfish (*Xiphias gladius*) steaks in retail packages containing CO₂ enriched atmospheres. *J. Food Prot.* **45**, pp 1197-1203.
- Lannelongue, M., Finne, G., Hanna, M.O., Nickelson, R. and Vanderzant, G. (1982b) Storage characteristics of brown shrimp (*Panaeus aztecus*) stored in retail packages containing CO₂ enriched atmospheres. *J. Food Sci.* **47**, pp 911-918.
- Layrisse, M.E. and Matches, J.R. (1984) Microbiological and chemical changes of spotted shrimp stored under modified atmosphere. *J. Food Prot.* **47**, pp 453-458.
- Leblanc, R.J. and Leblance, E.L. (1992) The effect of super chilling with CO₂ snow on the quality of commercially processed cod (*Gadus morhua*) and pollack (*Pollachius virens*) fillets, in Quality Assurance in the Fish Industry, Proc. of the Technological Laboratory Dimond Jubilee Conf., Lyngby, 1991.
- Leistner, L. (1992) Food presevation by combined methods. *Food Res. Int.* **25**, pp 151-158.
- Lilly, T. and Kautter, D.A. (1990) Outgrowth of naturally occurring *Clostridium botulinum* in vacuum packed fresh fish. *J. Assoc. Off. Anal. Chem.* **73**, pp 211-212.
- Lima dos Santos, C.A.M. (1978) Bacteriological spoilage of Iced Amazonian Freshwater Catfish (*Brachyplatistoma vaillanti valenciennes*). M.Sc. Thesis. Loughborough University of Technology, England.
- Lopez-Galvez, D.E., Hoz, L. de la and Ordenez, J.A. (1995) Effect of carbon dioxide and oxygen enriched atmospheres on microbiological and chemical changes in refrigerated tuna (*Thunnus alalunga*) steaks. *J. Agric. Food Chem.* **43**, pp 483-490.
- Lopez-Galvez, D.E., Hoz, L. de la., Blanco, M. and Ordenez, J.A. (1998) Refrigerated storage (2°C) of sole (*Solea solea*) fillets under CO₂ enriched atmospheres. *J. Agric. Food Chem.* **46**, 1143-1149.
- Lyver, A., Smith, J.P., Tarte, I., farber, J.M. and Nattress, F.M. (1998). Challenge studies with *Listeria monocytogenes* in a value added seafood product stored
- Lyver, A., Smith, J.P., Tarte, I., Furber, J.M. & Nattress, F.M. (1998). Challenge studies with *Listeria monocytogenes* in a value added seafood product stored under modified atmospheres. *Food Microbiology*, **15**, pp 379-389

- Martin, R.E. (ed.) (1981) Proceedings First National Conference on Modified and Controlled Atmosphere Packaging of Seafood Products, San Antonio, The National Fisheries Institute, Washington DC.
- Martin, R.E. (ed.) (1982) *Proceedings First National Conference on Seafood Packaging and Shipping*, Washington and Seattle, The National Fisheries Institute, Washington DC.
- Matches, J.R. and Layrisse, M.E. (1985) Controlled atmosphere storage of spotted shrimp (*Pandalus platyceros*). *J. Food Prot.* **48**, pp 709-711.
- McMullen, L.M. and Stiles, M.E. (1991) Changes in microbial parameters and gas composition during modified atmosphere storage of fresh pork loin cuts. *J. Food Prot.* **54**, pp 778-783.
- Molin, G., and Stenstrom, I.M. (1984) Effect of temperature on the microbial flora of herring fillets stored in air or carbon dioxide. *J. Appl. Bacteriol.* **56**, pp 275-282.
- Moran, T., Smith, E.C. and Tomkins, R.G. (1932) The inhibition of mould growth on meat by carbon dioxide. *J. Soc. Chem. Ind.* **51**, pp 114-120.
- Nelson, R.W. and Barnett, H.J. (1971) Fish preservation in refrigerated sea water modified with carbon dioxide, in *Proceedings XIII International Congress Refrigeration Vol. 3*, Washington DC, pp. 57-64.
- Nelson, R.W. and Tretsven, W.I. (1977) Storage of Pacific salmon in controlled atmosphere, in *Proceedings XIV International Congress Refrigeration* Moscow, USSR, 1975, Vol. 3, Int. Inst. Ref., pp. 736-743.
- Oberlender, V., Hanna, M.O., Miget, R., Vanderzant, C. and Finne, G. (1983) Storage characteristics of fresh swordfish steaks stored in carbon dioxide enriched controlled (flow through) atmospheres. *J. Food Prot.* **46**, pp 434-440.
- Ogrydziak, D.M. and Brown, W.D. (1982) Temperature effects in modified atmosphere storage of seafoods. *Food Technol.* **36**, pp 86-96.
- Ozogul, F., Polata, A. and Ozogul, Y. (2004) The effects of modified atmosphere packaging and vacuum packaging on chemical, sensory and microbiological changes of sardines (*Sardina pilchardus*). *Food Chem.* **85**, pp 49-57.
- Ozogul, F., Taylor, K.D.A., Quantick, P. and Ozogul, Y. (2000 a) A rapid HPLC determination of ATP related compounds and its application to herring stored under modified atmosphere. *Int. J. Food Sci. and Technol.* **35**, pp 549-554.
- Ozogul, F., Taylor, K.D.A., Quantick, P. and Ozogul, Y. (2000 b) Chemical, Microbiological and Sensory Evaluation of Atlantic herring (*Clupea harengus*) stored in ice, modified atmosphere and vacuum pack. *Food Chem.* **71**, pp 267-273.
- Palumbo, S. (1986) Is refrigeration enough to restore foodborne pathogens? *J. Food Prot.* **49**, pp 1003-1011.
- Papadopoulos, V., Chouliara, I., Badeka, A., Savvaidis, I.N. and Kontominas, M.G. (2002) Effect of gutting on microbiological, chemical, and sensory properties of aquacultured sea bass (*Dicentrarchus labrax*) stored in ice. *Food Microbiol.* **20**, pp 411-420.
- Parkin, K.L. and Brown, W.D. (1982) Preservation of seafoods with modified atmospheres. In *"Chemistry and biochemistry of marine food products"* AVI Publishing Co., Westport, CT.
- Parkin, K.L., Wells, M.J. and Brown, W.D. (1981) Modified atmosphere storage of rockfish fillets. *J. Food Sci.* **47**, pp 181-195.
- Parry, R.T. (1993) Introduction. In: Parry, R.T. (ed.). *Principles and applications of modified atmosphere packaging of food*. London: Blackie. pp. 1-18.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, L.M. (1998) Influence of sodium chloride and modified atmosphere packaging on microbiological, chemical and sensorial properties in ice storage of slices of hake (*Merluccius merluccius*). *Food Chem.* **61**, pp 23-28.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, L.M. (1996) Effect of modified atmosphere on shelf life of iced fresh hake slices. *J. Sci. Food Agr.* **71**, pp 541-547.
- Pedersen, L. and Snabe, L. (1995) Isolation of bacteriocin-producing Lactic Acid Bacteria

from Chilled, Vacuum- packaged Temperate and Tropical Fish Products. M.Sc. Thesis. Danish Institute for Fisheries Research, Lyngby, and The Royal Veterinary and Agricultural University of Copenhagen, Denmark.

- Penney, N., Bell, R.G. & Cummings, T.I. (1994). Extension of the chilled storage life of smoked blue cod (*Parapercis colias*) by carbon dioxide packaging. *Int. J. Food. Sci. Technol*, **31**, pp 463-479.
- Post, L.S., Lee, D.A., Solberg, M., Furgang, D., Specchio, J. and Graham, C. (1985) Development of botulinal toxin and sensory deterioration during storage of vacuum and modified atmosphere packaged fish fillets. *J. Food Sci.* **50**, pp 990-996.
- Pothuri, P., Marshall, D.L. & McMillin, K.W. (1996). Combined effects of packaging atmosphere and lactic acid on growth and survival of *Listeria monocytogenes*. Crayfish tail meat at 4°C. *J. Food protect.*, **59**, pp 253-256.
- Przybyrski, L.A., Finerty, M.W., Grodner, R.M. and Gerdes, D.L. (1989) Extension of shelf-life of iced fresh channel catfish fillets using modified atmosphere packaging and low dose irradiation. *J. Food Sci.* **54**, pp 269-278.
- Randell, K., Abvenainen, R. and Hattula, T. (1995) Effect of gas/product ratio and CO₂ concentration on the shelf life of MA packed fish. *Packaging Technol. and Sci.* **8**, pp 205-218.
- Randell, K., Hattula, T., Skytta, E., Sivertsvik, M., Bergslien, H. and Ahvenainen, R. (1999) Quality of filleted salmon in various retail packages. *J. Food Quality.* **22**, pp 485-497.
- Rccourt, J & Cossart, P (1997). *Listeria monocytogenes*. In :Food Microbiology- Fundamentals and Frontiers (edited by M.E. Doyle, L.R. Beuchat and T.J. Montville). pp 337-352. Washington, DC: ASM Press.
- Reddy, N.R., Paradis, A., Roman, M.G., Solomon, H.M. and Rhodehamel, E.J. (1996) Toxin development by *Clostridium botulinum* in modified atmosphere packaged fresh tilapia fillets during storage. *J. Food Sci.* **61**, pp 632-635.
- Reddy, N.R., Roman, M.G., Solomon, H.M., Kauter, D.A. and Rhodehamel, E.J. (1997a) Shelf life and *Clostridium botulinum* toxin development during storage of modified atmosphere packaged fresh catfish fillets. *J. Food Sci.* **62**, pp 878-884.
- Reddy, N.R., Solomon, H.M., Yep, H., Roman, M.G. and Rhodehamel, E.J. (1997b) Shelf life and toxin development by *Clostridium botulinum* during storage of modified atmosphere packaged fresh aqua cultured salmon fillets. *J. Food Prot.* **60**, pp 1055-1063.
- Reddy, N.R., Villanueva, M. and Kautter, D.A. (1995) Shelf life of modified atmosphere packaged tilapia fillets stored under refrigeration and temperature abuse conditions. *J. Food Prot.* **58**, pp 908-914.
- Reddy, N.R., Armstrong, D.J., Rhodehamel, E.J. and Kautter, D.A. (1992) Shelf-life extension and safety concerns about fresh fishery products packaged under modified atmospheres: a review. *J. Food Saf.* **12**, pp 87-118.
- Reddy, N.R., Solomon, H.M. and Rhodehamel, E.J. (1999) Comparison of margin of safety between sensory spoilage and onset of *Clostridium botulinum* toxin development during storage of modified atmosphere-packed fresh marine cod fillets with MA packaged aqua cultured fish fillets. *J. Food Saf.* **19**, pp 171-183.
- Ryder, J.M. (1985) Determination of adenine triphosphate and its breakdown products in fish muscle by high performance liquid chromatography. *J. Agric. Food Chem.* **33**, pp 678-680.
- Sacks, B. and Gore, A. (1987) Gas packaging-techniques and trends, *Food Rev.*, **14**, pp 3-9.
- Saito, T., Arai, K. and Matsuyoshi, M. (1959) A new method for estimating the freshness of fish. *Bull. Japan. Soc. Sci. Fish.* **24**, pp 749-750.
- Savvaiddis, I.N., Skandamis, P.N., Riganakos, K.A., Panagiotakis, N. and Kontominas, M.G. (2002) Control of natural microbial flora and *Listeria monocytogenes* in Vacuum packaged trout at 4°C and 10°C using irradiation. *J. Food Prot.* **65**, pp 515-522.
- Schweisfurth and Kalle Aktiengesellschaft (1970) Process for the manufacture of a portion package of fresh meat. Patent Specification 1, 186, 978, London.
- Scott, D.N., Fletcher, G.C. & Hogg, M.G. (1986). Storage of snapper fillets in modified

- atmosphere s at -1°C . *Food Technol. Australia*, **38**, pp 234-238.
- Scott, D.N., Fletcher, G.C. & Summers, G. (1984). Modified atmosphere and vacuum packaging of snapper fillets. *Food Technol. Australia*, **36**, pp 330-334
- Shewan, J.M. (1950) Improving the quality of white fish by the use of gas storage. *Fishing News*. 1946, pp 14-21
- Shewan, J.M. (1977) The bacteriology of fresh spoiling fish and biochemical changes induced by bacterial action. In *Proceedings of the Conference on the Handling, Processing and Marketing of Tropical Fish*. Tropical products Institute, London, pp 511-566.
- Shewan, J.M. (1949) Some bacteriological aspects of handling, processing, and distribution of fish. *J. Roy. Sanit. Inst.* **69**, pp394-402.
- Silliker, J.H. and Wolfe, S.K. (1980) Microbiological safety considerations in controlled atmosphere storage of meats. *Food Technol.* **34**, pp. 59-63.
- Silva, J.L. and White, T.D. (1994) Bacteriological and color changes in modified atmosphere packaged refrigerated channel catfish. *J. Food Prot.* **57**, pp 715-719.
- Smith, J.P., Hosahalli, S., Ramaswamy, H.S. and Simpson, B.K. (1990) Developments in food packaging technology. Part II: Storage aspects. *Trends Food Sci. Technol.* **1**, pp 111-118.
- Sorensen, N.K., Solberg, T. and Hansen, G.T. (1990) Storage of wet, iced, salmon under modified atmosphere, in *Chilling and Freezing of New Fish Products*, Proc. Int. Refrig., Comm. C2, Aberdeen 1990-3, pp. 167-171.
- Stammen, K., Gerdes, D. and Caporaso, F. (1990) Modified atmosphere packaging of seafood. *Critical reviews in Food Science and Technology*, **29**, pp 301-331.
- Stansby, M.E. and Griffiths, F.P. (1935) Carbon dioxide in handling fresh fish. *Ind. Eng. Chem.* **27**, pp 1452-1468
- Statham, J.A., Bremner, H.A. and Quarmby, A.R. (1985) Storage of morwong (*Nemadactylus macropterus* Bloch and Schneider) in combinations of polyphosphate, potassium sorbate and carbon dioxide at 4°C . *J. Food Sci.* **50**, pp 1580-1587.
- Statham, J.A. and Bremner, H.A. (1989) Shelf life extension of packaged seafoods- a summary of a research approach. *Food Aust.* **41**, pp 614-620.
- Stenstrom, I.M. (1985) Microbial flora of cod fillets (*Gadus morhua*) stored at 2°C in different mixtures of carbon dioxide and nitrogen/oxygen. *J. Food Prot.* **48**, pp 585-589.
- Stenstrom, I.M., Molin, G. (1990) Classification of the spoilage flora of fish, with special reference to *Shewanella putrefaciens*. *J. Appl. Bacteriol.* **68**, pp 601-618.
- Strasidine, G.A., Mah, G., and Monteith, D., (1982) Modified atmosphere storage-microbiology of cod fillets, Tech. Rep. No. 7, Division of Fisheries Technology, B.C. Research, Vancouver, B.C.
- Tarr, H.L.A. (1954) Microbiological deterioration of fish postmortem: its detection and control. *Bacteriol. Rev.* **18**, pp 1-21.
- Taylor, L.Y., Cann, D.D. and Welch, B.J. (1990) Antibotulinal properties of nisin in fresh fish packaged in an atmosphere of carbon dioxide. *J. Food Prot.*, **53**, pp 953-957.
- Tiffiney, P. and Mills, A. (1982) Storage trials of controlled atmosphere packaged fish products. Sea Fish Industry Authority. Technical Report No. **191**.
- Tomkins, R.G. (1932) The inhibition of the growth of meat attaching fungi by carbon dioxide. *J. Soc. Chem. Ind.* **51**, pp 261-275.
- Uchiyama, H., Ehira, S., Kawahara, T. and Shimidu, W. (1970) Significance in measuring volatile base and trimethyl amine nitrogen and nucleotides in fish muscle as indices of freshness. *Bull Jap Soc Sci Fish* **36**, pp 177-181.
- Van Spreekens, K.J.A. (1974) The suitability of a modification of Long and Hammerss medium for the enumeration of more fastidious bacteria from fresh fishery products. *Arch. Lebensm. Hyg.* **25**, pp 213-219.
- Varga, S., Keith, R.A., Michalik, P., Sims, G.G. and Reiger, L.W. (1980) Stability of lean and fatty fish fillets in hypobaric storage. *J. Food Sci.* **45**, pp 1487-1491.

- Villemure, G., Simard, R.F. and Picard, G. (1986) Bulk storage of cod fillets and gutted cod (*Gadus morhua*) under carbon dioxide atmosphere. *J. Food Sci.* **51**, pp 317-320.
- Wang, M.Y. and Orgydzia, D.M. (1986) Residual effect of storage in an elevated carbon dioxide atmosphere on the microbial flora of rock cod (*Sebastes spp.*). *Appl. Environ. Microbiol.* **52**, pp 727-729.
- Wang, M.Y. and Brown, W.O. (1983) Effect of elevated CO₂ atmosphere on storage of freshwater crayfish (*Pacifastacus leniusculus*). *J. Food Sci.* **48**, pp 158-159.
- Woyewoda, A.D., Bligh, E.G. and Shaw, S.J. (1984) Controlled and modified atmosphere storage of cod fillets. Canadian Institute Food Science and Technology, **17**, pp 24-27.