



Dietary Quercetin Mitigates Aflatoxin B₁-Induced Toxicity in *Labeo rohita*: Evidence from Physiological and Tissue Micro-Architecture Responses

Deepa Bhatt^{1*}, Abhed Pandey^{1,4}, Shanthanagouda, A. Holeyappa^{2,5}, Neelam Bansal³, Sachin Onkar Khairnar^{1,4} and Jaspal Singh Hundal⁶

¹Department of Aquaculture, College of Fisheries, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana - 141 004, Punjab, India

²Department of Aquatic Environment, College of Fisheries, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana - 141 004, Punjab, India

³Department of Veterinary Anatomy, College of Veterinary Sciences, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana - 141 004, Punjab, India

⁴Department of Aquaculture, College of Fisheries, Bihar Animal Sciences University, Patna, Arrabari, District- Kishanganj - 855 107, Bihar (India)

⁵Inland Fisheries Unit, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Navule Shivamogga - 577 204, Karnataka (India)

⁶Department of Animal Nutrition, College of Veterinary Sciences, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana - 141 004, Punjab, India

Abstract

This study evaluated the protective effect of dietary quercetin (QUE) against aflatoxin B₁ (AFB₁) toxicity in *Labeo rohita* fingerlings during a 90-day feeding trial. Fish were assigned to different dietary treatments, including a control (CON), AFB₁-contaminated diets, and AFB₁ diets supplemented with QUE (200 mg kg⁻¹). Growth performance, biochemical parameters, antioxidant enzyme activities, and tissue micro-architecture of the liver, and kidney were assessed. Results revealed that QUE supplementation significantly improved growth, enhanced biochemical and antioxidant responses, and reduced tissue damage, particularly at lower AFB₁ exposure. However, higher AFB₁ levels caused noticeable physiological disturbances and histopathological alterations. QUE was effective in alleviating oxidative stress and preserving tissue integrity in exposed fish. Overall, the study demon-

strates that dietary QUE can mitigate AFB₁-induced toxicity and improve fish health, suggesting its potential as a natural feed additive for enhancing resilience against mycotoxin contamination in aquaculture systems.

Keywords: Flavonoid supplementation, mycotoxin detoxification, oxidative stress modulation, physiology

Introduction

Mycotoxin contamination in aquaculture feeds has become a major concern affecting fish health, feed safety, and overall productivity. These toxic secondary metabolites, produced by various fungi, commonly contaminate agricultural commodities and feed ingredients, with nearly 25% of global crops affected annually (Eskola et al., 2020). Among them, aflatoxins, ochratoxin A, fumonisins, deoxynivalenol, and zearalenone are the most significant in animal production systems (Pitt, Taniwaki, & Cole, 2013). Common plant-based feed ingredients such as corn, soybean meal, wheat, DDGS, canola, and cottonseed are particularly prone to fungal contamination. The growing reliance on these ingredients in aquafeeds has increased the risk of mycotoxin exposure in fish,

Received 25 March 2026; Revised 13 June 2026; Accepted 29 June 2026

Handling editor: Dr. Tincy Varghese

*Email: deepabhatt94@gmail.com

highlighting the need for effective mitigation strategies to ensure sustainable aquaculture practices. Among mycotoxins, AFB₁ is one of the most toxic and carcinogenic, primarily produced by *Aspergillus flavus* and *A. parasiticus* (Bhatt & Pandey, 2022). Its contamination can occur both during crop growth and storage, especially under warm and humid conditions (Peng et al., 2022). The presence of AFB₁ in aquafeeds poses a serious risk to fish health and food safety, as it negatively affects growth, immune function, and metabolic activities (Gonçalves et al., 2018). Numerous studies across various fish species have demonstrated its effects in species such as tilapia (*Oreochromis aureus* × *O. niloticus*) (Deng et al., 2010; Hussain, Mateen, & Gatlin, 2017), rohu (*L. rohita*) (Bhatt, Holeyappa et al., 2024; Bhatt, Pandey, Holeyappa, Bansal, & Khairnar, 2024; Bhatt, Pandey, Holeyappa, Hundal, & Khairnar, 2024), and common carp (*Cyprinus carpio*) (Tasa, Imani, Moghanlou, Nazdar, & Moradi-Ozarlou, 2020; Nasirin et al., 2023). AFB₁ exposure has been shown to impair growth performance, disrupt hepatic function, and alter haematological and physiological parameters, highlighting its significant impact on aquaculture systems.

In recent years, natural bioactive compounds, particularly polyphenols, have gained considerable attention as dietary supplements to improve fish health and mitigate toxic stress in aquaculture. These compounds are known to enhance growth performance, antioxidant capacity, immune responses, and disease resistance in fish (Ahmadifar et al., 2021; Nasirin et al., 2023; Terzi, Tahliluddin, & Kadak, 2023). Among them, QUE, a flavonoid widely present in fruits and vegetables such as onions, apples, and cabbage, has attracted significant interest due to its strong antioxidant, anti-inflammatory, and protective properties (Dong et al., 2020). QUE acts as an efficient scavenger of reactive oxygen and nitrogen species, thereby protecting biological systems from oxidative damage (Boots, Haenen, & Bast, 2008), and has been shown to regulate antioxidant defence mechanisms (Pan et al., 2023). Studies in fish such as zebrafish, tilapia, and silver catfish (*Rhamdia quelen*) have demonstrated that QUE supplementation alleviates oxidative stress, inflammation, and lipid peroxidation while enhancing antioxidant status (Pês et al., 2018; Deng et al., 2019; Zhang, Jiang, Wang, & Qi, 2021). Additionally, it may improve gut health, nutrient absorption, and growth performance (Shabbir et al., 2021; Xin et al., 2024). However, complete preven-

tion of AFB₁ contamination in aquafeeds remains challenging, and its adverse effects can only be partially controlled, highlighting the need for effective dietary interventions.

Therefore, developing effective nutritional strategies to mitigate AFB₁-induced toxicity in aquaculture is essential, as information on the protective role of QUE in Indian major carps remains limited. *L. rohita*, a key species in South Asian aquaculture, is highly susceptible to mycotoxin-contaminated feeds. This study evaluated the effects of dietary AFB₁ exposure and the protective role of QUE on growth, biochemical responses, antioxidant status, and tissue integrity to support improved fish health and sustainable aquaculture.

Materials and Methods

AFB₁ and QUE (≥ 95% purity), were purchased from Sigma-Aldrich (St. Louis, MO, USA). Five experimental diets (CON, AFB₁100, AFB₁25+QUE, AFB₁50+QUE, and AFB₁100+QUE) were formulated to be isonitrogenous and isocaloric along with a negative CON (basal feed+AFB₁0), and a positive control (AFB₁100). The dietary AFB₁ concentrations (25, 50, and 100 ppb) were selected based on previous studies reporting toxicological effects of AFB₁ in fish at comparable exposure levels (Ahmed et al., 2020; Hoseyni et al., 2024). These concentrations represent low to moderate contamination levels that may occur in aquafeeds under practical farming conditions. The quercetin supplementation level (200 mg kg⁻¹ diet) was selected based on previous studies demonstrating its beneficial effects on growth performance, antioxidant status, and immune responses in fish (Xu et al., 2019; Armobin et al., 2023). All basal feed ingredients were finely ground using a grinder and passed through a **60-mesh sieve** to obtain uniform particle size. QUE was carefully weighed and thoroughly mixed with the other feed ingredients to ensure uniform distribution before pellet formation. Pellets were formulated with varying ingredient levels according to the formulations outlined in Table 1. Precisely weighed ingredients were thoroughly blended in a grinder and pressure-cooked in an autoclave at 115 °C pounds per square inch (psi) for 15 minutes. Following cooking, the dough was cooled to room temperature, and AFB₁ dilutions was dissolved in 70% methanol to create a stock solution, which was then serially diluted (25, 50, and 100 ppb). These solutions were then uniformly sprayed onto the

Table 1. Details of treatments with respect to feed formulation and their inclusion level

Details of diet groups					
Diet groups	Dosage				
AFB ₁ 0 (CON)	Basal diet* without AFB ₁ kg ⁻¹ feed (CON)				
AFB ₁ 100 (+ve control)	Basal diet with 100ppb AFB ₁ kg ⁻¹ feed				
AFB ₁ 25+QUE	Basal diet with 25ppb AFB ₁ and 200 mg QUE kg ⁻¹ feed				
AFB ₁ 50+QUE	Basal diet with 50ppb AFB ₁ + 200 mg QUE kg ⁻¹ feed				
AFB ₁ 100+QUE	Basal diet with 100ppb AFB ₁ + 200 mg QUE kg ⁻¹ feed				
Ingredient formulations and compositions of experimental diets					
Ingredients	CON	AFB ₁ 100	AFB ₁ 25+QUE	AFB ₁ 50+QUE	AFB ₁ 100+QUE
Soyabean (%)	40	40	40	40	40
Rice bran (%)	30	30	30	30	30
Mustard meal (%)	26	26	26	26	26
Vitamin-mineral mixture (%)	2.0	2.0	2.0	2.0	2.0
Carboxy methyl cellulose (%)	2.0	2.0	2.0	2.0	2.0
QUE (mg)	-	-	200	200	200
AFB ₁ (ppb)	-	100	25	50	100

QUE: Quercetin; AFB₁: Aflatoxin B₁

cooked feed ingredients using a hand sprayer. Subsequently, the ingredients were left to undergo methanol evaporation in the shade. Once dried, all ingredients were meticulously mixed with the prescribed amounts of turmeric, vitamin-mineral premix, and binder, ensuring homogeneous blending, and hand-kneaded to achieve the desired consistency, adding sufficient water to form dough. The dough was then extruded through a mechanical pelletizer. The resultant pellets were dried in a hot air oven at 40 °C overnight until the moisture content reached 10%. Each diet was individually packed in high-density polythene bags, labelled, and stored in a refrigerator at 4 °C until further use, consistent with the methodology previously described (Bhatt, Holeyappa et al., 2024). Proximate composition analysis of diets (moisture, crude protein, crude fat, crude fiber, and ash) was performed using standard AOAC (2012) methods, and results are presented in Table 2.

Healthy fish fingerlings were sourced from Fish Farm, College of Fisheries, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, Punjab (India) and then acclimated for 20 days in fibre-reinforced plastic (FRP) experimental tanks under a controlled temperature range of 24-28 °C.

During this period, they were fed a standard basal diet and a constant temperature (24-28 °C) was maintained before the experiment. A total of 180 fish (mean initial length: 12.40 ± 0.06 cm; mean initial weight: 17.31 ± 0.03 g) were allocated into 15 FRP (1.5x1.0x0.75 m) tanks (12 fish/tank), with triplicate groups assigned to each dietary treatment and maintaining a stocking density of 12 fish per tank with continuous aeration. The fish were fed twice daily at 2-3% of their body weight (9-10 AM and 4-5 PM) for 90 days. Before each feeding, leftover feed and fecal matter were siphoned out to maintain water quality. Water in the tanks was partially exchanged (33%) with pre-aerated water every week to remove accumulated waste. The experimental diets and pellets were prepared following the methods described in previous studies (Bhatt, Holeyappa et al., 2024; Bhatt, Pandey, Holeyappa, Bansal, & Khairnar, 2024). The water quality parameters were maintained as follows: dissolved oxygen >6.0 mg L⁻¹, total ammonia nitrogen <0.2 mg L⁻¹, temperature 28-30 °C, and pH 7.4-7.8.

At the end of the 90-day feeding trial, the experimental fish were starved for 24 h to clear the gut contents prior to sampling. Subsequently, the fish were anesthetized using 2-phenoxyethanol at a

Table 2. Proximate composition of different feed ingredients and experimental diet (% of dry weight)

Ingredients/ diets	^[1] CP (%)	EE (%)	CF (%)	Ash (%)	NFE (%)
Rice bran	12.38	1.28	15.64	11.51	59.19
Mustard meal	38.91	1.93	11.54	8.16	39.46
Soybean meal	42.9	1.13	4.01	7.2	44.76
QUE	9.53	0.97	0.07	2.3	87.13
(Experimental diets)					
CON	25.47	3.32	19.82	9.40	41.99
AFB ₁ 100	23.38	3.12	19.12	9.12	45.26
AFB ₁ 25+QUE	24.26	3.28	19.91	9.24	43.31
AFB ₁ 50+QUE	24.41	3.25	19.89	9.21	43.24
AFB ₁ 100+QUE	23.38	3.13	19.13	9.16	45.2

^[1] CP: Crude protein; EE: Ether extract; CF: Crude fiber; NFE: Nitrogen-free extract

concentration of 0.4 mL L⁻¹. Each fish was then individually weighed and measured to evaluate growth performance parameters before being returned to their respective tanks. For hematological analysis, blood samples were collected from three randomly selected fish per tank through the caudal vein using 26G × 1/2" (0.45 × 13 mm) disposable syringes (Hindustan Syringes & Medical Devices Ltd.). The collected blood was transferred into EDTA-coated tubes using heparinized syringes to prevent clotting and was used for the determination of hematological parameters. For serum biochemical analysis, blood samples were collected from three fish per tank without anticoagulant. The samples were kept at 4 °C for 24 h to allow clot formation and were subsequently centrifuged at 3000 rpm for 10 min at 4 °C. The obtained serum (supernatant) was carefully separated, aliquoted, and stored at -80 °C until further biochemical analysis. For histological examination, approximately 1 cm tissue segments of liver and kidney, were collected from four randomly selected fish. The tissues were immediately fixed in 10% neutral buffered formalin (NBF) and stored at room temperature until further histological processing and analysis.

Growth performance indices were calculated using standard formulas (Halver, 1957):

Specific growth rate (SGR; % weight gain day⁻¹) = $\log(\ln) \text{ final body weight (g)} - \log(\ln) \text{ initial body weight (g)} / t \text{ (time interval in days)} \times 100$

Feed conversion ratio (FCR) = feed intake (g)/ weight gain (g)

Weight gain (%) = $[\text{Final body weight (g)} - \text{Initial body weight}] / \text{Initial body weight (g)} \times 100$

For proximate composition analysis, fish samples were subjected to autoclave treatment (120 °C for 20 min), followed by homogenisation, oven drying, and grinding into fine powder prior to chemical analysis. The proximate composition, including moisture, crude protein, crude lipid, crude fibre, and ash content, was determined following standard procedures recommended by the Association of Official Analytical Chemists (AOAC, 2012).

Serum biochemical parameters, including alanine aminotransferase (ALT; Cat. #120207), aspartate aminotransferase (AST; Cat. #120204), total protein (TP; Cat. #120231), glucose (GLU; Cat. #120235), and albumin (Cat. #120223), were determined using commercially available ERBA® Mannheim diagnostic kits. The analyses were performed using a CHEM-7 biochemical analyser, following the manufacturer's instructions. For each biochemical parameter, 50 µL of serum was used for analysis according to the standard assay procedure provided with the diagnostic kits. Both standards and serum samples were analysed in triplicate to ensure analytical accuracy and reproducibility of the results. The activities of antioxidant enzymes, including superoxide dismutase (SOD) and catalase (CAT), were determined following established stan-

Table 3. Growth performances of *L. rohita* in different treatments at the termination of experiment

Parameters ^[1]	Treatments				
	CON	AFB ₁ 100	AFB ₁ 25+QUE	AFB ₁ 50+QUE	AFB ₁ 100+QUE
Initial body weight (g)	17.36±0.13	17.31±0.08	17.34±0.10	17.31±0.15	17.32±0.19
Final body weight (g)	30.59 ^a ±0.11	26.25 ^d ±0.06	30.31 ^a ±0.05	29.15 ^b ±0.14	28.38 ^c ±0.12
NBWG (g)	13.26 ^a ±0.73	8.94 ^c ±0.12	12.97 ^{ab} ±0.12	11.84 ^b ±0.05	11.06 ^{bc} ±0.04
SGR	0.89 ^a ±0.02	0.59 ^d ±0.01	0.81 ^b ±0.04	0.69 ^c ±0.03	0.64 ^c ±0.07
FCR	1.19 ^c ±0.01	1.26 ^a ±0.01	1.20 ^b ±0.02	1.21 ^b ±0.01	1.25 ^a ±0.05
K-value	1.23 ^a ±0.01	0.64 ^d ±0.01	1.22 ^a ±0.02	1.04 ^b ±0.01	0.92 ^c ±0.03
Survival rate (%)	100.00 ^a ±0.00	79.76 ^e ±2.77	98.21 ^b ±2.77	94.59 ^c ±0.45	91.54 ^d ±2.77

^[1] NBWG: net body weight gain. SGR: specific growth rate. FCR: feed conversion ratio. K: condition factor. Values are mean ± S.E. ($p \leq 0.05$). ^{a, b, ..., e}: values with same superscript in a row do not differ significantly ($p \leq 0.05$)

dard protocols (Nicholls, 1962; Nishikimi, Rao, & Yagi, 1972). Prior to analysis, tissue samples were subjected to appropriate homogenisation and preparation procedures. All steps related to sample preparation, tissue homogenisation, reagent preparation, and enzyme assay procedures were performed strictly according to the manufacturer's guidelines supplied with the assay kits to ensure accuracy, reliability, and consistency of the analytical results.

Fish from the CON and experimental groups were fixed in 10% NBF for 24 hours and washed under running tap water. The tissues were dehydrated in graded alcohol (70% to absolute), cleared in xylene, and embedded in paraffin wax using molds. Thin sections (4 µm) were prepared using a rotary microtome (Leica RM2125 RT, Germany) and mounted onto Super Frost slides (#G10106, Abdos, India). After drying overnight at ~33 °C, the sections were stained with hematoxylin and eosin, mounted with DPX (Product Code 23140, Molychem, Mumbai), and images were captured using a Leica DM3000 LED light microscope.

All experimental data were expressed as mean ± standard error (SE). Statistical analysis was performed using appropriate statistical software. Differences among treatment groups were evaluated using one-way analysis of variance (ANOVA), followed by suitable post hoc tests to determine significant differences between means. A significance level of $p < 0.05$ was considered statistically significant.

Results and Discussion

During the experiment, no mortality was observed in the CON group. Growth parameters, including net body weight gain (NBWG) and SGR, were highest in the control group; however, the differences were not statistically significant ($p > 0.05$) compared to the other treatment groups. (Table 3). However, growth indicators declined notably in the AFB₁100 group exposed to 50 ppb AFB₁. Interestingly, dietary supplementation with 25 ppb AFB₁ and 200 mg QUE per kg of diet (AFB₁25) significantly enhanced growth indices in *L. rohita* fingerlings, with NBWG and SGR increasing by 12.97% and 0.81%, respectively, compared to other AFB₁-exposed treatments (AFB₁50 and AFB₁100). Among the AFB₁-exposed groups, the AFB₁25 treatment showed the highest NBWG; however, the difference was not statistically significant ($p > 0.05$), highlighting the potential protective effects of QUE.

In addition to growth impairment, prolonged dietary exposure to AFB₁ adversely affected survival, with the AFB₁100 group exhibiting the lowest survival rate among the treatments. These findings indicate that higher dietary AFB₁ concentrations negatively influence both growth and survival in *L. rohita* fingerlings. The observed reduction in growth may be attributed to impaired protein metabolism and disruptions in glucose and lipid utilisation, as reported by Zaki and Fawzy (2012) and Zychowski et al. (2013). Conversely, dietary quercetin supplementation improved growth performance and survival, suggesting a protective role against AFB₁-induced toxicity. This beneficial effect may be

Table 4. Biochemical Parameters and Anti-oxidant Enzymes of *L. rohita* in different treatments at the termination of experiment

Biochemical Parameters ^[2]	Treatments				
	CON	AFB ₁ 100	AFB ₁ 25+QUE	AFB ₁ 50+QUE	AFB ₁ 100+QUE
Blood glucose (mg dL ⁻¹)	67.4 ^d ±0.05	98.15 ^a ±0.18	72.85 ^c ±0.19	74.46 ^c ±0.05	82.20 ^b ±0.12
Total protein (g dL ⁻¹)	3.75 ^a ±0.01	2.24 ^d ±0.03	3.71 ^a ±0.02	3.24 ^b ±0.04	2.97 ^c ±0.06
Albumin (g dL ⁻¹)	1.47 ^a ±0.04	0.54 ^d ±0.02	1.25 ^b ±0.05	1.04 ^b ±0.02	0.76 ^c ±0.05
Globulin (g dL ⁻¹)	2.27 ^b ±0.03	1.71 ^d ±0.01	2.46 ^a ±0.01	2.19 ^c ±0.02	2.20 ^b ±0.01
A/G Ratio	0.65 ^a ±0.04	0.31 ^d ±0.03	0.51 ^b ±0.04	0.47 ^c ±0.01	0.34 ^d ±0.002
ALT /SGPT (IU L ⁻¹)	21.46 ^d ±0.03	26.75 ^a ±0.01	23.16 ^c ±0.04	24.26 ^b ±0.06	26.14 ^a ±0.05
AST/SGOT (IU L ⁻¹)	97.28 ^e ±0.02	140.35 ^a ±0.01	105.49 ^d ±0.03	115.51 ^c ±0.02	131.29 ^b ±0.05
Anti-oxidant Enzymes					
CAT (U mg Hb ⁻¹)	2.14 ^d ±0.03	3.00 ^a ±0.03	2.32 ^c ±0.01	2.68 ^b ±0.02	2.92 ^b ±0.01
SOD (U mg Hb ⁻¹)	0.35 ^c ±0.02	0.60 ^a ±0.04	0.38 ^c ±0.05	0.56 ^b ±0.03	0.58 ^b ±0.04

^[2] ALT: Alanine aminotransferase, SGPT: Serum glutamic pyruvic transaminase, AST: Aspartate aminotransferase, SGOT: Serum glutamic-oxaloacetic transaminase, SOD: Superoxide dismutase. CAT: Catalase. Values with same superscripts in a row do not differ significantly ($p > 0.05$), using one-way ANOVA; Post hoc: Duncan's multiple comparison

associated with the antioxidant and anti-inflammatory properties of quercetin, as previously reported by Han and Huang (2006).

Biochemical measures showed a substantial difference between the AFB₁ and QUE treated groups. The highest blood GLU levels were seen in AFB₁100, followed by the lowest in AFB₁0. Similarly, AFB₁100 and AFB₁100+QUE had lower TP, albumin, and globulin levels than AFB₁0, AFB₁25, and AFB₁50. The enzymatic activity of AST, ALT, CAT, and SOD in fish subjected to varied amounts of AFB₁ feeding treatments correlated negatively with rising AFB₁ inclusion levels. The treatment that received (combined dose) AFB₁ and QUE exhibited lower antioxidant levels as observed in the AFB₁25 group. As AFB₁ concentrations increased, enzymatic activity also increased, with the AFB₁100 group exhibiting the highest levels, followed by the AFB₁100+QUE group. The differences among the treatment groups were statistically significant (Table 4).

Significant changes in biochemical and antioxidant parameters were observed in AFB₁ and QUE-treated groups, including reduced TP, globulin, and A/G ratio, along with increased GLU, ALT, AST, SOD, and CAT levels. These alterations indicate a stress response involving metabolite mobilisation (Koolhaas et al., 2011). Elevated GLU may result from

enhanced glycogenolysis and reduced insulin activity, as reported by El-Boshy, El-Ashram, and El-Ghany (2008) in *O. niloticus*, while similar responses have been linked to immune activation by Akrami, Gharaei, Mansour, and Galeshi (2015). Evaluation of liver enzymes is essential for assessing toxicological effects (Sudová, Piačková, Kroupová, Pijáček, & Svobodová, 2008), while dietary bioactive compounds can further enhance physiological performance (Hossain, Small, Kumar, & Hardy, 2024).

The dietary intake of AFB₁ resulted in a non-significant increase in AST and ALT activities ($p > 0.05$), indicating potential liver dysfunction or swelling, leading to enzyme leakage into the bloodstream (Fawole et al., 2020). These enzymes serve as biomarkers for cellular injury, including degeneration and necrosis (Akrami et al., 2015). However, co-administration of AFB₁ and QUE mitigated the elevated enzyme levels. Similarly, Akrami et al. (2015) reported reduced ALT and AST activity in juvenile beluga (*Huso huso*) fed 1% onion, attributed to onion's antioxidant properties, which protect cell membranes from lipid peroxidation and enzyme diffusion. SOD and CAT activity increased with higher AFB₁ levels but decreased with QUE, consistent with QUE's role in regulating endogenous antioxidants and suppressing apoptosis (Moloi, Ziqubu, Mazibuko-Mbeje, Mabaso, & Ndlovu, 2024).




Fig. 1. Photomicrograph of H & E stained kidney sections of *L. rohita* showing A: (CON: 0 ppb AFB₁) Distal convoluted tubule (DCT), Proximal convoluted tubule (PCT), B: (Positive control, 100 ppb AFB₁) Evacuation of glomeruli (EVG), Haematopoietic tissue (HT); C: (25 ppb AFB₁+200 mg QUE) Distal convoluted tubule (DCT), Proximal convoluted tubule (PCT), Glomerulus (G); D: (50 ppb AFB₁+200 mg QUE) Distal convoluted tubule (DCT), Lumen (L), Nucleus (N), Glomerulus (G), Brush border (BB); E: (100 ppb AFB₁+200 mg QUE) Glomerulus (G), Enlarged proximal convoluted tubule (EPCT), (D) Scale bar = 50µm, H & E staining

SOD catalyzes O^{2•} into O₂ and H₂O₂, whereas CAT and glutathione peroxidase (GPx) disintegrate H₂O₂ into O₂ and H₂O, effectively eliminating excessive free radicals and preventing oxidative stress (Engwa, EnNwekegw, & Nkeh-Chungag, 2022; Jerald, Ravindran, & Babu, 2024) (Fig. 3).

In this study, a comprehensive examination of photomicrographs depicting the tissue microarchitecture of kidney structures was conducted across different experimental groups. The CON group (AFB₁0) had regular characteristics, including normal glomerulus, proximal convoluted tubule, and distal convoluted tubule architecture. Conversely, significant deviations were observed in the AFB₁-exposed group (AFB₁100), including the evacuation of glomeruli and the presence of hematopoietic tissue. Group AFB₁25 displayed degenerative changes primarily in the proximal and distal tubules, while the glomerulus maintained its typical structure with no significant alterations noted. Notably, group AFB₁50 showcased nearly normal proximal tubules with centrally placed lumens, a typical glomerular morphology and a well-defined brush border of epithelial cells. Con-

versely, group AFB₁100 presented enlarged proximal tubules and glomerulus, accompanied by a slight joining of tubules (Fig. 1).

The kidneys of teleost fish are vital hematopoietic and osmoregulatory organs and are highly susceptible to pollutants (Shahid et al., 2022; Smorodinskaya et al., 2023). In this study, high AFB₁ exposure caused significant alterations in kidney microarchitecture, indicating physiological stress and metabolic disturbances at cellular and tissue levels (Liu et al., 2024). Similar studies have linked flavonoid-rich compounds to nephroprotective effects through their antioxidant and free radical scavenging properties (Adeneye & Benebo, 2008). QUE, due to its polyphenolic nature, may exert similar protective effects by reducing oxidative stress and restoring tissue integrity in AFB₁-exposed fish.

The CON group's liver cell microarchitecture during the experimental period showed typical hepatocyte morphology, characterized by uniform cytoplasm, centrally located nuclei, and a discernible central vein (Fig. 2A). In contrast, the liver tissue of the




Fig. 2. Photomicrograph of H & E stained liver sections of *L. rohita* showing A: (CON: 0 ppb AFB₁) Hepatocytes (H), Central vein (CV), B: (Positive control, 100 ppb AFB₁) Invasion of hepatocytes in central vein (IH), Deposition (D) Vacuolization (V), C: (25 ppb AFB₁+200 mg QUE) Kupffer cells (KC), Nucleus (N), Sinusoids (S), Hepatocytes shows Karyorrhexis (HKRX) D: (50 ppb AFB₁+200 mg QUE) Kupffer cells (KC), Hepatic sinusoids (HS), Hemolysis of blood vessels (HBV); E: (100 ppb AFB₁+200 mg QUE) Kupffer cells (KC), Blood congestion (BC) Scale bar= 50μm, H & E staining

AFB₁100 group exhibited several notable alterations, notably including hepatocyte congestion and infiltration into the central vein, vacuolisation, intravascular hemolysis, and dilation of hepatic sinusoids, forming a distinctive tree-like pattern (Fig. 2B). Conversely, observations of the AFB₁25 group showed an existence of Kupffer cells, sinusoidal structures, and karyorrhexis (aggregation of chromatin material within hepatocytes) (Fig. 2C). Furthermore, the liver tissue of the AFB₁50 group displayed features such as hepatic sinusoids, Kupffer cells within macrophages, and complete occlusion of the central vein lumen with blood (hemolysis) (Fig. 2D). Notably, the AFB₁100 group demonstrated the presence of Kupffer cells and congestion within the central vein (Fig. 2E).

Evaluation of liver microarchitecture is a reliable indicator of health impairment in fish (Mokhtar, 2021). In the present study, higher AFB₁ concentrations caused hepatic injury and structural alterations, including dilated sinusoids and congestion of major veins. This disruption in blood flow from the hepatic portal vein and artery to the central vein can lead to cellular degeneration and necrosis

(Hinton, 2021). These findings are consistent with Azzam and Gababl (1997) and Tessari, Oliveira, Cardoso, Ledoux, and Rottinghaus (2006) who reported similar effects at comparable AFB₁ levels. QUE, known for its strong antioxidant and free radical scavenging properties (Wu et al., 2021), showed a protective effect when co-administered with AFB₁ by enhancing antioxidant enzyme activity and reducing tissue damage (Venkataraman, Krishnamoorthy, Selvakumar, & Arunakaran, 2009). The study highlights the impact of AFB₁ on organ integrity and provides valuable insights into the mitigating role of QUE in *L. rohita*, contributing to future research on AFB₁ toxicity.

The present study was designed to evaluate the protective efficacy of dietary quercetin against AFB₁-induced toxicity in *L. rohita*, the experimental design focused primarily on assessing its effects under toxin-challenged conditions. Inclusion of a quercetin-only group and dose-matched AFB₁-only controls (25 and 50 ppb) would have provided additional insight into the intrinsic effects of quercetin and enabled a more precise assessment of its protective efficacy across different toxin exposure




Fig. 3. Schematic representation of enzymatic reactions generating free radicals and antioxidant system during oxidative stress

levels. Nevertheless, the consistent improvements observed in growth performance, physiological and biochemical responses, antioxidant status, and tissue microarchitecture underscore the potential of quercetin as a dietary strategy for mitigating AFB₁-induced stress in fish. Future studies incorporating these additional control groups would further strengthen mechanistic understanding and support its application in aquaculture.

Overall, the present study demonstrated that dietary quercetin supplementation (200 mg kg⁻¹ diet) effectively mitigated the adverse effects of aflatoxin B₁ (AFB₁) in *L. rohita*. Quercetin improved growth performance, physiological and biochemical responses, antioxidant status, and tissue microarchitecture in AFB₁-exposed fish, indicating its protective role against mycotoxin-induced stress. The beneficial effects were observed across all AFB₁ exposure levels evaluated, supporting its potential as a natural feed additive in aquaculture. Although the extent of recovery varied among treatments, this variation is likely associated with differences in toxin burden rather than quercetin efficacy. The study was conducted under controlled conditions; therefore, further validation under field conditions

is required. Future studies incorporating quercetin-only groups, dose-matched AFB₁ controls, and molecular approaches such as transcriptomics and proteomics are recommended to elucidate the mechanisms of protection.

Acknowledgements

The authors express their sincere gratitude to the Dean, College of Fisheries, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, Punjab (India) for providing the resources and facilities necessary to carry out this research. The authors also thank the editors and anonymous reviewers for their valuable recommendations and constructive comments, which considerably improved the quality of the manuscript.

References

- Adeneye, A. A., & Benebo, A. S. (2008). Protective effect of the aqueous leaf and seed extract of *Phyllanthus amarus* on gentamicin and acetaminophen-induced nephrotoxic rats. *Journal of Ethnopharmacology*, 118(2), 318–323. <https://doi.org/10.1016/j.jep.2008.04.025>
- Ahmadifar, E., Yousefi, M., Karimi, M., Raieni, R. F., Dadar, M., Yilmaz, S., Dawood, M. A. O., & Abdel-Latif, H. M. R. (2021). Benefits of dietary polyphenols

- and polyphenol-rich additives to aquatic animal health: An overview. *Reviews in Fisheries Science & Aquaculture*, 29(4), 478–511. <https://doi.org/10.1080/23308249.2020.1818689>
- Ahmed, S., Baten, M. A., Hossain, M. M., Rahim, M. M., Rasul, M. G., Sultana, R., Hossain, M. M., & Bapary, M. A. J. (2020). Effects of aflatoxin-contaminated feed on the fingerlings of tilapia (*Oreochromis niloticus* Linnaeus, 1758). *Archives of Agriculture and Environmental Science*, 5(3), 390–396. <https://doi.org/10.26832/24566632.2020.0503022>
- Akrami, R., Gharaei, A., Mansour, M. R., & Galeshi, A. (2015). Effects of dietary onion (*Allium cepa*) powder on growth, innate immune response and hemato-biochemical parameters of beluga (*Huso huso* Linnaeus, 1754) juvenile. *Fish & Shellfish Immunology*, 45(2), 828–834. <https://doi.org/10.1016/j.fsi.2015.06.005>
- Armobin, K., Ahmadifar, E., Adineh, H., Samani, M. N., Kalhor, N., Yilmaz, S., Hoseinifar, S. H., & Van Doan, H. (2023). Quercetin application for common carp (*Cyprinus carpio*): I. Effects on growth performance, humoral immunity, antioxidant status, immune-related genes, and resistance against heat stress. *Aquaculture Nutrition*, 2023(1), Article 1168262. <https://doi.org/10.1155/2023/1168262>
- Association of Official Analytical Chemists. (2012). *Official methods of analysis* (19th ed.).
- Azzam, A. H., & Gabal, M. A. (1997). Interaction of aflatoxin in the feed and immunization against selected infectious diseases. I. Infectious bursal disease. *Avian Pathology*, 26(2), 317–325. <https://doi.org/10.1080/03079459708419214>
- Bhatt, D., Holeyappa, S. A., Pandey, A., Bansal, N., Hundal, J. S., & Khairnar, S. O. (2024). Growth performance, physiological response, and tissue microarchitecture of the carp *Labeo rohita* challenged with AFB1 are improved by supplementing with turmeric. *Spanish Journal of Agricultural Research*, 22(2), e0501–e0501. <https://doi.org/10.5424/sjar/2024222-19974>
- Bhatt, D., & Pandey, A. (2022). Aflatoxins and their Repercussions in Aquaculture: A review. *Fishery Technology*, 59(1), 69–78.
- Bhatt, D., Pandey, A., Holeyappa, S. A., Bansal, N., & Khairnar, S. O. (2024). Histomorphological alterations in the liver of *Labeo rohita* exposed to aflatoxin B₁ and quercetin-enriched diets. *National Academy Science Letters*, 47(4), 425–432. <https://doi.org/10.1007/s40009-023-01369-x>
- Bhatt, D., Pandey, A., Holeyappa, S. A., Hundal, J. S., & Khairnar, S. O. (2024). Protective effect of quercetin on haematology and flesh quality in *Labeo rohita* against aflatoxin B1. *National Academy Science Letters*, 47, 133–137. <https://doi.org/10.1007/s40009-023-01333-9>
- Boots, A. W., Haenen, G. R. M. M., & Bast, A. (2008). Health effects of quercetin: From antioxidant to nutraceutical. *European Journal of Pharmacology*, 585(2–3), 325–337. <https://doi.org/10.1016/j.ejphar.2008.03.008>
- Deng, S. X., Tian, L. X., Liu, F. J., Jin, S. J., Liang, G. Y., Yang, H. J., Du, Z. Y., & Liu, Y. J. (2010). Toxic effects and residue of aflatoxin B1 in tilapia (*Oreochromis niloticus* × *O. aureus*) during long-term dietary exposure. *Aquaculture*, 307(3–4), 233–240. <https://doi.org/10.1016/j.aquaculture.2010.07.029>
- Deng, Y., Qiu, M., Wang, Y., Wang, R., Lu, P., Sun, L., Li, X., & Gooneratne, R. (2019). Protective effect of antioxidant-enriched diets on T-2-toxin-induced damage in tilapia (*Oreochromis niloticus*). *Aquaculture*, 506, 341–349. <https://doi.org/10.1016/j.aquaculture.2019.03.066>
- Dong, Y., Hou, Q., Lei, J., Wolf, P. G., Ayansola, H., & Zhang, B. (2020). Quercetin alleviates intestinal oxidative damage induced by H₂O₂ via modulation of GSH: In vitro screening and in vivo evaluation in a colitis model of mice. *ACS Omega*, 5(14), 8334–8346. <https://doi.org/10.1021/acsomega.0c00804>
- El-Boshy, M. E., El-Ashram, A. M. M., & El-Ghany, N. A. A. (2008). Effect of dietary beta-1, 3 glucan on immunomodulation on diseased *Oreochromis niloticus* experimentally infected with aflatoxin B₁. In H. Elghobashy, K. Fitzsimmons, & A. S. Diab (Eds.), *From the Pharaohs to the Future: 8th international symposium on Tilapia in aquaculture* (pp. 1109–1127). World Aquaculture Society.
- Engwa, G. A., EnNwekegw, F. N., & Nkeh-Chungag, B. N. (2022). Free radicals, oxidative stress-related diseases and antioxidant supplementation. *Alternative Therapies in Health and Medicine*, 28(1), 114–128.
- Eskola, M., Kos, G., Elliott, C. T., Hajšlová, J., Mayar, S., & Krska, R. (2020). Worldwide contamination of food-crops with mycotoxins: Validity of the widely cited 'FAO estimate' of 25%. *Critical Reviews in Food Science and Nutrition*, 60(16), 2773–2789. <https://doi.org/10.1080/10408398.2019.1658570>
- Fawole, F. J., Adeoye, A. A., Tiamiyu, L. O., Ajala, K. I., Obadara, S. O., & Ganiyu, I. O. (2020). Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture*, 518, Article 734849. <https://doi.org/10.1016/j.aquaculture.2019.734849>
- Gonçalves, R. A., Cam, T. D., Tri, N. N., Santos, G. A., Encarnação, P., & Hung, L. T. (2018). Aflatoxin B₁ (AFB₁) reduces growth performance, physiological response, and disease resistance in Tra catfish (*Pangasius hypophthalmus*). *Aquaculture International*, 26(3), 921–936. <https://doi.org/10.1007/s10499-018-0259-x>

- Halver, J. E. (1957). Nutrition of salmonoid fishes: III. Water-soluble vitamin requirements of chinook salmon. *The Journal of Nutrition*, 62(2), 225–243. <https://doi.org/10.1093/jn/62.2.225>
- Han, A. Y., & Huang, R. L. (2006). Effects of dihydromyricetin on growth performance, immune organ indexes and serum biochemical parameters of broilers. *China Feed*, 10, 20–22.
- Hinton, D. E. (2021). Toxicologic histopathology of fishes: A systemic approach and overview. In J. A. Couch & J. W. Fournie (Eds.), *Pathobiology of marine and estuarine organisms* (pp. 177–215). CRC Press.
- Hoseyni, S. Z., Imani, A., Vazirzadeh, A., Moghanlou, K. S., Farhadi, A., & Razi, M. (2024). Dietary aflatoxin B₁ and zearalenone contamination affected growth performance, intestinal and hepatopancreas gene expression profiles and histology of the intestine and gill in goldfish, *Carassius auratus*. *Aquaculture Reports*, 34, Article 101887. <https://doi.org/10.1016/j.aqrep.2023.101887>
- Hossain, M. S., Small, B. C., Kumar, V., & Hardy, R. (2024). Utilization of functional feed additives to produce cost effective, ecofriendly aquafeeds high in plant based ingredients. *Reviews in Aquaculture*, 16(1), 121–153. <https://doi.org/10.1111/raq.12824>
- Hussain, D., Mateen, A., & Gatlin, D. M. (2017). Alleviation of aflatoxin B₁ (AFB₁) toxicity by calcium bentonite clay: Effects on growth performance, condition indices and bioaccumulation of AFB₁ residues in Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 475, 8–15. <https://doi.org/10.1016/j.aquaculture.2017.04.003>
- Jerald, I., Ravindran, J., & Babu, M. M. (2024). Fish in focus: Navigating organ damage assessment through analytical avenues—A comprehensive review. *Toxicology Reports*, 13, Article 101841. <https://doi.org/10.1016/j.toxrep.2024.101841>
- Koolhaas, J. M., Bartolomucci, A., Buwalda, B., de Boer, S. F., Flügge, G., Korte, S. M., Meerlo, P., Murison, R., Olivier, B., Palanza, P., Richter-Levin, G., Sgoifo, A., Steimer, T., Stiedl, O., van Dijk, G., Wöhr, M., & Fuchs, E. (2011). Stress revisited: a critical evaluation of the stress concept. *Neuroscience & Biobehavioral Reviews*, 35(5), 1291–1301. <https://doi.org/10.1016/j.neubiorev.2011.02.003>
- Liu, H., Xie, R., Huang, W., Yang, Y., Zhou, M., Lu, B., Li, B., Tan, B., & Dong, X. (2024). Effects of dietary aflatoxin B₁ on hybrid grouper (*Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*) growth, intestinal health, and muscle quality. *Aquaculture Nutrition*, 2024(1), Article 3920254. <https://doi.org/10.1155/2024/3920254>
- Mokhtar, D. M. (2021). *Fish histology: From cells to organs*. CRC Press.
- Moloi, T. P., Ziqubu, K., Mazibuko-Mbeje, S. E., Mabaso, N. H., & Ndlovu, Z. (2024). Aflatoxin B₁-induced hepatotoxicity through mitochondrial dysfunction, oxidative stress, and inflammation as central pathological mechanisms: A review of experimental evidence. *Toxicology*, 509, Article 153983. <https://doi.org/10.1016/j.tox.2024.153983>
- Nasirin, C., Najm, M. A., Chen, T. C., Dhamija, A., Lionardo, A., Bokov, D. O., & Naserabad, S. S. (2023). The protective effects of quercetin on the physiological responses in malathion-exposed common carp, *Cyprinus carpio*. *Tropical Animal Health and Production*, 55(1), Article 22. <https://doi.org/10.1007/s11250-022-03429-8>
- Nicholls, P. (1962). The reaction between aminotriazole and catalase. *Biochimica et Biophysica Acta*, 59(2), 414–420. [https://doi.org/10.1016/0006-3002\(62\)90191-9](https://doi.org/10.1016/0006-3002(62)90191-9)
- Nishikimi, M., Rao, N. A., & Yagi, K. (1972). The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochemical and Biophysical Research Communications*, 46(2), 849–854. [https://doi.org/10.1016/S0006-291X\(72\)80218-3](https://doi.org/10.1016/S0006-291X(72)80218-3)
- Pan, E., Chen, H., Wu, X., He, N., Gan, J., Feng, H., Sun, Y., & Dong, J. (2023). Protective effect of quercetin on avermectin induced splenic toxicity in carp: Resistance to inflammatory response and oxidative damage. *Pesticide Biochemistry and Physiology*, 193, Article 105445. <https://doi.org/10.1016/j.pestbp.2023.105445>
- Peng, K., Chen, B., Sun, Y., Chen, X., Wang, Y., & Huang, W. (2022). Response of growth performance, serum metabolites, intestinal tight junction structure and bacterial microbiomes to the long-term intervention of aflatoxin B₁ in *Lateolabrax maculatus* diets. *Aquaculture Reports*, 22, Article 101005. <https://doi.org/10.1016/j.aqrep.2022.101005>
- Pês, T. S., Saccol, E. M. H., Londero, É. P., Bressan, C. A., Ourique, G. M., Rizzetti, T. M., Prestes, O. D., Zanella, R., Baldissierotto, B., & Pavanato, M. A. (2018). Protective effect of quercetin against oxidative stress induced by oxytetracycline in muscle of silver catfish. *Aquaculture*, 484, 120–125. <https://doi.org/10.1016/j.aquaculture.2017.10.043>
- Pitt, J. I., Taniwaki, M. H., & Cole, M. B. (2013). Mycotoxin production in major crops as influenced by growing, harvesting, storage and processing, with emphasis on the achievement of food safety objectives. *Food Control*, 32(1), 205–215. <https://doi.org/10.1016/j.foodcont.2012.11.023>
- Shabbir, U., Rubab, M., Daliri, E. B. M., Chelliah, R., Javed, A., & Oh, D. H. (2021). Curcumin, quercetin, catechins and metabolic diseases: The role of gut microbiota. *Nutrients*, 13(1), Article 206. <https://doi.org/10.3390/nu13010206>

- Shahid, S., Sultana, T., Sultana, S., Hussain, B., Al-Ghanim, K. A., Al-Bashir, F., Riaz, M. N., & Mahboob, S. (2022). Detecting aquatic pollution using histological investigations of the gills, liver, kidney, and muscles of *Oreochromis niloticus*. *Toxics*, 10(10), Article 564. <https://doi.org/10.3390/toxics10100564>
- Smorodinskaya, S., Kochetkov, N., Gavrilin, K., Nikiforov-Nikishin, D., Reznikova, D., Vatlin, A., Klimuk, A., Odorskaya, M., Nikiforov-Nikishin, A., Ponomarev, A., Marsova, M., & Danilenko, V. (2023). The effects of acute bisphenol A toxicity on the hematological parameters, hematopoiesis, and kidney histology of zebrafish (*Danio rerio*). *Animals*, 13(23), Article 3685. <https://doi.org/10.3390/ani13233685>
- Sudová, E., Piačková, V., Kroupová, H., Pijáček, M., & Svobodová, Z. (2009). The effect of praziquantel applied per os on selected haematological and biochemical indices in common carp (*Cyprinus carpio* L.). *Fish Physiology and Biochemistry*, 35(4), 599–605. <https://doi.org/10.1007/s10695-008-9269-3>
- Tasa, H., Imani, A., Moghanlou, K. S., Nazdar, N., & Moradi-Ozarlou, M. (2020). Aflatoxicosis in fingerling common carp (*Cyprinus carpio*) and protective effect of rosemary and thyme powder: Growth performance and digestive status. *Aquaculture*, 527, Article 735437. <https://doi.org/10.1016/j.aquaculture.2020.735437>
- Terzi, E., Tahiluddin, A. B., & Kadak, A. E. (2023). Evaluation of the antibacterial activity of cultivated Caucasian whortleberry (*Vaccinium arctostaphylos* L.) against fish pathogens. *Fisheries & Aquatic Life*, 31(2), 79–86. <https://doi.org/10.2478/aopf-2023-0009>
- Tessari, E. N. C., Oliveira, C. A. F., Cardoso, A. L. S. P., Ledoux, D. R., & Rottinghaus, G. E. (2006). Effects of aflatoxin B₁ and fumonisin B₁ on body weight, antibody titres and histology of broiler chicks. *British Poultry Science*, 47(3), 357–364. <https://doi.org/10.1080/00071660600756071>
- Venkataraman, P., Krishnamoorthy, G., Selvakumar, K., & Arunakaran, J. (2009). Oxidative stress alters creatine kinase system in serum and brain regions of polychlorinated biphenyl (Aroclor 1254) exposed rats: Protective role of melatonin. *Basic & Clinical Pharmacology & Toxicology*, 105(2), 92–97. <https://doi.org/10.1111/j.1742-7843.2009.00406.x>
- Wu, X., Wang, L. J., Hou, Y., Guo, R. Y., Liu, M., Yang, L., & Zhang, J. L. (2021). Different action mechanisms of low-and high-level quercetin in the brains of adult zebrafish (*Danio rerio*). *Ecotoxicology and Environmental Safety*, 223, Article 112597. <https://doi.org/10.1016/j.ecoenv.2021.112597>
- Xin, Y., Li, X., Ping, K., Xiang, Y., Li, M., Li, X., Yang, H., & Dong, J. (2024). Pesticide avermectin-induced hepatotoxicity and growth inhibition in carp: Ameliorative capacity and potential mechanisms of quercetin as a dietary additive. *Aquatic Toxicology*, 268, Article 106859. <https://doi.org/10.1016/j.aquatox.2024.106859>
- Xu, Z., Li, X., Yang, H., Liang, G., Gao, B., & Leng, X. (2019). Dietary quercetin improved the growth, antioxidation, and flesh quality of grass carp (*Ctenopharyngodon idella*). *Journal of the World Aquaculture Society*, 50(6), 1182–1195. <https://doi.org/10.1111/jwas.12663>
- Zaki, M. S., & Fawzy, O. (2012). Effect of aflatoxins B₁ on endocrine status in cat fish (*Clarias lazera*). *Life Science Journal*, 9(1), 419–422.
- Zhang, C., Jiang, D., Wang, J., & Qi, Q. (2021). The effects of TPT and dietary quercetin on growth, hepatic oxidative damage and apoptosis in zebrafish. *Ecotoxicology and Environmental Safety*, 224, Article 112697. <https://doi.org/10.1016/j.ecoenv.2021.112697>
- Zychowski, K. E., Pohlenz, C., Mays, T., Romoser, A., Hume, M., Buentello, A., Gatlin, D. M., & Phillips, T. D. (2013). The effect of NovaSil dietary supplementation on the growth and health performance of Nile tilapia (*Oreochromis niloticus*) fed aflatoxin-B₁ contaminated feed. *Aquaculture*, 376, 117–123. <https://doi.org/10.1016/j.aquaculture.2012.11.020>