PART II SCIENTIFIC AND TECHNICAL

ON THE RESPONSE OF ZOOPLANKTON IN AN ELECTRICAL FIELD

G. N. MITRA

Ministry of Food, Agriculture, C. D. and Co-operation Govt. of India, Krishibhavan, New Delhi

AND

K. P. BISWAS*

Pilot Power Fishing in Sea, Paradeep, Cuttack, Orissa.

Zooplankton showed directional movement towards the positive electrode in an electrical field till the current continued to flow. Examination of the plankton twentyfour hours after their exposure for one minute to different current densities showed considerable mortality, the percentage of which was more in the case of the larger size groups.

Introduction

Electrical fishing in ponds and tanks might have possible adverse effects on their plankton population. Threshold current density for galvanotaxis of Daphnia magna has been reported to be 0.98 by previous workers. The anodic movement of these organisms during galvanotaxis also has been observed by them. The present series of experiments were conducted in the laboratory in order to study the percentage of mortality of zooplankton due to exposure to different current densities.

MATERIALS AND METHODS

The apparatus used for studying the effect of electric current on zooplankton consisted of 11 nos of cylindrical jars of 15 1 capacity each, containing 10 1 of tap

water. The electrodes consisted of two rectangular copper plates, 15.5 cm x 8 5 cm size kept immersed in the jars at a distance of 21 cm from each other. 0.7 ml of of zooplankton collected from a tank was introduced into each of the eleven The jars were divided into five groups, two in each leaving one as control. Direct current was used as source of The voltage was reduced by a step down transformer and controlled with the help of a rheostat. The input voltage was read on voltmeter on the switch board and the current drawn measured with a mA meter, connected in the circuit. Electrical stimulation was given for one minute in each jar with variable current The only visible immediate densities. effect observed was the attraction and congregation of plankton around the positive

^{*} Present address: Directorate of Fisheries, Cuttack-7, Orissa.

TABLE I THE EFFECT OF ELECTRICAL FIELD ON ZOO-PLANKTON

Water temp: 28°C		Time of ex	Time of exposure 1 min.			Vol. of water 10 l.		
All the second s		Total		Observations at	fter 24 hours			
Expt. No.	Current density in 8	population of plankton in nos.	No. of plankton living	Nos. of plankton dead	% of living	% of dead		
		Voltage	input - 15					
1	3.2	8150	2550	5600	31	69		
2	-do-	13400	6900	6500	51.5	48.5		
		Voltage	input - 24					
1	8	11750	2850	8900	24	76		
2	-do-	10850	2650	8200	24.5	75.5		
		Voltage	input - 35					
1	15.2	19700	4000	15650	21	79		
2	-do-	12700	5200	7500	41	59		
		Voltage	input - 47					
1	19.2	11600	4150	7450	36	64		
2	-do-	14100	4900	9200	35	65		
		Voltage	input = 55					
1	23.2	13650	3700	9950	27	73		
2	-do-	15400	3400	12000	23	77		
Control	0	15000	8650	5350	64	36		

TABLE II PERCENTAGE OF MORTALITY & SURVIVAL OF INDIVIDUAL ZOOPLANKTON AFTER ELECTRICAL TREATMENT

Water temp: 28°C		Time of exposure: 1 min.				Vol. of water 10 l.	
Current	Voltage in put	Copepod		Ciadocerans		Nauplius	
density in 8		% of living	% of dead	% of living	% of dead	% of living	% of dead
3.2	15	37	37.5	13	12.5		***************************************
-do-	-do-	20	35.5	15.5	14.5	5.5	-
8	24	34	36	7	14	9	,
do-	-do-	40.5	36.5	5	13	4.5	0.5
15.2	35	30.5	46.5	5.5	3.5	14	guarantii d
-do-	-do-	35.5	45.5	8	4.5	6.5	
19.2	47	39.5	39	10.5	11	Filomotod	personal
- d o-	-do-	41	44	5	6	4	
23.2	55	44	36	6	14	Extracted	-
-do-	-do-	35,5	41	10.5	9	4	
Control	0	38.5	44.5	11.5	5.5	_	Standill his

TABLE III THE % MORTALITY OF INDIVIDUAL ZOOPLANKTON AFTER ELECTRICAL TREATMENT IN RELATION TO SIZE

Water temp. 28°C		Tim	Time of exposure 1 min.			Vol. of water 10 1		
Current density in 8	Voltage in put	Cope 1.95-0.75 mm	pods 0.74-0.49 mm	Clade 0.82-0.52 mm	0.51-0.39 mm	Nauplius 0.19 to 0.15 mm		
3.2	15	27.5	10	10.5	2			
-do-	-do-	28	7.5	11.5	3			
8	24	26	10	12	4			
-do-	-do-	31.5	5	5	8	5		
15.2	35	40	6.5	2	1.5	—		
-do-	-do-	30.5	1.5	4	.5	_		
19.2	47	31	8	5	6			
-do-	-do-	38.5	5.5	6	name of the last o			
23.2	55	29.5	6.5	14	-			
-do-	-do-	32.5	8.5	6.5	.4	4		
Control	0	30.5	14	4.5	1	harmond .		

TABLE IV THE % OF SURVIVAL OF ZOOPLANKTON AFTER ELECTRICAL TREATMENT

		the state of the s	Anna Anna Anna Anna Anna Anna Anna Anna			Nauplius
Current	Voltage		Copepods		Cladocerans	
density	in put	1.95 - 0.75	0.74 - 0.49	0.82 - 0.52	0.51 - 0.39	0.19 - 0.15
in 8	m pur	$\mathbf{m}\mathbf{m}$	$\mathbf{m}\mathbf{m}$	$\mathbf{m}\mathbf{m}$	mm	mm
3.2	15	7	30	.7	6	—
-do-	-do-	13.5	15.5	12	3.5	5.5
8	24	8	26	4	3	9
-do-	-do-	16	24.5	1	4	4.5
15.2	35	6	24.5	3	2.5	14
-do-	-do-	15.5	20	3.5	5.5	6.5
19.2	47	15	24.5	6	4.5	
-do-	-do-	-do-	26	4	1	4
23.2	55	10.5	33.5	6		
-do-	-do	8	27.5	6.5	4	4
Control	0	25	13.5	-do-	5	

electrode when the current was switched on. The scattered movements of the plankton were converted into directional movement towards the positive electrode for the whole period of flow of the current.

To observe the subsequent effect, they were allowed to stay in the jars for twenty-four hours, after which the dead planktons were separated from the living ones and fixed in formalin for further study. They

were analysed both for quantity and different species. Each species was measured for length and the percentages of mortality and survival were calculated.

RESULTS AND DISCUSSION

Zooplankton showed a directional movement towards the positive electrode so long as they were in the electric field. It was also observed that larger planktons like cladocerans and diaptomus moved faster to the positive electrodes and con-

gregrated around them, while the smaller ones took longer time for reaching the electrodes. No immediate effect was observed after the electrical field was removed.

The numbers of dead and living planktons were determined after twentyfour hours and their percentages calculated for different current densities. (Table I). It was observed that dead planktons were more than the living ones in current densities ranging from 3.2 to 23.28, except in one case where the current density was 3.28. Increased current density did not show any significant effect on the percentage of mortality. But in the control, survivors were more than the dead. difference was observed in the percentages of mortality and survival among copepods. cladocerans and nauplius from those of control (Table II). The mortality was found to depend on the size of the organisms irrespective of species (Table III). It varied from 26 to 40% in copepods of 1.95 to 0.75 mm length and 5 to 14% in those of 0.74 to 0.49 mm while in cladocerans of 0.82 to 0.52 mm it was 2 to 14%

and in those of 0.51 to 0.39 mm the figure was 2 to 8%. The percentages of survival increased with decrease in size (Table IV). Daphnia magna in an electric field have been reported to show anodic movement by some workers while some others were of opinion that these organisms moved towards the cathod during galvanotaxis. In our experiments we have observed a movement towards the positive electrode as long as the flow of current continued. The faster movement of larger planktons towards the positive electrode was possibly due to the lower voltage gradient required by them to exhibit electrotaxis. as the flow was cut off, the planktons exhibited scattered movement.

CONCLUSION

Zooplanktons like copepods and cladocerans exhibited directional movement towards positive electrode in an electric field till the current continued to flow. Exposure to electric field for one minute showed more mortality than survival after 24 hours. Larger sizes showed higher rates of mortality.