### NOTE III

# QUALITY LOSS IN PRAWNS DUE TO DOUBLE FREEZING

Icing is the practice for preserving prawns on board fishing boats in India. Majority of these boats need to preserve the catch only for a few hours because of the short duration of the fishing trip. However, with the anticipated introduction of a considerable number of bigger fishing vessels which can remain in the fishing ground for longer periods-more than fortnight - preservation methods other than icing are required to retain prime quality. Freezing and cold storage of whole prawns on board followed by thawing and processing on land is a possible proposition. The extent of quality loss in prawns during these operations is one of the important points to be considered. Hence, laboratory scale studies were undertaken on double freezing of prawns and the results are dealt with in this communication.

Prawns - m. dobsoni and p. stylifera collected from Cochin were washed in sea water and half of the lot was frozen whole at -40°C and the other half was converted to Headless and Peeled and Deveined (2 kgs. and 0.3 kg. blocks) and frozen at -0°C, the latter served as the once frozen samples. The whole frozen material was thawed in running water after a month, then processed as HL and PD forms, this served as the twice frozen In order to collect information blocks. regarding drip loss, some blocks were kept in airtight polythene bags, then thawed in running water and the material quality and drip loss were assessed. The frozen blocks were drawn at intervals, thawed at 4°C for 17 hours and the material was analysed for total nitrogen, and free  $\alpha$ -amino nitrogen in addition to the organoleptic and physical characteristics by the methods given earlier. (C. George, 1973).

Results of experiments carried out with m. dobsoni are given in tables. The thawed yield dropped from 97 to 93% in 40 weeks of storage in the case of once frozen sample and from 94 to 90% in the case of HL double frozen sample. The initial thawed yields in PD once frozen and double frozen samples are 86.1 and 80.3% respectively; which decreases to 80 and 70% in 40 weeks of storage. percentage of pieces showing black spots increases during frozen storage and it is more in double frozen samples as is seen in the tables. The cooked characteristics, texture and flavour shows appreciable difference between the once and double frozen lots. The percentage loss of total nitrogen is from 1.4 to 3.2% in HL once frozen lot and from 2.5 to 5.1% in double frozen HL lot in 40 weeks of storage and from 2.8 to 6.6% in PD once frozen lot and from 8.0 to 14.2% in double frozen PD lot. The free  $\alpha$ -amino nitrogen shows marked change during frozen storage and it varies from 1.3 to 22.6% in once frozen control HL lot and from 3.1 to 43.1% in HL double frozen lot and from 32.5 to 43.6% in PD once frozen lot and from 46.8 to 57.1% loss in PD double frozen lot in 40 weeks of storage.

Experiments conducted with p, stylifera shows similar pattern of changes, but black spot development was absent through-

## Quality loss in prawns due to double freezing

# CHEMICAL, PHYSICAL AND ORGANOLEPTIC CHARACTERISTICS OF ONCE FROZEN (a) AND, DOUBLE FROZEN (b) PRAWNS M. DOBSONI:

TABLE I

HL - Form

Weeks of frozen storage	Percentage loss of total Nitrogen		Percentage loss of α—NH <sub>2</sub> N		Percentage showing blackening		Thawed yield %	
	(a)	(b)	(a)	(b)	(a)	(b)	(a)	(b)
0	1.4	2.5	1.3	3.1	6.95	13.35	97.0	94.0
15	2.1	3.9	5.6	9.9	17.95	19.23	96.4	93.2
30	2.5	4.4	18.7	28.9	17.82	25.2	95.0	92.0
40	3.2	5.1	22.6	43.1	33.3	65.0	93.4	90.6

## M. dobsoni. HL Form.

Weeks of	Appearance		Cooked characteristics				
frozen			Texture		Flavour		
storage	(a)	(b)	(a)	(b)	(a)	(b)	
0	Good	G. F.	Soft & firm	F.	G. F.	F.	
15	G. F.	F.	F.	S. tough	F.	F.	
30	Fair	F. P.	F.	Mod. tough	F.	F. P.	
40	F. P.	Poor	Sl. tough	Tough	F. P.	Poor	

TABLE II

PD - Form

Percenta total N	ge loss of itrogen.	Percenta $\alpha - NF$	ige loss of H2—N.		ge showing kening	Thawed %	yield
(a)	(b)	(a)	(b)	(a)	(b)	(a)	(b)
2.83	8.0	32.5	46.8	0.0	11.1	86.1	80.3
4.24	9.4	34.1	48.4	1.2	17.0	81.3	76.8
5.35	12.2	42.0	56.3	3.4	21.8	80.9	74.7
6.67	14.2	43.6	57.1	4.0	31.7	80.0	69.7

### Quality loss in prawns due to double freezing

M. dobsoni. PD Form

		Cooked characteristics				
Appearance		Тех	ture	Flavour		
(a)	(b)	(a)	(b)	(a)	(b)	
G. F.	F.	Soft & firm	F.	G. F.	F.	
F.	F.P.	Sl. hard	Sl. hard	F.	F. P	
F.	F.P.	Sl. tough	Mod tough	F.	F. P	
Poor	Poor	Sl. tough	Mod tough	P.	Poor	

out the storage period, as this phenomenon is of species specific.

The foregoing studies clearly indicate that some changes are brought out by refreezing and when compared to the quality loss in icing, it is less. The major quality defects are high drip loss and high rate of blackening.

I am much grateful to Shri. M R.

Nair, Fishery Scientist of this Institute for his valuable suggestions during the course of this work.

### Reference:

Chinnamma George 1973 Technological aspects of preservation and processing of edible shell fishes. III. Freezing characteristics of crab. Fishery Technology, Vol. 10, No. 1.

Central Institute of Fisheries Technology, Ernakulam, Cochin-11.

Chinnamma George