

## Water Soluble Nitrogenous Component from Squilla (*Orato squilla nepa*)\*

A. LEKSHMY NAIR and P. V. PRABHU

Central Institute of Fisheries Technology, Cochin-682 029

A water soluble hygroscopic powder has been isolated from squilla in good yield, ranging from 3.5 to 5.0 per cent of the fresh raw material, by a simple direct method. The process consists of homogenising squilla with an equal quantity of water, removal of chitinous matter from the slurry by filtration, heating the filtrate at 0.7 kg/sq.cm steam pressure for 15–20 minutes, removal of the precipitated protein by filtration and concentration and final drying in vacuum of the filtered cooled liquor. The pale brown powder so obtained consists mainly of peptones and proteoses and has been found to be comparable to BDH peptone for growth of bacteria, ability to serve as source for tryptophan for indole production and to provide substrate for the production of hydrogen sulphide. Comparative studies have been made on similar water soluble fractions from two species of prawns, namely, *Metapenaeus affinis* and *Parapenaeopsis stylifera*.

Squilla (*Orato squilla nepa*) is available in plenty throughout the west coast of India and is caught along with prawn in trawl catches. Although exact data on the total landings of squilla is not available, it is present as a by-catch in sufficiently large quantity to warrant development of suitable methods for its utilisation. Since it does not contain appreciable amount of meat and great difficulty is encountered in separating the meat from the shell, the squilla available in India has not been put to any commercial use. Recently, attempts have been made for development of methods for better utilisation of squilla. Madhavan & Nair (1975) have described a method for preparation of chitosan from squilla. A simple method for the isolation of protein from squilla without any chemical treatment has been reported by Garg *et al.* (1977). During earlier studies on squilla, it was observed that a major portion of the nitrogen present in squilla came out in water soluble fraction during the precipitation and separation of squilla protein. The present paper reports the yield and composition of the water soluble nitrogenous

fraction of squilla. A comparative evaluation has also been carried out between the water soluble nitrogenous components of squilla and two species of prawns, *Metapenaeus affinis* and *Parapenaeopsis stylifera*.

### Materials and Methods

Squilla caught by the Central Institute of Fisheries Technology fishing boats from the sea off Cochin was washed, iced immediately and brought to the laboratory. It was stored in crushed ice and used the next day or was frozen and stored at –20°C until taken out for further processing. The prawns used in the experiments were purchased fresh from the local market and kept heavily iced for about a day before taking for experiments the next day. All the samples used were in organoleptically good condition.

Isolation of the chitinous matter and protein from the samples was done according to the method described earlier (Garg *et al.*, 1977). The water soluble portion obtained after separation of the precipitated protein was concentrated on a water bath and finally dried under vacuum (750 mm of Hg at 45–50°C). A pale brown hygroscopic powder was obtained. Decolourisation of the cooled filtrate with activated

\* Paper presented at the symposium on "By-products from Food Industries—Utilization and Disposal" 29-30 May, 1980 at Central Food Technological Research Institute, Mysore.

charcoal was also attempted to improve the colour of the final product, with little success. Total nitrogen and chitin nitrogen of raw material were estimated by the method of Garg *et al.* (1977). Coagulable nitrogen, proteose nitrogen and peptone nitrogen present in the water soluble fraction were determined according to Winton & Winton (1958).

The samples prepared from squilla were incorporated into various bacteriological media, both liquid and solid and their

growth promoting properties were studied using selected bacterial strains.

### Results and Discussion

Results of a few typical trials on the isolation of water soluble nitrogenous component from squilla and prawn are summarised in Table 1. Yield of the total water soluble fraction from squilla varied between 3.5 and 5% of the fresh raw material and that from the two prawns between 1.8 and 2% respectively. Table 2 presents

**Table 1.** Yield of water soluble nitrogen fraction from squilla and prawn

| Raw material                    | Wt. of fresh material | Wt. of vacuum dried powder from water soluble fraction | Yield |
|---------------------------------|-----------------------|--------------------------------------------------------|-------|
|                                 | g                     | g                                                      | %     |
| Squilla Batch 1                 | 1000                  | 50.0                                                   | 5.0   |
| Squilla Batch 2                 | 1000                  | 35.0                                                   | 3.5   |
| Squilla Batch 3                 | 1000                  | 42.0                                                   | 4.2   |
| Squilla Batch 4                 | 1000                  | 40.0                                                   | 4.0   |
| Squilla Batch 5                 | 1000                  | 35.0                                                   | 3.5   |
| Squilla Batch 6                 | 1000                  | 38.0                                                   | 3.8   |
| Prawns ( <i>P. styliifera</i> ) | 1000                  | 18.8                                                   | 1.9   |
| Prawns ( <i>M. affinis</i> )    | 1000                  | 21.0                                                   | 2.1   |

**Table 2.** Yield and distribution of different nitrogen components in the water soluble nitrogen fraction from squilla and prawns

| Raw material                    | Nitrogen present in raw material |           |           | Nitrogen recovered in the water soluble fraction | Percentage of nitrogen recovered in the water soluble fraction | Percentage yield in the water soluble fraction of |                   |                           |
|---------------------------------|----------------------------------|-----------|-----------|--------------------------------------------------|----------------------------------------------------------------|---------------------------------------------------|-------------------|---------------------------|
|                                 | Total TN                         | Chitin CN | TN-CN A-B |                                                  |                                                                | Coagulable nitrogen                               | Proteose nitrogen | Proteose peptone nitrogen |
|                                 | mg                               | mg        | mg        |                                                  |                                                                |                                                   |                   |                           |
| Squilla                         | 1530                             | 180       | 1350      | 640                                              | 47.4                                                           | Not detectable                                    | 13.0              | 67.7                      |
| Squilla                         | 1730                             | 210       | 1520      | 640                                              | 42.1                                                           | Not detectable                                    | 17.6              | 60.7                      |
| Squilla                         | 1310                             | 200       | 1110      | 580                                              | 52.3                                                           | Traces                                            | 15.0              | 70.0                      |
| Squilla                         | 1710                             | 200       | 1510      | 730                                              | 48.3                                                           | Not detectable                                    | 14.1              | 69.5                      |
| Prawns ( <i>P. styliifera</i> ) | 1730                             | 100       | 1630      | 550                                              | 33.7                                                           | 16.5                                              | 2.0               | 40.3                      |
| Prawns ( <i>M. affinis</i> )    | 2150                             | 100       | 2050      | 550                                              | 26.8                                                           | 16.2                                              | 8.6               | 29.9                      |

the relationship of nitrogen recovered in the water soluble fraction to the total nitrogen present in the raw material excluding the chitin nitrogen. Distribution of the individual nitrogenous components in the aqueous extract after removal of protein is shown in Table 2. Recovery as water soluble nitrogen fraction varied between 42–52% in the squilla and between 26.8 and 33.8% in prawns. It is seen from Table 2 that the water soluble fraction from squilla consists mainly of peptones and proteoses and that coagulable protein nitrogen was not detectable in this fraction. In the case of prawns appreciable amounts of coagulable proteins are present and peptones and proteoses are available to a limited extent.

Studies on the growth promoting properties of the samples from squilla have shown that these compared very favourably with Difco and BDH peptones (Iyer, K.M. & Rao, C.C.P., personal communication) in their ability to promote bacterial

growth, to serve as source for tryptophan for indole production and to provide a substrate for the production of hydrogen sulphide. The investigations on the effectiveness of the squilla product for incorporation in bacteriological media is not yet completed and the results reported here should be regarded as preliminary.

The authors are thankful to the late Shri G. K. Kuriyan, Director, Central Institute of Fisheries Technology, Cochin for permission to publish the paper.

### References

- Garg, D. K., Lekshmy Nair, A. & Prabhu, P. V. (1977) *Fish. Technol.* **14**, 53
- Madhavan, P. & Ramachandran Nair, K. G. (1975) *Fish. Technol.* **12**, 81
- Winton, A. L. & Winton, K. B. (1958) *The Analysis of Foods*, p. 800, John Wiley and Sons, Inc., New York