



## District level crop yield estimation with reduced number of crop cutting experiments

KAUSTAV ADITYA\*, HUKUM CHANDRA, PRADIP BASAK, VANDITA KUMARI and SHRILA DAS

ICAR-Indian Agricultural Statistics Research Institute, New Delhi 110 012, India

Received: 17 September 2019; Accepted: 27 September 2019

### ABSTRACT

It was observed that around 1300000 Crop Cutting Experiments (CCE) were conducted in India every year to find out the crop yield estimates of several major and minor crops under General Crop Estimation Surveys (GCES). Due to shortage of manpower and huge bulk of work day by day the data quality is becoming questionable. To tackle this problem, a pilot study was conducted by ICAR-IASRI, New Delhi sponsored by Directorate of Economics and Statistics (DES), Ministry of Agriculture and Farmers Welfare (MoA & FW), Govt. of India to generate district level estimates of major crop yield from a reduced sample size of villages selected from the states. With the reduction in number of villages, the problem of no sample size in some districts were faced during the study where common design based estimates of crop yield cannot be generated. To tackle this problem Aggregate level Small Area Estimation (SAE) was used to tackle this problem. The results obtained from this study in the state of Uttar Pradesh for two major crops, i.e. rice and wheat for two seasons, i.e. *kharif* and *rabi* of Agriculture Year 2015-16 and for paddy in Assam for *kharif* of the Agriculture Year (AY) 2015-16 in India were discussed. The yield estimates were compared with the estimates released under GCES for AY 2015-16. It was found that the estimates obtained from reduced sample size of number of CCEs w.r.t. GCES, produced similar estimates with acceptable level of precision.

**Keywords:** Agricultural statistics, Crop cutting experiments, Pilot survey, sample Size, Small area estimation

In view of the predominant position of the Agricultural Sector, collection and maintenance of Agricultural Statistics assume great importance. The yield rates of principal crops are estimated through GCES. Under the GCES, CCEs are done in a sample of Timely Reporting Scheme (TRS) villages by one or more state agencies (Report of Expert Committee on Agricultural Statistics 2011). The intention was to generate estimates of per ha yields of various crops within a reasonable, specified margin of standard error. During past few agricultural years, it was found that around 1300000 CCEs covering 52 food and 16 non-food crops were conducted every year in different States/UTs in India. Therefore a study was carried out to evaluate the possibility of generating reliable estimate of major crop yield from a reduced number of CCEs. The results of this study were based on the data collected independently in state of Uttar Pradesh for two major crops, i.e. rice and wheat for two seasons, i.e. *kharif* and *rabi* of Agriculture Year 2015-16 by the Department of Agriculture and Crop Insurance, Lucknow, Govt. of Uttar Pradesh and for paddy in Assam for *kharif* of the Agriculture Year 2015-16 by Directorate of

Economics and Statistics, Guwahati, Govt. of Assam. The results were compared with the estimates released under GCES for the year 2015-16.

### MATERIALS AND METHODS

Under this study, a survey was undertaken in 5 states of India namely Assam, Uttar Pradesh, Odisha, Karnataka and Gujarat. The survey was conducted for the Agricultural year 2015-16 in India for both *kharif* and *rabi*. In this paper we will concentrate on two states namely Assam and Uttar Pradesh. The total number of villages and number of crop cutting experiments are around 6% of villages in the state of Assam and 10% of villages in Uttar Pradesh with respect to the sample size selected under TRS scheme. Accordingly, the total number of CCEs will also get reduced in the same proportion.

To efficiently implement the proposed sample size to generate direct estimates of crop yield, the sampling design adopted is that of stratified multi-stage random sampling as detailed below:

- i. District in each participating State has been considered as a stratum.
- ii. 50% of the Tehsils/Taluks in a district have been selected as First stage Sampling Units (FSUs) by simple random sampling without replacement (SRSWOR).

\*Corresponding author e-mail: [Kaustav.aditya@icar.gov.in](mailto:Kaustav.aditya@icar.gov.in)

Villages within a FSU are taken as Second stage Sampling Units (SSUs), and within each SSU, survey/sub-survey numbers have been taken as Third stage Sampling Units (TSUs). Experimental plot of specified size in the selected survey/sub-number is considered the ultimate stage unit for yield estimation. The SSUs and TSUs have been selected using SRSWOR. For construction of frame for selection of the CCE plots, 100 survey numbers have been selected in the form of 20 clusters of 5 survey numbers each within a selected village.

- iii. Each selected survey number has been visited by the primary worker for plot to plot enumeration of crop during *kharif* and *rabi* seasons of AY 2015-16.
- iv. Crop-wise frame for selections of survey number for CCE has been generated on the basis of 100 survey numbers selected randomly. For each major crop, two survey/sub-survey numbers have been selected for conducting CCEs using SRSWOR.

Selection of sample units up to the level of second stage units (SSUs) namely villages has been done by IASRI, New Delhi, whereas the selection of third stage units (TSUs), namely survey/sub-survey numbers for both area enumeration and CCEs, have been done by concerned State field officials. For field data collection, one schedule for preparation of frame and two schedules for crop cutting experiments have been designed. The methodology for estimation of average yield at district level using design based estimator was derived from Sukhatme *et al.* (1984) and Singh *et al.* (1996) while for Small Area Estimation (SAE) the Aggregate Level SAE method (EBLUP model) proposed by Fay *et al.* (1979) was used.

## RESULTS AND DISCUSSION

In the state of Uttar Pradesh data on two major crops, viz. Rice and Wheat were collected by the state agency in two crop seasons *Kharif* and *Rabi* respectively during Agriculture year 2015-16. After receiving the data, while cleaning and compiling the data it was found that very less or no data on CCE was recorded in 10 districts in case of Rice in *Kharif* 2015-16 and in 7 districts in case of Wheat in *Rabi* 2015-16 out of total 75 districts. In the

state of Assam in 3 districts very scanty data was recorded for Rice in *Kharif* 2015-16 out of total 26 districts. Due to conduct of less number of CCEs than the proposed, the sample size got reduced and the percentage Coefficient of Variation (% CV) of the proposed estimates got increased in few districts. Further, due to reduction of number of CCEs, in some districts sufficient sample size was not found to produce the district level direct estimates of the crop yield. Those districts are treated as the non-sample districts and the estimates are predicted using EBLUP estimator along with the estimates of all other districts with improved precision. For fitting SAE model, the district level fertilizer consumption data of the year 2015-16 for the state of Uttar Pradesh and Assam, released by Fertilizer Statistics (2015-16) was used as auxiliary information. The data on total fertilizer dose of Nitrogen (N), Phosphorus (P) and Potassium (K) fertilizers was used as the auxiliary information for construction of the SAE estimator. It was found that the correlation between the yield of rice and fertilizer dose ranged between 0.1-0.2 in the state of Uttar Pradesh, while in case of wheat the same resulted around 0.1 and in Assam the correlation was found in the tune of around 0.05. The district level yield estimates are generated using proposed design based methodology and is found to be almost same with the estimates released under the GCES scheme with acceptable % CV in the respective states in the AY 2015-16 for the sample districts. In the state of Uttar Pradesh estimates of rice and wheat was generated, whereas in Assam estimates of Paddy yield was generated. The average number of CCEs conducted for Rice crop in the state of Uttar Pradesh under this study was 42, whereas the same for wheat is 48 and in the state of Assam the average number of CCEs conducted under this study for winter paddy is only 10. Now the average number of CCEs at district level conducted under GCES for major crops like Paddy/Rice and wheat are in the tune of 100-120. So it can be seen that in the state of Uttar Pradesh the number of CCEs proposed in this study is less than half the sample size in GCES while in the state of Assam it's only around 10%.

A close perusal of data (Table 1) depicts that the % CV of the design based estimator of Rice varies from 1.3-

Table 1 District-wise Estimates of Yield (kg/ha) with % CV for Rice and Wheat crop in Uttar Pradesh for AY 2015-16.

| District     | Rice  |      |       | Wheat |      |       |      |       |      |      |
|--------------|-------|------|-------|-------|------|-------|------|-------|------|------|
|              | Yield | % CV | EBLUP | % CV  | GCES | Yield | % CV | EBLUP | % CV | GCES |
| Agra         |       |      | 2181  | 22.93 | 2399 | 2491  | 6.78 | 2491  | 0.52 | 3226 |
| Aligarh      | 2163  | 1.89 | 2164  | 1.89  | 2091 | 3400  | 2.70 | 3400  | 0.29 | 3177 |
| Allahabad    | 2384  | 4.08 | 2377  | 4.04  | 1899 | 2379  | 6.83 | 2379  | 0.55 | 1912 |
| Ambedkar Ngr | 2052  | 6.58 | 2059  | 6.36  | 2111 | 2934  | 1.66 | 2934  | 0.24 | 3183 |
| Amethi       | 1973  | 3.89 | 1978  | 3.84  | 1967 | 2176  | 9.17 | 2176  | 0.64 | 2375 |
| Amroha       | 2103  | 2.97 | 2104  | 2.95  | 2257 | 3316  | 6.17 | 3316  | 0.42 | 2962 |

Contd.

Table 1 (Continued)

| District    | Rice  |       |       |       |      | Wheat |      |       |       |      |
|-------------|-------|-------|-------|-------|------|-------|------|-------|-------|------|
|             | Yield | % CV  | EBLUP | % CV  | GCES | Yield | % CV | EBLUP | % CV  | GCES |
| Auraiya     | 3022  | 3.93  | 2974  | 3.9   | 2820 | 3434  | 2.98 | 3434  | 0.29  | 3085 |
| Azamgarh    | 2241  | 7.02  | 2240  | 6.74  | 1970 | 2753  | 3.74 | 2753  | 0.36  | 2676 |
| Badaun      | 1922  | 3.35  | 1928  | 3.32  | 1847 | 4091  | 3.33 | 4091  | 0.29  | 4140 |
| Baghpat     | 2747  | 2.43  | 2736  | 2.41  | 2746 | 2734  | 5.81 | 2734  | 0.48  | 2661 |
| Bahraich    | 1658  | 5.29  | 1676  | 5.13  | 1888 | 2695  | 8.62 | 2695  | 0.56  | 2707 |
| Ballia      | 2263  | 4.32  | 2260  | 4.25  | 1694 | 3242  | 9.02 | 3242  | 0.52  | 2389 |
| Balrampur   | 1508  | 9.03  | 1555  | 8.49  | 1803 | 1564  | 9.39 | 1564  | 0.77  | 1200 |
| Banda       | 1449  | 5.68  | 1468  | 5.52  | 1295 | 3023  | 4.09 | 3023  | 0.36  | 2973 |
| Barabanki   | 2675  | 6.43  | 2625  | 6.21  | 2751 |       |      | 2883  | 29.26 | 2930 |
| Bareilly    | 2029  | 4.94  | 2037  | 4.81  | 2608 | 2738  | 3.24 | 2738  | 0.33  | 2615 |
| Basti       | 1807  | 1.76  | 1809  | 1.77  | 1880 | 3244  | 9.43 | 3244  | 0.52  |      |
| Bhadrohi    | 1938  | 6.05  | 1952  | 6.01  |      | 2936  | 2.14 | 2936  | 0.27  | 3100 |
| Bijnor      | 2546  | 2.65  | 2540  | 2.64  | 2590 | 3668  | 3.49 | 3668  | 0.31  | 2831 |
| Bulandshahr | 2432  | 4.34  | 2422  | 4.25  | 2688 | 1804  | 6.06 | 1804  | 0.55  | 3867 |
| Chandauli   | 3586  | 6.82  | 3303  | 6.69  | 3085 | 1487  | 6.22 | 1487  | 0.67  | 1918 |
| Chitrakoot  | 867   | 12.34 | 922   | 11.39 | 1156 | 2493  | 2.78 | 2493  | 0.32  | 1414 |
| Deoria      | 1172  | 9.73  | 1222  | 9.08  | 776  | 2874  | 3.43 | 2874  | 0.35  | 2545 |
| Etah        | 2171  | 4.77  | 2170  | 4.71  | 1545 | 3370  | 1.28 | 3370  | 0.21  | 2901 |
| Etawah      | 2306  | 3.80  | 2301  | 3.74  | 2466 | 2569  | 6.89 | 2569  | 0.51  | 3326 |
| Faizabad    | 2040  | 5.35  | 2046  | 5.23  | 2283 | 3408  | 3.93 | 3408  | 0.35  | 2854 |
| Farrukhabad | 2563  | 4.61  | 2539  | 4.53  | 2246 | 1617  | 3.78 | 1617  | 0.49  | 3171 |
| Fatehpur    | 1703  | 5.79  | 1722  | 5.63  | 2047 | 3331  | 2.87 | 3331  | 0.32  | 2485 |
| Firozabad   |       |       | 2311  | 22.55 | 2397 | 3269  | 6.05 | 3269  | 0.43  | 3287 |
| GTBnag      | 2789  | 3.11  | 2768  | 3.07  | 2616 | 3397  | 6.95 | 3397  | 0.44  | 3423 |
| Ghaziabad   | 2670  | 3.96  | 2645  | 3.93  | 2568 | 2105  | 7.17 | 2105  | 0.57  | 3610 |
| Gazipur     | 2016  | 7.06  | 2029  | 6.75  | 2165 | 2459  | 4.89 | 2459  | 0.45  | 2386 |
| Gonda       | 2174  | 4.19  | 2176  | 4.14  | 2061 | 2246  | 6.23 | 2246  | 0.53  | 2766 |
| Gorakhpur   | 1541  | 8.84  | 1586  | 8.32  | 1621 | 1351  | 6.01 | 1351  | 0.67  | 2519 |
| Hamirpur    |       |       | 2124  | 23.68 | 1156 | 3746  | 2.24 | 3746  | 0.24  | 1752 |
| Hapur       | 2784  | 4.79  | 2738  | 4.71  | 2836 | 3183  | 2.73 | 3183  | 0.28  | 3840 |
| Hardoi      | 2318  | 2.51  | 2317  | 2.51  | 2256 | 3684  | 2.72 | 3684  | 0.27  | 3001 |
| Hathras     | 2020  | 4.02  | 2023  | 3.95  | 1966 | 2965  | 5.34 | 2965  | 0.44  | 3495 |
| Jalaun      |       |       | 2181  | 22.93 | 1795 | 2336  | 7.53 | 2336  | 0.56  | 1880 |
| Jaunpur     | 1787  | 5.17  | 1802  | 5.05  | 1760 |       |      | 2879  | 29.24 | 2661 |
| Jhansi      |       |       | 2205  | 22.72 | 1683 | 3590  | 2.69 | 3590  | 0.28  | 1918 |
| Kannauj     | 2020  | 3.69  | 2022  | 3.66  | 2351 |       |      | 2821  | 29.49 | 3250 |
| Kanpur(D)   | 2338  | 2.23  | 2336  | 2.18  | 2635 | 3299  | 7.97 | 3299  | 0.49  | 2124 |
| Kanpur (N)  | 2296  | 5.41  | 2286  | 5.29  | 2274 | 3078  | 5.82 | 3078  | 0.42  | 3012 |
| Kasganj     |       |       | 2120  | 23.77 | 1917 | 2266  | 4.53 | 2266  | 0.44  | 3084 |

Contd.

Table 1 (Concluded)

| District      | Rice  |       |       |       |      | Wheat |       |       |       |      |
|---------------|-------|-------|-------|-------|------|-------|-------|-------|-------|------|
|               | Yield | % CV  | EBLUP | % CV  | GCES | Yield | % CV  | EBLUP | % CV  | GCES |
| Kaushambi     | 1506  | 6.43  | 1529  | 6.21  | 2119 | 2555  | 2.01  | 2555  | 0.27  | 2176 |
| Kushinagar    | 1691  | 7.34  | 1722  | 7.03  | 1583 | 3968  | 9.08  | 3967  | 0.48  | 2336 |
| Lakhimpur     | 2762  | 2.21  | 2757  | 2.18  | 2629 | 3177  | 2.25  | 3177  | 0.25  | 3393 |
| Lalitpur      |       |       | 2192  | 22.81 | 635  | 2236  | 7.29  | 2236  | 0.58  | 1738 |
| Lucknow       | 2152  | 4.83  | 2150  | 4.83  | 2195 | 676   | 10.86 | 676   | 1.33  | 2619 |
| Maharajganj   | 2392  | 5.79  | 2377  | 5.64  | 2464 |       |       | 2820  | 29.51 | 2996 |
| Mahoba        |       |       | 2148  | 23.32 | 1156 | 3494  | 6.69  | 3494  | 0.43  | 787  |
| Mainpuri      | 2460  | 4.61  | 2445  | 4.54  | 2636 | 3129  | 7.26  | 3129  | 0.48  | 3332 |
| Mathura       | 2643  | 6.62  | 2591  | 6.41  | 2224 | 5595  | 0.35  | 5595  | 0.07  | 3453 |
| Mau           | 1919  | 2.32  | 1921  | 2.29  | 1456 |       |       | 2844  | 29.27 | 2503 |
| Meerut        | 2881  | 3.84  | 2848  | 3.79  | 2711 | 1708  | 7.13  | 1708  | 0.64  | 3874 |
| Mirzapur      | 1545  | 7.24  | 1574  | 6.92  | 1730 | 3102  | 1.97  | 3102  | 0.26  | 1829 |
| Moradabad     | 2431  | 3.15  | 2425  | 3.13  | 2338 | 3698  | 1.96  | 3698  | 0.22  | 3142 |
| Muzaffarnagar | 2609  | 2.35  | 2603  | 2.34  | 2596 | 3990  | 1.84  | 3990  | 0.23  | 3565 |
| Pilibhit      | 3480  | 2.89  | 3429  | 2.89  | 3010 | 1679  | 5.24  | 1679  | 0.54  | 3877 |
| Pratapgarh    | 1649  | 7.18  | 1677  | 6.86  | 1799 | 1884  | 5.79  | 1884  | 0.53  | 2188 |
| Raebareli     | 1951  | 6.31  | 1963  | 6.31  | 2126 | 3670  | 4.80  | 3670  | 0.35  | 1969 |
| Rampur        | 2339  | 2.25  | 2337  | 2.22  | 2260 |       |       | 2767  | 30.67 | 3423 |
| Saharanpur    | 2403  | 2.44  | 2400  | 2.37  | 2264 | 3517  | 1.80  | 3517  | 0.23  | 3579 |
| Sambhal       | 2204  | 2.72  | 2204  | 2.72  | 2301 |       |       | 2828  | 29.41 | 3070 |
| SantKabirngr  | 1847  | 10.85 | 1887  | 9.91  | 1755 | 2471  | 4.86  | 2471  | 0.45  | 2542 |
| Shahjahanpur  | 3257  | 1.31  | 3250  | 1.29  | 3149 | 2442  | 9.79  | 2442  | 0.61  | 3509 |
| Shamli        | 1903  | 7.72  | 1920  | 7.34  | 2498 | 4045  | 2.71  | 4045  | 0.25  | 4071 |
| Shravasti     | 1668  | 7.91  | 1698  | 7.54  | 1950 | 2312  | 6.02  | 2312  | 0.52  | 2656 |
| Sidharthnagar | 2025  | 6.59  | 2035  | 6.34  | 2262 | 3241  | 3.76  | 3241  | 0.34  | 3342 |
| Sitapur       | 2025  | 6.59  | 2320  | 8.36  | 2158 | 2400  | 2.12  | 2400  | 0.29  | 2485 |
| Sonbhadra     | 1905  | 19.78 | 1985  | 15.31 | 1487 | 1334  | 13.15 | 1334  | 0.97  | 1590 |
| Sultanpur     | 2042  | 4.92  | 2047  | 4.84  | 2077 | 2765  | 5.55  | 2765  | 0.43  | 2838 |
| Unnao         | 1650  | 4.99  | 1665  | 4.86  | 1584 | 2567  | 1.78  | 2567  | 0.27  | 2222 |
| Varanasi      | 2045  | 9.08  | 2054  | 8.52  | 2256 | 2007  | 6.71  | 2007  | 0.60  | 2414 |

19.78 with an average of 5.35, whereas the % CV of the SAE estimates (EBLUP) varies from 1.29-15.31 for all the sampled districts and 22.55-23.77 for non-sampled districts. For wheat crop, it can be seen that, the % CV of the design based estimator varies from 0.35-10.86 with an average of 5.10, whereas the % CV of the SAE estimates varies from 0.07- 1.33 for all the sampled districts and 29.24-30.67 for non-sampled districts. For the state of Assam it is visible that the % CV of the design based estimator varied from 1.83-16.51 with an average of 8.07, whereas the % CV of the SAE estimates varies from 1.83-10.97 for all the sampled districts with an average of 6.74 (Table 2). For the non-sampled districts, the % CV of the SAE estimator lies in the

range of 15.84-16.63. As the % CV of the GCES estimates are not available hence no conclusion can be drawn about the acceptability of these estimates where from our study for around 90% of districts of Uttar Pradesh we are able to generate precise estimates of yield of Rice and Wheat crop from around 40% of CCEs on an average conducted under GCES in each district. In case of Assam, for around 96% of districts, we are able to generate precise estimates of yield of Wheat crop from around 10% of CCEs on an average conducted under GCES in each district.

The CCE is termed as a gold standard method for estimation of crop yield by FAO (Sud *et al.* 2017) and it cannot be replaced with any other existing methods. Hence,

Table 2 District wise estimate of Yield (kg/ha) along with % CV in the state of Assam for AY 2015-16.

| District    | Yield   | % CV  | EBLUP   | % CV  | GCES |
|-------------|---------|-------|---------|-------|------|
| Baksa       | 2975.30 | 4.00  | 3009.97 | 3.88  | 3267 |
| Barpeta     | 2193.20 | 13.57 | 2392.52 | 10.05 | 2272 |
| Bongaigaon  | 2595.12 | 10.56 | 2803.90 | 8.97  | 2510 |
| Cachar      | 3169.66 | 5.76  | 3256.39 | 5.39  | 3345 |
| Chirang     | 3087.56 | 4.29  | 3046.86 | 4.20  | 2679 |
| Darrang     | 3236.45 | 11.81 | 3278.41 | 9.94  | 3469 |
| Dhemaji     | 2465.29 | 3.10  | 2241.58 | 3.07  | 2293 |
| Dhubri      | 3989.76 | 5.61  | 3943.58 | 5.41  | 2658 |
| Dibrugarh   | 4350.06 | 3.32  | 4282.31 | 3.31  | 3042 |
| Dima Hassao |         |       | 3620.83 | 15.84 | 3766 |
| Goalpara    | 4061.14 | 16.29 | 3749.91 | 10.97 | 3295 |
| Golaghat    | 3271.85 | 7.52  | 3353.06 | 6.84  | 3438 |
| Hailakandi  | 3514.34 | 8.17  | 3474.35 | 7.55  | 3725 |
| Jorhat      | 3547.44 | 10.03 | 3112.68 | 8.62  | 3404 |
| K. Anglong  |         |       | 2807.98 | 14.46 | 2575 |
| Kamrup (M)  |         |       | 3366.51 | 16.63 | 3395 |
| Karimganj   | 3594.01 | 16.51 | 3681.08 | 10.77 | 2635 |
| Kokrajhar   | 2458.64 | 8.40  | 2610.83 | 7.53  | 2595 |
| Lakhimpur   | 3789.12 | 5.06  | 3754.29 | 4.86  | 3138 |
| Morigaon    | 2718.85 | 16.02 | 2877.89 | 10.88 | 2960 |
| Nagaon      | 3015.46 | 4.70  | 3050.81 | 4.62  | 3377 |
| Nalbari     | 2957.99 | 5.04  | 2955.38 | 4.89  | 3192 |
| Sivasagar   | 3200.00 | 1.83  | 3185.68 | 1.83  | 3155 |
| Sonitpur    | 3539.60 | 9.05  | 3304.97 | 7.99  | 3294 |
| Tinsukia    | 2989.91 | 8.55  | 3152.84 | 7.43  | 2890 |
| Udalguri    | 3343.35 | 6.47  | 3360.67 | 6.09  | 3298 |

to reduce the workload and to improve the data quality with existing man power, the only way left was to reduce the total number of crop cutting experiments conducted under the GCES and we can conclude that our study resulted into success. Further, the use of SAE technique has improved the precision of the estimates. Hence we can conclude that estimates of crop yield with acceptable precision can be generated from less than 50% CCEs conducted under GCES for major food grain crops.

#### ACKNOWLEDGEMENTS

We acknowledge the contributions made by Late Mr Sushil Kumar, Scientist, ICAR-IASRI, New Delhi in data analysis of this paper.

#### REFERENCES

- Fay R E and Herriot R A. 1979. Estimation of income from small places: An application of James-Stein procedures to census data. *Journal of the American Statistical Association* **74**: 269-77.
- FAI. 2016. *Fertilizer Statistics*. 2015-16. The Fertilizer Association of India, New Delhi.
- MOA. 2011. *Report of Expert Committee on Agricultural Statistics*. 2011. Directorate of Economics and Statistics, Ministry of Agriculture and Farmers welfare, Govt. of India.
- Singh R and Mangat N P S. 1996. *Elements of Survey Sampling*. Kluwer Academic Publishers, Springer-Verlag.
- Sud U C, Ahmad T, Gupta V K, Chandra H, Sahoo P M, Aditya K, Singh M and Biswas A. 2017. *Crop1&2: Measuring Crop Area and Yield under Pure Stand, Mixed and Continuous Cropping*. Working Paper No. 16. Global Strategy, FAO Publication, FAO, Rome. <http://gsars.org/wp-content/uploads/2017/01/WP-16.01.2017-Findings-from-the-Field-Tests-Conducted-in-Three-Countries.pdf>.
- Sukhatme P V, Sukhatme B V, Sukhatme S and Asok C. 1984. *Sampling Theory of Surveys with Applications*, 3<sup>rd</sup> Edition, Iowa State University Press, USA.