



## Disease management in horticulture crops through microbial interventions: An overview

POPY BORA<sup>1</sup> and L C BORA<sup>2</sup>

Assam Agricultural University, Jorhat, Assam 785 013, India

Received: 9 March 2020; Accepted: 9 June 2020

### ABSTRACT

Horticultural crops-based nutritional security is now a buzz word to Indian agriculture. Significant level of progress has been made towards the management of some pandemic pathogens affecting the production economics of horticultural crops. Some of these pathogens are : *Fusarium oxysporum*, *Ralstonia solanacearum*, *Pythium* spp., *Phytophthora* spp, *Xanthomonas* spp., *Colletotrichum* spp., *Penicillium* spp , *Alternaria* spp etc. Many soil-borne diseases as well as post harvest fungi are effectively addressed through a variety of bioagents belonging to predominant microbial genera represented by *Trichoderma* spp., *Bacillus* spp., *Gliocadium* spp. and fluorescent *Pseudomonads*. Our thumping success with different microbial bioformulations showed an effective control of both pests and diseases in major crops of northeast India comprising tea, banana, turmeric, ginger, in addition to bacterial wilt of *Bhoot jolokia*, tomato, hydroponic lettuce and *Rhizoctonia*- induced root rot management of chilli and patchouli. These area wide and crop wide results put forth a renewed claim in favour of developing a vibrant organic horticulture in years to come. All these issues are analysed highlighting the future use of rhizospheric and endophytic plant beneficial microbes as dominant communities.

**Key words:** Horticultural crops, Microbial interventions, Organic horticulture, Post-harvest diseases, Soil borne diseases

Horticultural crops have carved an indelible mark on the daily dietary requirement as an alternative source of nutritional security to the extent that the per capita consumption of cereal crops have reduced remarkably. Horticulture is ably recognized to have the potential to raise the farm income, provide livelihood security and earn foreign exchange, considering as much as 37% contribution of horticulture out of the total export of agricultural commodities ( NHB 2020). The importance of horticulture in day-to-day life is like a panacea, made unexpected strides in recent past, thereby, surpassed the food grain production by nearly 30 million tons, which in coming years is expected to make some bigger strides. India's horticulture production registered a 1% rise during 2018-19 to record 314.67 million tons from 25.87 million ha area (NHB 2020). Despite, such revealing success, quality production coupled with low productivity and profitability in several horticultural crops are still a cause

of serious concern. Of them, outbreak of diseases account for maximum damage to crops.

Most of the growers basically rely on chemicals for disease management as a quick remedy. But development of resistance against chemicals by pathogenic microbes coupled with environmental and public concern over hazardous chemicals have posed newer challenges to disease management of horticultural crops. The world trend in recent times is moving towards chemical residue free fruits, vegetables and tea plantations (Bora *et al.* 2016a). The production without chemical pesticides, technology based on natural resources has immense potential wherein exploring plant beneficial microorganisms is one such novel approach (Sharma *et al.*, 2012). Significant progress has been made worldwide in research for identification of efficient and aggressive biocontrol agents and exploited in disease management programmes due to their persistent nature in soil, phyllosphere, rhizosphere, high multiplication rate, easiness in handling and mass multiplication without any adverse effects on non-targeted organisms (Bora *et al.* 2016b). Voluminous literatures on rhizospheric antagonists are available which signifies its implications on plant health management in horticultural crops. Many microbes residing inside plants have (Popularly known as endophytes) caught researchers' attention in recent times

<sup>1</sup>Junior Scientist (Plant Pathology), ICAR-AICRP on Fruits (email:pбора.sonitpur10@gmail.com), <sup>2</sup>Professor and Head (email:boralohitc@gmail.com), Department of Plant Pathology, Assam Agricultural University, Jorhat, Assam 785 013.

due to their ability to suppress plant pathogens (Bora *et al.* 2019).

#### Microbial bioagents : The background

Both rhizospheric and endophytic microbes hold greater promise as sustainable option for plant health management in horticultural crops. Some of the plant beneficial microbes used as bioagent facilitate crop production through disease suppression as well as enhanced plant growth. *Trichoderma* spp. fluorescent *Pseudomonads*, *Bacillus* spp. etc. are established as plant growth promoter and commercially used as growth enhancer too (Bora *et al.* 2013). Horticultural crops including vegetables, fruits and medicinal plants harbor many endophytic microbes and many of them have been reported to possess antimicrobial properties. Compant *et al.* (2011) reported that bacterial genera *Pseudomonas* and *Bacillus* isolated from the interior of flowers, fruits and seeds of grapevine showed antagonistic property against pathogenic microbes.

Microbial biological control agents (BCA) provide protection to plants through different modes of action, viz. direct antagonism against the pathogens and priming host defense against pathogenic attack (Conrath *et al.* 2015). This has been resulted from a microbe-plant cross talk that involves an array of bioactive metabolites produced by the antagonist (Woo *et al.* 2006; Alfano *et al.* 2007). BCAs interact directly with the pathogen by hyperparasitism, antibiosis, production of antimicrobial secondary metabolites or indirectly through competition for nutrients and space (Bora *et al.* 2013, Singh *et al.* 2014).

Fluorescent *pseudomonads* such as *Pseudomonas fluorescens* and *P. putida* have been reported to exhibit biocontrol efficacy against a wide range of phytopathogens. They are aggressive colonizers of the rhizosphere of various crop plants and their antagonistic activity is attributed to antibiotics, siderophore production and ability to compete for nutrition or space besides producing an array of antifungal metabolites and lytic enzymes (chitinase and glucanase) that inhibit the mycelial growth of pathogenic fungi (Bora and Deka 2007, Bora *et al.* 2016a). *Bacillus* spp. also produce a wide range of antibiotics effective against many plant pathogens. These bacterial bioagents are effective against many bacterial and fungal pathogens (Bora and Bhagabati 1999, Bora *et al.* 2019). *Trichoderma* spp. (*T. viride*, *T. harzianum*, *T. koningi*, *T. hamatum*, *T. polysporum* and *T. longibrachiatum*) are reported successful against many soil borne diseases, employing an array of mechanisms, viz. hyperparasitism, competition for nutrients, antibiosis and induction of host defense mechanism in plants (Harman *et al.* 2004).

The plant beneficial microbes offer priming to the host plant through induced systemic resistance (ISR) regulated through salicylic acid independent mechanisms. However, PGPR triggering an SA-dependent type of ISR resembling pathogen induced SAR have also been reported. *P. fluorescens* strain WCS374 applied in the greenhouse grown radish induced ISR and suppressed Fusarium wilt

disease and also increased yield in radish (Pieters *et al.* 2014).

#### Microbial interventions and disease management

**Fruit crops:** Some of the destructive fruit diseases causing huge losses to the fruit industry worldwide include apple scab, mango malformation, guava wilt, fire blights, *Fusarium* wilt of banana, brown rot of stone fruits, crown gall, downy and powdery mildews etc. (Singh 1996, Bora and Deka 2014). Significant progress have been made for biological management of these diseases through BCAs like *Trichoderma* spp., *Gliocladium* spp., *Verticillium* spp., *Bacillus subtilis*, *Agrobacterium radiobacter*, *Pseudomonas* spp. (Rishbeth 1988). Successful management of crown gall pathogen *A. tumefaciens* through microbial strategy is a classic success story. The excellent control of bacterial crown gall disease of grapevine, pome and stone fruits have been achieved worldwide by using strain of K84 and its derivatives K 1026 of *Agrobacterium radibactor* (Bell *et al.* 1995, Thind 2017). These strains produce Agrocin 84 that kill *A. tumefaciens* and prevents the disease from further spread.

Microbial antagonists are more suited for control of soil borne pathogenic genera (*Pythium*, *Phytophthora*, *Rhizoctonia*, *Fusarium* and *Sclerotium*), where *Trichoderma* spp and *Gliocladium* spp., *P. fluorescens* are the most extensively used. *T. harzianum*, *T. viride*, *T. koningii*, *Pseudomonas putida* and *Mycothecium roroidum* provided an effective control of *Phytophthora* infestation in citrus (Menge 1993). Various fungal antagonists, *Ophisotoma* spp, *Chaetomium* sp, *Aureobasidium* sp and *Phoma* spp. inhibit the growth of the apple scab fungus (Outmet *et al.* 1997). While other antagonists such as *B. subtilis* and *T. koningii* have been also used for controlling apple canker (Lima *et al.* 1994).

Fusarium wilt in banana is a major concern for banana industry due to the devastating nature of the pathogen *F. oxysporum* f. sp. *cubense*. Management of Race 1 of the pathogen have been successfully addressed by several workers using strains of *Trichoderma* spp (Baruah *et al.* 2018). Anuratha *et al.* (1990) reported management of Moko disease, another disease of banana caused by bacterial pathogen *R. solanacearum* through application of *P. fluorescens* and *B. subtilis*. Citrus canker caused by *Xanthomonas citri* pv. *citri* is considered as another economically important disease as it affects the marketability of fruits. *T. viride* and *P. fluorescens* based formulation showed significant disease reduction (72%) with enhanced fruit yield when applied as soil application and foliar spray (Saikia 2018). *B. subtilis* and *P. fluorescens* significantly lowered the disease symptoms of citrus mal secco disease (*Phoma tracheiphila*) and the stem inoculated sour orange seedlings maintained higher population of these BCAs of the host plant (Wilhelm *et al.* 1998). Satisfactory progress has been made in addressing apple scab (*Venturia inaequalis*) management through microbe-based strategies. Burr *et al.* (1996) reported that isolate of *Pseudomonas syringae*

(isolate 508) prevented conidial germination and effectively suppressed apple scab pathogen to a level comparable to the fungicide captan. Efficacy of another antagonistic isolate, *Cladosporium cladosporioides* H39 was demonstrated by Kohl (2015) under field condition of organic apple orchard where 48-92% reduction in apple scab incidence was observed.

Besides fungal and bacterial bioagents, AM Fungi are also being explored against many fruit diseases. In an experiment papaya seedling treated with AM fungi *Glomus mosseae* and *G. deserticola* significantly suppressed the foot rot caused by *Pythium aphanidermatum* and also enhanced the seedling growth (Olawuyi *et al.* 2013). Combined treatment of melon seedlings by *Glomus intraradices* and *T. harzianum* provided a greater control against *Fusarium* wilt as compared to *T. viride* treated alone (Medina *et al.* 2016).

**Vegetable crops:** Ecofriendly management of vegetable disease through microbe based strategy has been quite successful and being practised in organic and IDM practices. Bacterial wilt (*R. solanacearum*), fungal wilt (*Fusarium* spp.), damping off and root rot (*Pythium* spp, *Phytophthora* spp, *Rhizoctonia solani*), anthracnose (*Colletotrichum* spp) etc. are few examples of diseases with successful management through different bioagents under field condition (Bora and Deka 2007). Bioagents have been explored against *R. solanacearum*, a pathogen enlisted among top 10 economically important bacterial pathogen with unusually wide host range that includes solanaceous vegetables. Bora *et al.* (2016c) reported formulation of *P. fluorescens* and *Trichoderma* spp. combination to reduce percent wilt incidence with increased yield attributes in brinjal and chilli. Few new and uncommon microbes, viz. *Chryso bacterium indologenes*, *C. luteola*, *Ralstonia pickettii*, *Streptomyces rochei* are reported to show inhibitory effect against the pathogen (Yuliar *et al.* 2015). However, their mode of action and risk assessment needs further investigation. Hydroponically grown vegetables, also

face pathogenic challenges that can be addressed through bioagents. Khan *et al.* (2018) reported that consortia of *T. viride* and *P. fluorescens* could significantly reduce root rot incited by *F. oxysporum* f.sp. *lectucae* with enhanced growth parameters in hydroponically grown lettuce. Utilisation of different microbial bioagents for successful control of various diseases of vegetable crops have further been summarised (Table 1).

**Spices, plantation and medicinal crops:** With the growing demand of organic spices, tea, coffee etc., biocontrol has become inevitable as disease management in these crops. *Trichoderma* spp. are widely used against many diseases of spice crops including ginger, turmeric, pepper, cardamom, cumin, turmeric etc (Chattopadhyay *et al.* 2018). *T. harzianum* ( Mass multiplied on wheat bran preparation in 1:1 ratio and applied at 50g per vine) was reported promising solution for the management of foot rot in black pepper (Jahagirdar and Siddaramaiah 2000). In tea, microbial bioformulations are predominantly used against diseases and insect pests. Some successful application of bioagents in medicinal, spices and other horticultural crops have further been enlisted (Table 2).

#### Post-harvest diseases

Exploiting the immune capacity of plants has been described as a plausible route to prevent post-harvest diseases, since plants have evolved the capacity to sensitize their immune system for a better expression of induced defence mechanism (Conrath *et al.* 2006). Induced post-harvest resistance through the use of safer chemicals from biological origin, and physical means have also been reported (Petriacq *et al.* 2018). Quite strikingly, however, more than a third of fruit and vegetables fails to reach the customer due to infections with pathogenic microbes (Oerke *et al.* 2004). Post-harvest losses can dramatically impact fruit production and quality (Romanazzi *et al.* 2016), resulting in losses of an average of 22.5% of the yield in developing countries. This phenomenon is known as priming of defence

Table 1 Biological control agents used against some popular diseases of vegetable crops

Vegetable	Disease and pathogen	BCA	Reference
Tomato	Fungal wilt ( <i>F. oxysporum</i> f.sp. <i>lycopersici</i> ) Bacterial wilt	<i>Trichoderma</i> spp. <i>Bacillus amyloliquefaciens</i>	Someshwar <i>et al.</i> (2013) Wei <i>et al.</i> (2011)
Tomato	Early blight ( <i>Alternaria solani</i> )	<i>T. virid</i>	Roopa <i>et al.</i> (2014)
Chilli	Fruit rot ( <i>C. dematium</i> ) Root rot ( <i>R. solani</i> )	<i>T. viride</i> <i>T. viride</i> and <i>B. subtiis</i>	Belge <i>et al.</i> (2009) Ahmed <i>et al.</i> (2003)
Brinjal	Verticillium wilt	<i>B. cereus</i>	Li <i>et al.</i> (2008)
Okra	Root rot ( <i>R. solani</i> )	<i>P. fluorescens</i>	Adhikari <i>et al.</i> (2013)
Cucumber	Damping off ( <i>Pythium ultimum</i> )	<i>B. subtilis</i> <i>P. fluorescens</i>	Khabbaz and Abbasi (2014)
French bean	Anthracnose ( <i>Colletotrichum lindemuthianum</i> )	<i>T. viride</i> , <i>T. harzianum</i> and <i>P. fluorescens</i>	Amin <i>et al.</i> (2003)
Cauliflower	Black rot ( <i>Xanthomonasa campestris</i> pv. <i>campestris</i> )	<i>Bacillus</i> spp	Luna <i>et al.</i> (2002)

Table 2 Successful bioagents against major diseases of spices, plantation and aromatic crops

Crop	Disease and pathogen	BCA	Reference
Ginger	Rhizome rot ( <i>Pythium aphanidermatum</i> )	<i>T. viride</i> and <i>P. fluorecens</i>	Sarma (2000)
	Bacterial wilt ( <i>R. solanacearu</i> )	<i>T. harzianum</i> and <i>P. fluorescens</i>	Bora <i>et al.</i> (2016 a)
Cardamom	Clump rot ( <i>P. vexans</i> , <i>R. solani</i> )	<i>Trichoderma</i> spp	Reddy (2014)
	Phytophthora rot ( <i>Phytophthora</i> spp.)	<i>Trichoderma</i> spp.	Thomas (2003)
Black pepper	Foot rot ( <i>Phytophthora capsici</i> )	<i>T. viride</i> , <i>T. harzianum</i> and <i>Glomus fasciculatum</i>	Dhanapal and Thomas (2009)
Fennel	Damping off and seedling rot ( <i>F. oxysporum</i> f. sp. <i>solani</i> , <i>R. solani</i> , <i>Pythium</i> spp)	<i>Trichoderma</i> sp. and <i>Bacillus subtilis</i>	Gebily (2015)
Vanilla	Root rot ( <i>Phytophthora meadii</i> )	<i>T. harzianum</i> and <i>P. fluorecens</i>	Reddy (2010)
	Wilt ( <i>F. oxysporum</i> f. sp. <i>vanillae</i> )	<i>T. harzianum</i> and <i>P. fluorescens</i>	Athul <i>et al.</i> (2012)
Tea	Grey blight	<i>T. viride</i> , <i>B. subtilis</i> and <i>P. fluorescens</i>	Deb <i>et al.</i> (2017)
	Phomopsis canker ( <i>Phomopsis theae</i> )	<i>Gliocladium virens</i> and <i>T. harzianum</i>	Ponmurugan and Baby (2007)

and is understood as an adaptive part of induced resistance (Mauch-Mani *et al.* 2017).

Devastating post-harvest fungal pathogenic genera such as *Penicillium* and *Botrytis* cause post-harvest fruit decay in considerable proportions. *P. digitatum* and *P. italicum* are responsible for green mould (or green rot) in citrus fruit and *P. expansum* cause blue mould which significantly affects orchard fruits (Kumar *et al.* 2018). The mildews caused by *Plasmopara viticola* and *Erysiphe necator*, and rots caused by *A. alternate*, also result in considerable post-harvest decay of grapes and tomatoes, respectively. The infections with *Clavibacter michiganensis* sub sp. *Sepedonicus* causing ring rot in potato and subsp. *michiganensis*, causing bacterial wilt and canker of tomato (Eichenlaub and Garteman 2011) causing quite a substantial economic losses are reported. The other diseases, such as bacterial spot in tomatoes and peppers caused by *Xanthomonas axonopodis*, also result in serious damage to the fruits (Basim *et al.* 2004) Yeast cultures are the most preferred option of biocontrol of post-harvest diseases against fungal pathogens (Wilson and Wisniewski 2003). The strains such as *Pichia membranefaciens* resulted in the up or down regulation of 25 proteins, which included antioxidants and PR proteins in peach fruit (Chan *et al.* 2007). Similarly, strains of *Aureobasidium pullulans* and *Cryptococcus laurentii* induced resistance in cherry tomatoes and peach, respectively, through the activation of host antioxidants metabolism. In addition, induced resistance activity triggered by *A. pollutants* has been linked to the accumulation of GLU, CHI and PAL enzymes (Ippolito *et al.* 2000). The bacteria, *Bacillus cereus* was proven effective to control anthracnose disease caused by *Colletotrichum acutatum* in loquat fruit through induced production of phenolic compounds and H<sub>2</sub>O<sub>2</sub> (Wange *et al.* 2014). Priming-based induced resistance has several benefits: i. it does not trigger a direct activation of defences therefore does not incur in major costs in growth and development; ii. it provides broad-spectrum resistance;

and iii. it is long-lasting and can reach the fruiting stage (Petriacq *et al.* 2018).

*Microbial bioformulations and disease management:* Our experiences of quality biopesticides with aggressive strains and their survival during storage are two key factors governing success of biological control in organic agriculture as well as IDM Programmes in conventional agriculture. Assam Agricultural University, the pioneer in biopesticide research in NE India greatly contributed to microbe mediated pest management through development of a few bioformulation, viz. Bioveer (*T. viride*), Biofor Pf2 (*P. fluorescens*+ *T. viride*), Biozin PTB (*P. fluorescens*+ *T. harzianum* + *B. brevis*), Biotime (*P. fluorescens*+ *T. viride*+ *Metarhizium anisoplae*), Biozium (*T. harzianum*), Biogreen-5 (*P. fluorescens*+ *T. viride*+ *B. subtilis* + *M. anisoplae* + *B. thuringiensis*). A bioformulation carrying antagonists as well as entomopathogens could be more sustainable and cost effective solution from the farmer's perspective as it would address both insect pests and disease problem simultaneously. Compatibility assay of different antagonistic microbes and entomopathogens such as *T. viride*, *P. fluorescens*, *Bacillus* spp and *M. anisoplae* have been carried out at Assam Agricultural University (Bora and Deka 2007, Deuri 2013, Khan *et al.*, 2016). Based on these studies we developed the consortia formulations and different field studies established that combination of different bioagents observed the higher magnitude of disease reduction as compared to single bioagent.

Microbial bioformulations developed at AAU, Jorhat have been found successful against some pests and diseases of economically important horticultural crops in Assam and NE India, viz. banana (Dutta 2015), tea (Deb *et al.* 2018), turmeric and ginger (Deuri 2013, Lalfakawma *et al.* 2014). Sarkar (2015) used aggressive strains of four different bioagents *Trichoderma viride*, *T. parareesei*, *M. anisoplae* and *B. thuringiensis* against *R. solanacearum*

causing bacterial wilt of *Bhoot jolokia* (*Capsicum chinense*) and recorded significant reduction of wilt incidence (83.87%) with highest yield (9.84 t/ha). Record of yield enhancement as well as suppression of bacterial wilt incidence in *Bhoot jolokia* (*Capsicum assamicum*) using consortia of *Trichoderma* along with some PGP microbes was also made by Bora *et al.* (2016a), bacterial wilt in tomato by Nath *et al.* (2016) and Bora *et al.* (2016b). Bora *et al.* (2016c) also recorded *Rhizoctonia* root management in chilli (*Capsicum frutescens*) using *Trichoderma*-based microbial consortia. Khan *et al.* (2016) used microbial consortia containing *T. viride* against wilt causing pathogen *Fusarium oxysporum* f. sp. *lactucae* in hydroponically grown lettuce (*Lactuca sativa*).

Gogoi *et al.* (2017) observed that soil application of Biofor-pf-2 (containing *T. viride* and *P. fluorescens*) @ 100 g/pot showed least root rot incidence (44.40%) of patchouli caused by *Fusarium solani* as compared to inoculated control (100%). Seed treatment of vegetable seeds with nano-bio formulation of *Trichoderma asperellum* and chitosan NP suppressed the growth of *F. oxysporum*, *Sclerotium rolfisii* and *Rhizoctonia solani* as compared to recommended chemicals (of patchouli caused by *Fusarium solani*).

#### Future line of research

Microbes from different genera of rhiospheric and endophytes have shown some precision in disease management. However, root colonization efficiency of effective microbes should also be further studied to increase their effectiveness as bioinoculants via. consortia inoculants as priority research. The major thrust area in years to come would be: concentration on development of novel formulation, understanding the impact of environmental factors on population dynamics of bio-control agents, mass production of bio-control microorganisms and use of biotechnology and nano-technology in improvement of bio-control mechanism and strategies and development of a multi-utility bioformulation by integrating micronutrient supplements. Micronutrients can enhance disease suppression indirectly by stimulating indigenous populations of antagonistic microorganisms (Huber 2007) and directly by activating systemic acquired resistance in host plant systems (Dordas 2007). Exploration of antagonists and bio-fertiliser agents as consortia can significantly reduce crop diseases in field along with enhanced growth and yield (Srivastava *et al.* 2010).

The researches around the world are showing concern about the probable risks of biopesticides also, correlating the facts that biopesticides showing toxic property against target pests may also transfuse these properties to plant physiology and biochemistry. Therefore, toxicological studies of microbial biopesticides are now essential through accumulation of biological characteristics and chemistry data of the microbes used for preparation of biopesticides.

Role of AMF in providing an additional resilience to rhizosphere's ability of carbon accreditation within rhizosphere and associated development of plant's antioxidant profile

as a defense mechanism (Wu *et al.* 2013) should divert the research studying strong mycorrhizal dependency of fruit crops. Rhizosphere specific AMF-based microbial consortium would add a new dimension in providing newer options for raising the productivity potential of fruit crops as well as enhancing plant immune system to add the required physiological preparedness to the plant fight against pathogens. Researches are using different approaches for screening rhizobacteria to select effective PGPR including promotion of root/shoot growth under genotobiotic condition, *in-vitro* production of plant growth regulators/biologically active substances and assessing of deaminase activity of the rhizobacteria (Zahir *et al.* 2003). Fungal biofertilizer agent arbuscular mycorrhizal applied along with *T. harzianum* based formulation could lead to a significant reduction of disease along with enhanced growth and yield of crops like tomato in field (Srivastava *et al.* 2010).

A system-based approach emphasizes the incorporation of different practices involving: (i). Prevention of introduction and spread of pathogens; (ii.) Reduction of inoculum to the point, manageable under natural system; (iii.) Promotion of disease suppressive soil microbial communities and (iv.) Use of an integrative approach to interfere with life cycle of pathogens and minimising the disruptive actions of pesticides/biopesticides. These approaches must go hand-in-hand while addressing various diseases in horticultural crops in order to maintain sustainable production of zero risk horticultural crops in the days to come.

#### REFERENCES

- Adhikari A, Dutta, S and Nandi S. 2013. Antagonistic potentiality of native rhizobacterial isolates against root rot disease of okra incited by *Rhizoctonia solani*. *African Journal of Agricultural Research* **8**(4):405–412
- Ahmed A, Ezziyyani M and Candela, M E. 2003. Effect of chitin on biological control activity of *Bacillus* spp. and *Trichoderma harzianum* against root rot disease in pepper (*Capsicum annum*) plants. *European Journal of Plant Pathology* **106** (6): 633-637
- Alfano G, Ivey M LL, Cakir C, Bos J I B, Miller S A, Madden L V, Kamoun S and Hoitink H A J. 2007. Systemic modulation of gene expression in tomato by *Trichoderma hamatum*. *Phytopathology* **97**: 429-437.
- Amin M, Ayalew A, Dechassa N and Negeri M. 2013. Effect of integrated management of anthracnose (*Colletotrichum lindemuthianum*) on plant and seed health of common bean in Hararge highlands, Ethiopia. *Journal of Science and Sustainable Development* **1**:1-19
- Anuratha C S and Gnanamanickam S S. 1990. Biological control of bacterial wilt caused by *Pseudomonas solanacearum* in India with antagonistic bacteria. *Plant & Soil* **124**:109–116.
- Athul S, Aju R and Asok A J. 2012. Biocontrol of fusarium wilt of vanilla (*Vanilla planifolia*) using combined inoculation of *Trichoderma* sp. and *Pseudomonas* sp. *International Journal of Pharmacology and Biological Sciences* **3**(3B): 706 – 716.
- Baruah N, Bhattacharyya A, Thangavellu R and Puzari K C. 2018. *In vitro* screening of native banana rhizospheric microbes and endophytes of Assam against *Fusarium oxysporum* f. sp. *cubense*. *International Journal of Current Microbiology Applied*

*Sciences* 7: 1575-1583

- Basim H, Basim E, Jones J B, Minsavage G V and Dickstein E R. 2004. Bacterial spot of tomato and pepper caused by *Xanthomonas axonopodis* pv. *vesicatoria* in the Western Mediterranean Region of Turkey. *Plant Disease* **88** : 85-92
- Bell C, Dickie G, Harvey W and Chan J. 1995. Endophytic bacteria in grapevine. *Canadian Journal of Microbiology*. 41: 46–53
- Belgee B A and Padghan P R. 2009. Efficacy of bioagents and fungicides against *Colletotrichum dematium* - the incitant of fruit rot of chilli. *Environment and Ecology* **29**:1391-1394
- Bora L C and Bhagabati K N. 1999. Quantitative aspects of assessing virulence of *Xanthomonas campestris* pv. *citri* (Hasse) Dye, the incitant of citrus canker in Assam. *Intl. Symp. citriculture*, National Research. Centre for Citrus, Nagpur, Maharashtra, p 82.
- Bora L C and Bora Popy. 2008. Vermicompost based bioformulation for management of bacterial wilt of tomato in poly house. *Journal of Mycology and Plant Pathology* **38**(3): 527-530.
- Bora L C and Deka S N. 2007. Wilt disease suppression and yield enhancement in tomato (*Lycopersicon esculentum*) by application of *Pseudomonas fluorescens* based bio pesticide (Biofor-Pf) in Assam. *Indian Journal of Agricultural Sciences* **77**: 490-494.
- Bora L C, Katak L, Talukdar K and Khan P. 2016c. Yield enhancement and bacterial wilt suppression in Bhut Jolokia using consortia of microbial antagonists. *Biopesticides International* **12**(2): 165-171.
- Bora L C, Katak L, Talukdar K, Nath B C and Sarkar R. 2015. Molecular characterizations of Microbial antagonists and development of bioformulations for management of bacterial wilt of Naga chilli (*Capsicum chinens*) in Assam. *Journal of Experimental Biology Agricultural Science* **3**(2): 109-122.
- Bora Popy and Bora L C. 2016a. *Hand Book of Organic Farming*. Sabda Publishers, Guwahati, Assam.
- Bora Popy, Bora L C and Begum M. 2013. Eco-friendly management of soil borne diseases in brinjal through application of antagonistic microbial population. *Journal of Biological Control* **27**: 29–34.
- Bora Popy, Bora L C and Deka P C. 2016c. Efficacy of substrate based bioformulation of microbial antagonists in the management of bacterial disease of some solanaceous vegetables in Assam. *Journal of Biological Control* **30**: 49-54.
- Bora Popy and Deka P C. 2014. Biointensive approaches for disease management of *rabi* vegetables: Agriculture for sustaining livelihood. Bhattacharya H, Neog M and Saud B (Eds). Assam Agricultural University Press, p 123-126.
- Bora Popy, Deka P C and Sarmah A K. 2016b. Efficacy of *Pseudomonas fluorescens* and *Trichoderma viride* based bioformulation for management of bacterial wilt disease of ginger. *International Journal of Plant Science*. 11 (2): 34-39.
- Bora Popy, Saikia K, Hazarika H and Gavas R. 2019. Exploring potential of bacterial endophytes in disease management of horticultural crops. *Current Horticulture* **7**: 32-37.
- Boruah S. 2017. Development of next-gen nano-bioformulation of *Trichoderma* for seed treatment of vegetable crops. Ph D (Agri.) thesis, Assam Agric. Univ., Jorhat, Assam.
- Burr T J, Matteson M C, Smith C A, Corral-Garcia M R and Huang T C. 1996. Effectiveness of bacteria and yeasts from apple orchards as biological control agents of apple scab. *Biological Control* **6**:151-157.
- Chan Z, Qin G, Xu X, Li B and Tian S. 2007. Proteome approach to characterize proteins induced by antagonist yeast and salicylic acid in peach fruit. *Journal of Proteome Research*. **6**: 1677–1688.
- Chattopadhyay A, Purohit J, Patel D S and Tatarwal M L. 2018. Management of diseases of spice crops through microbes. (In) *Biological Control of Crop Diseases: Recent Advances & Perspective*. Singh D, Chakraborty B N, Pandey R N and Sharma P (Eds). Today & Tomorrow's Printers and Publishers, New Delhi, India, pp 249-270.
- Compant S, Mitter B, Colli-Mull J G, Gangl H and Sessitsch A. 2011. Endophytes of grapevine flowers, berries, and seeds: identification of cultivable bacteria, comparison with other plant parts, and visualization of niches of colonization. *Microbial Ecology* **62**: 188–197.
- Conrath W, Beckers G J M, Langenbach C J G, and Jaskiewicz M R. 2015. Priming for enhanced defense. *Annual Review of Phytopathology* **53**: 97–119.
- Deb R. 2017. PGPM formulation mediated biochemical changes and defense response of tea against major diseases and pests. M Sc (Agri.) thesis, Assam Agricultural University, Jorhat, Assam.
- Deuri D. 2013. Bio-intensive approach for management of bacterial wilt of ginger (*Zinziber officinale*). M Sc (Agri.) thesis, Assam Agricultural University, Jorhat, Assam.
- Dhanapal K and Thomas J. 2009. Role of biocides in spices disease management. (In) *Role of Biocontrol Agents for Disease Management in Sustainable Agriculture*, pp. 27–33. Ponnuragan P and Deepa MA (Eds). Research India Publications, Delhi, India.
- Dordas C. 2007. Role of nutrients in controlling disease in sustainable agriculture: A review. *Sustainable Development* **28**: 33-46.
- Dutta R. 2015. Eco-friendly management of *Alternaria* leaf blight of Mandukaparni (*Centella asiatica* L.). MSc (Agri.) thesis, Assam Agricultural University, Jorhat, Assam.
- Eichenlaub R, Gartemann K H. 2011. The clavibacter michiganensis species: Molecular investigation of gram-positive bacterial plant pathogens. *Annual Review of Phytopathology* **49**: 455-465.
- Gebily D. 2015. Studies on root rot disease of fennel under organic farming regulations. M Sc thesis, Faculty of Agriculture. Fayoum University, Fayoum, Egypt, p 121.
- Gogoi M, Sarmah D K, Bora L C and Barua I C. 2017. Management of root rot disease of patchouli caused *Fusarium solani* (Mart) Sacc. Through fungicide, bioagent and oilcake in field. *International Journal Current Microbiology and Applied Science* **6**: 3385-3397.
- Harman G E, Howell C R, Viterbo A, Chet I and Lorito M. 2004. *Trichoderma* species opportunistic, avirulent plant symbionts. *Nature Reviews Microbiology* **2**: 43-56.
- Huber D M and S Hanekklans. 2007. Managing nutrition to control plant disease. *Landbanforsch Volkenrode* **57**: 313-322.
- Ippolito A, El Ghaouth A, Wilson C L and Wisniewski M. 2000. Control of postharvest decay of apple fruit by *Aureobasidium pullulans* and induction of defense responses. *Postharvest Biology and Technology* **19**: 265–272.
- Jahagirdar S and Siddaramaiah A L. 2000. Association of pathogens in foot rot complex of black pepper in Karnataka. *Karnataka Journal of Agricultural Science* **13**: 757-760.
- Khabbaz S E and Abbasi P A. 2014. Isolation, characterization, and formulation of antagonistic bacteria for the management of seedlings damping-off and root rot disease of cucumber. *Canadian Journal of Microbiology* **60**: 25–33.
- Khan P, Borah P K, Bora Popy and Bora L C. 2016. *Fusarium oxysporum* f. sp. *lactucae* (FoL) a potential threat to

- hydroponically grown lettuce (*Lactuca sativa*) and its management using microbial consortia. NE zonal meeting. Indian Phytopathological Society. p 35.
- Kohl J. 2015. Towards an integrated use of biocontrol by *Cladosporium cladospoides* in apple scab management. *Plant Disease* **99** : 534-539.
- Kumar D, Tannous J, Sionov E, Keller N and Prusky D. 2018. Apple intrinsic factors modulating the global regulator, Lae a, the patulin gene cluster and patulin accumulation during fruit colonization by *Penicillium expansum*. *Frontiers in Plant Science* **9**:1094.
- Lalfakawma C, Nath B C, Bora L C, Srivastava S and Singh J P. 2014. Integrated disease management of *Zingiber officinale* Rosc. rhizome rot. *Bioscan* **9**: 265-269.
- Li J G, Jiang, J Q, Xu L P, Sun P and Gu J H. 2008. Characterization of chitinase secreted by *Bacillus cereus* strain CH2 and evaluation of its efficacy against *Verticillium* wilt of eggplant. *Applied Soil Ecology* **53**:931-944.
- Lima G, Ippolito A and Nigro F. 1994. Attempting at biological control of citrus mal secco (*Phoma tracheiphila*) with endophytic bacteria. *Difesa delle piante* **17**: 43-49.
- Luna C L, Mariano R L R and Souto-Maior A M. 2002. Production of a biocontrol agent for Crucifers black rot disease. *Brazilian Journal of Chemical Engineering* **19**: 133-140.
- Mauch-Mani B, Baccelli I, Luna E and Flors V. (2017). Defense priming: An adaptive part of induced resistance. *Annual Review of Plant Biology* **68**: 485-512.
- Medina A M, Fernandez I, Lok G B, Pozo M J, Pieterse C M J and Wees S C M Van 2016. Shifting from priming of salicylic acid- to jasmonic acid-regulated defences by *Trichoderma* protects tomato against the rootknot nematode *Meloidogyne incognita*. *New Phytologist* **213**: 1363-137.
- Menge J A. 1993. Improved root rot control with mulches. *California Citrograph* **79**: 16-17.
- Nath B C, Bora L C, Katak L, Talukdar K, Sharma P, J Dutta and Khan P. 2016. Plant growth promoting microbes, their compatibility analysis and utility in biointensive management of bacterial wilt of tomato. *International Journal of Current Microbiology and Applied Science* **5**: 1007-1016.
- National Horticulture Board. 2020. Area and Production of Horticultural crops for 2018-19(1<sup>st</sup> Advance Estimate), www.nhb.gov.in
- Oerke E C and Dehne H W. 2004. Safeguarding production-losses in major crops and the role of crop protection. *Crop Protection* **23**: 275-285.
- Olwuyi O J, Odebode A C, Oyewole I U, Akanmu A O and Afolabi O. 2013. Effect of arbuscular mycorrhizal fungi on *Pythium aphanidermatum* causing foot rot disease on pawpaw (*Carica Papaya* L.) seedlings. *Archives of Phytopathology and Plant Protection* **47**: 1-10
- Outmet A, Carisse, O and Neumann P. 1997. Evaluation of fungal isolates for inhibition of vegetative growth of *Venturia inaequalis*. *Canadian Journal of Botany* **7**: 626-631.
- Petricaq P, Lopez A and Luna E. 2018. Fruit decay to diseases: Can induced resistance and priming help?. *Plants* **77**:doi:10.3390/plants7040077.
- Pieters C, Zamioudis C, Berendsen R and Wees S V. 2014. Induced systemic resistance by beneficial microbes. *Annual Review of Phytopathology* **52**:347-75.
- Ponmurugan P and Baby U I. 2007. Evaluation of fungicides and biocontrol agents against *Phomopsis* canker of tea under field conditions. *Australasian Plant Pathologists* **36**(1): 68-72.
- Reddy P P. 2014. *Biointensive Integrated Pest Management in Horticultural Ecosystems*, Springer, India.
- Reddy P P. 2010. Spice crops. (In) *Fungal Diseases and their Management in Horticultural Crops* pp 312-347, (Ed. Reddy PP), Scientific Publishers, Jodhpur, India.
- Rishbeth J. 1988. Biological control of air borne pathogens. (In) *Biological Control of Pests. Pathogens and Weeds: Development and Prospects. Proc. Royal Soc. Discussion Meeting, London (Eds. Wood, RKS and MJ Way)*. London Royal Society.
- Romanazzi G, Sanzani S M y, Tian S, Gutierrez Martinez and P, Alkan N. 2016. Induced resistance to control postharvest decay of fruit and vegetables. *Postharvest Boiology and Technology* **122**:82-94.
- Roopa R S and Yadahalli K B. 2014. Evaluation of natural plant extracts, antagonists and fungicide against early blight of tomato caused by *A. solani* in vitro. *Bioscan* **9**: 1309-1312.
- Saikia, S. 2018. Biointensive approach for citrus canker management. M Sc thesis, Department of Plant Pathology, Assam Agriculture University, Jorhat, Assam.
- Sarma Y R. 2000. Integrated management of rhizome rot of ginger. Final report of ICAR, AP Cess Fund Scheme, Indian Institute of Spice Research, Calicut, Kerala, India, pp 1-110.
- Sharma P, Patel A N, Saini M K and Deep S. 2012. Field demonstration of *Trichoderma harzianum* as a plant growth promoter in wheat (*Triticum aestivum* L). *Journal of Agricultural Science* **4**(8): 65-73.
- Singh S J. 1996. *Advances in Disease of Fruit Crops in India*. Kalyani Publisher, Ludhiana, India.
- Singh A, Sarma B K, Singh H B and Upadhyay R S. 2014. *Trichoderma*: A silent worker of plant rhizosphere. (In) *Biotechnology and Biology of Trichoderma*, pp 543-549. [Eds. Gupta et al.]. Elsevier.
- Someshwa B, Bambawale O M, Tripathi A K, Ahmad I and Srivastava R C. 2013. Biological management of fusarial wilt of tomato by *Trichoderma* spp. in Andamans. *Indian Journal of Horticulture* **70**: 397-403.
- Srivastava A K, Malhotra, S K and N K Krishna Kumar. 2015. Exploiting nutrient-microbe synergy in unlocking productivity potential of perennial fruits: A review. *Indian Journal of Agricultural Sciences* **85**: 459-81.
- Srivastava R, Khalid A, Singh U S and Sharma A K. 2010. Evaluation of arbuscular mycorrhizal fungus, *Fluorescent Pseudomonas* and *Trichoderma harzianum* formulation against *Fusarium oxysporum* f.sp. *lycopersici* for the management of tomato wilt. *Biological Control* **53**: 24-31.
- Thind S K. 2017. Principles of disease management in fruit crops. *International Clinical Pathology Journal* **5**:123-137.
- Thomas J. 2003. Biocontrol agents for diseases of spice crops. *Spice India* **16**: 9-12.
- Wang X, Wang L, Wang J, Jin P, Liu H and Zheng Y. 2014. *Bacillus cereus* AR156-induced resistance to *Colletotrichum acutatum* is associated with priming of defence responses in loquat fruit. *PLoS ONE* **9**: e112494.
- Wei Z, Yang X, Yin S and Xu Y. 2011. Efficacy of *Bacillus*-fortified organic fertiliser in controlling bacterial wilt of tomato in the field. **48**: 152-159.
- Wilhelm E, Arthofer W and Schafleitner R. 1998. *Bacillus subtilis* an endophyte of chestnut (*Castanea sativa*) as antagonist against chestnut blight (*Cryphonectria parasitica*). *Plant Cell Tissue Organ Culture* **12**:331-337.
- Wilson C and Wisniewski M. 2003. Biological control of postharvest diseases of fruits and vegetables: An emerging

- technology. *Annual Review of Phytopathology* **27**:425–441.
- Woo S L, Scala F and Ruocco M. 2006. The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology* **96**:181-185.
- Wu Qiang-Sheng, Srivastava A K and Ying-Ning Zou. 2013. AMF-induced tolerance to drought stress in citrus. A review. *Scientia Horticulturae* **164**: 77–87.
- Yuliar , Nion Y A and Toyota K. 2015. Recent trends in control methods for bacterial wilt disease cauded by *Ralstonia solanacearum*. *Microbes and Environments* **30**: 1-11.
- Zahir A, Zahir Mohmmad, Arshad, William T, and Frankenberger Jr. 2003. Plant growth promoting rhizobacteria : Applications and perspectives in agriculture. *Advances in Agronomy* **31**: 97–166.