Effect of conservation agriculture on soil organic carbon dynamics and mineral nitrogen under different fertilizer management practices in maize (*Zea mays*)-wheat (*Triticum aestivum*) cropping system

PRITI TIGGA¹, MAHESH C MEENA^{2*}, ABIR DEY³, B S DWIVEDI⁴, S P DATTA⁵, H S JAT⁶ and M L JAT⁷

ICAR-Indian Agricultural Research Institute, New Delhi 110 012, India

Received: 11 October 2019.; Accepted: 27 November 2019

ABSTRACT

Soil organic carbon (SOC) is the center of all physical, chemical and biological properties of soils, its maintenance and buildup in soils is necessary to sustain the intensive cereal-based cropping system of Indo Gangetic plains. Intensive tillage, residue removal and indiscriminate use of fertilizers led to a continuous deterioration of soil health in Indo Gangetic parts of the country. Under this context, a field experiment on conservation agriculture (CA) was conducted at Taroari, Karnal, comprising four combinations of tillage and residue management [i.e. conventional tillage (CT) with complete residue removal (CT-RR), CT with 20% residue incorporation of wheat, 50% residue incorporation of maize and incorporation of green gram residue (CT+RI+GI), permanent beds (PB) with 20% residue retention of wheat and 50% residue retention of maize (PB+RR), and PB with 20% residue retention of wheat, 50% residue retention of maize and retention of green gram residue (PB+RR+GR)] and three nutrient management options [viz. farmers fertilization practice (FFP), recommended fertilizer dose (RDF) and site-specific nutrient management (SSNM)]. The results showed that, there was an increase in SOC mainly the active SOC pools, permanganate oxidizable-C (1.70 g/kg), hot water extractable-C (0.32 g/kg) and soil microbial biomass C (310 mg/kg) under CA-based treatment (PB+RR+GR) compared to CT (0.58, 0.23 g/kg, 183 mg/kg, respectively). Also, the mineral N was invariably greater under residue treatments. Therefore, crops residue retention as well as balanced fertilization (RDF and SSNM) under CA helped in improving SOC; mineral N and soil aggregation stability which can lead to increased sustainability under cereal-based intensive cropping systems.

Key words: Conservation agriculture, Mineral nitrogen, Nutrient management, Soil organic carbon

Managing soil organic carbon (SOC) is a pre-requisite for restoring soil health and sustaining high productivity (Meena *et al.* 2012). Soil physical, chemical and biological properties are intrinsically linked to SOC (Thomas *et al.* 2018). The SOC levels of most of the Indian soils are extremely low, *i.e.* 0.3 to 0.4% resulting in poor soil health and low crop yields. Continuous intensive tillage, imbalanced fertilization and continuous cropping of ricewheat system is practiced especially in the Indo-Gangetic plains (IGP) caused further degradation of SOC (Das *et al.* 2017, Dey *et al.* 2018).

¹Ph D Scholar (prititigga31@gmail.com), ²*Senior Scientist and corresponding author (mcmeena@gmail.com), ³Scientist (abirdey21@gmail.com), ⁴Head (bsdwivedi@yahoo.com), ⁵Professor (profssac2017@gmail.com), Division of Soil Science and Agriculture Chemistry; ICAR-Indian Agricultural Research Institute, New Delhi; ⁶Senior Scientist (hsjat_agron@yahoo.com), CIMMYT-India, Karnal; ⁷Senior Cropping Systems Agronomist (M.Jat@cgiar.org), CIMMYT-India, New Delhi.

Balanced fertilization as an agricultural management strategy is being used to promote soil C storage (Mandal *et al.* 2013) which could directly or indirectly increase the SOC inputs and thereby change the availability of nutrients and soil turnover. In fact, Vanlauwe *et al.* (2014) proposed nutrient management as the fourth principle of CA and argued that appropriate use of fertilizer is required to define CA to enhance both crop productivity and produce sufficient crop residues to ensure soil cover.

Soil organic C consists of multiple compounds, from simple to more complex molecules having varying lability (Mandal *et al.* 2013). Since changes induced by soil management practices are often difficult to detect by total SOC measurement, assessing rapidly changing SOC pools, *i.e.* labile pool, might be more informative to assess soil quality (Awale *et al.* 2017, Nath *et al.* 2017). Therefore, different methods used to assess mainly the labile fractions of SOC such as potassium permanganate oxidizable C (KMnO₄-C), microbial biomass carbon (MBC), and hot water-extractable C (HWEC) have recently received more attention due to their sensitivity to management practices.

These labile fractions change rapidly with time and can provide an early assessment of SOC changes induced by management practices such as tillage, cropping system, and N fertilization (Chen et al. 2009). Bhattacharyya et al. (2019) advocated that short-term CA had positive effects on soil aggregation and nitrogen accumulation, which further had a role in increasing sustainability of cereal based cropping systems. This underlines the importance of such studies on CA and nutrient management under different cropping system and thereby, establishing specific fertilizer management for its successful adoption by farmers. Therefore, this study was conducted based on the hypothesis that retention of crop residues under CA would reflect early changes in various SOC pools and mineral N, with objectives to evaluate the impact of different tillage and residue management practice on SOC and N pools; and to outline the best fertilizer management option under CA.

MATERIALS AND METHODS

Study site characteristics

A field experiment on conservation agriculture (CA) was initiated in the year 2012 at a farmer's field located at village Taraori, Karnal to encourage the farmer's participation under Participatory Platform for Climate Smart Agriculture under the aegis of CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) of CIMMYT. The soil type is Typic Ustocrept. The climate of study area is semi-arid subtropical, characterized by very hot summers and cool winters. The average annual rainfall recorded is 700 mm of which 70-80% is received during July to September. The collected initial soil samples from uncultivated area (two depth, i.e. 0-5 and 5-15 cm soil depth) were analysed for their characteristics. The soil was non-saline (EC 0.39 dS/m), clay loam with mildly alkaline reaction (pH 8.02), and medium in Walkley-Black carbon (0.57%). The experimental soil was categorized low in KMnO₄-oxidizable N (157 kg/ ha), medium in 0.5M NaHCO₃-extractable P (17.6 kg/ha) and also medium in 1N NH₄OA_C-extractable K (235 kg/ ha). The initial soil was fairly permeable with a bulk density of 1.52 Mg/m^3 .

Treatments and experimental design

The field experiment was established in split-plot design under maize (*Zea mays* L.) - wheat (*Triticum aestivum* L.) cropping system with 12 treatments combinations. The main plot comprised of 4 treatment combinations of tillage, residues and legume, *viz.* (i) conventional tillage with complete residue removal (CT-RR), (ii) conventional tillage with 20% residue incorporation of wheat, 50% residue incorporation of maize and incorporation of green gram residue (CT+RI+GI), (iii) permanent beds with 20% residue retention of wheat and 50% residue retention of maize (PB+RR), and (iv) permanent beds with 20% residue retention of greengram residue (PB+RR+GR). The actual amount of crop residue retained/incorporated is given in

Table 1. The sub-plot consists of 3 treatments of different nutrient management options, *viz.* (i) Farmer's fertilizer practice (FFP), (ii) Recommended dose of fertilizers (RDF), and (iii) Site-specific nutrient management ((SSNM). Farmer's fertilizer practice (FFP) was based on a local survey conducted in that area for fertilizer application rates practiced by farmers in maize and wheat crops. Recommended dose of fertilizers (RDF) were fertilizers doses for maize and wheat as advocated by CCS Haryana Agricultural University, Hisar. The SSNM options were calculated based on a computer-based decision support system 'Nutrient Expert'. The amount of fertilizers applied to the crops as per treatments is given in Table 2.

Soil sampling and analysis

After completion of three cropping cycles, soil samples were collected from two depths, *i.e.* 0-5 cm and 5-15 cm of treated plots using core sampler at physiological maturity of wheat. Carbon fractions, *i.e.* (i) Very labile SOC ($C_{\rm VL}$): SOC oxidizable with 12 N H₂SO₄; (ii) Labile SOC ($C_{\rm L}$): Difference between SOC oxidizable with 18 N H₂SO₄ and 12 N H₂SO₄; (iii) Less labile SOC ($C_{\rm LL}$): Difference between SOC oxidizable with 24 N H₂SO₄ and 18 N H₂SO₄; and (iv) Non-labile SOC ($C_{\rm NL}$): Difference between TOC and 24 N H₂SO₄ were determined using Modified Walkley-Black method (Chan *et al.* 2001).

The permanganate oxidizable C (KMnO₄-C) was determined by oxidizing the samples with 33 mM KMnO₄ (Tirol-Padre and Ladha 2004), hot water extractable carbon (HWE-C) by Ghani $et\ al.$ (2003) and soil microbial biomass carbon (MBC) by fumigation-extraction method described (Jenkinson and Ladd 1981). Microbial extraction efficiency (K_{EC} of 0.25) was used for MBC determination.

Table 1 Treatment-wise amount of crop residues (t/ha) retained/ incorporated in the soil

Treatment*	FFP	RDF	SSNM
CT-RR	0	0	0
CT+RI+GI	4.14	4.31	4.63
PB+RR	3.39	3.56	3.74
PB+RR+GR	4.52	4.62	4.85

*CT: Conventional tillage; PB: Permanent beds; -RR: Residue retention; +RR: Residue removal; GI: Greengram residue incorporation; GR: Greengram residue retention

Table 2 Nutrient applied (kg/ha) in maize and wheat under different nutrient management practices

Treatment@		Maize			Wheat		
	N	P _O	K _O	N	PO	KO	
FFP	172.5	22.5	ő	230	22.5	ő	
RDF	187.8	24.5	62.5	177-184	23.5	60	
SSNM (NE)	172.5	18.0	60-75	150-164	15.3	81	

@ FFP: Farmer's fertilizer practice; RDF: Recommended dose of fertilizer; SSNM: Site-specific nutrient management

For mineral-N ($NH_4 + NO_3$) determination, moist soil sample was extracted with 2M KCl solution (soil: solution:: 1:5) (Rowell 1994). Aggregate stability, viz. mean weight diameter (MWD) and geometric mean diameter (GMD) were calculated as the indices of aggregate stability (van Bavel 1950; Kemper and Roseneau 1986) using following formula:

$$\begin{array}{c} \text{MWD=}\Sigma\text{xiwi}\\ \text{and, GMD =}\text{exp} \; [\Sigma \; (\text{wi log xi})/(\; \Sigma \; \text{wi}) \end{array}$$

where, wi is the weight of each aggregate class to the total soil weight; and xi the mean diameter (mm) of the class. The data recorded for different parameters were analyzed with the help of analysis of variance (ANOVA) technique (Gomez and Gomez 1984) for split-plot design using SAS 9.1 software. The least significant difference (LSD) test was used to interpret the effect of treatments at 5% level of significance (p=0.05).

RESULTS AND DISCUSSION

Soil organic C pools

The active SOC (C_{VL} and C_{L}) and recalcitrant pool (C_{LL} and C_{NL}), both declined in lower soil depth (5-15 cm) compared to surface 0-5 cm. These fractions followed the order as: $C_{VL} > C_{NL} > C_{LL} > C_{L}$. Tillage and residue management significantly influenced different SOC fractions in the surface soil depth (0-5cm), while in 5-15 cm depth no significant variation within treatments was observed, except $C_{\scriptscriptstyle T}$ fraction. The highest $C_{\scriptscriptstyle VL}$ was recorded under PB+RR+GR (2.84 g/kg) it was 25.7% higher than that under CT-RR and followed a trend as: PB+RR+GR>CT+RI+GI>PB+RR>CT-RR. Among the nutrient management options, the C_{VL} content was highest under RDF (2.81 g/kg). Unlike C_{VL}, C_L was not significantly affected by tillage, residue management or nutrient management options in surface depth. However, in 5-15 cm soil depth, the individual impact of tillage practices gave significantly higher C_I (0.66 g/kg) under

PB+RR+GR. The PB+RR showed significantly higher $C_{\rm LL}$ (1.92 g/kg) compared with other tillage practices in surface soil depth. The results indicated that both PB as well as CT treatments was at par with each other. The effect of nutrient management options was significant with relatively higher $C_{\rm LL}$ under SSNM (1.87 g/kg). The $C_{\rm NL}$ content under different tillage practices varied from 2.34 to 2.53 g/kg and 1.15 to 2.97 g/kg in 0-5 and 5-15 cm depths, respectively. The effect of nutrient management options was significant only for surface soil with comparatively higher $C_{\rm NL}$ (2.52 g/kg) content under SSNM, showing its superiority over FFP.

On an average, the contribution of active SOC was higher in the surface soil, whereas there was a shift towards recalcitrant pools in the lower depth. Labile organic matter in soil mainly originates from the decomposition of plant and faunal biomass, root exudates, and deceased microbial biomass (Bolan *et al.* 2011). Addition of organic matter and reduced tillage in the surface soil layer may likely to increase labile organic carbon in the soils (Cooper *et al.* 2016). Other researcher reported greater lability of C under CA compared with CT (Bhattacharya *et al.* 2012; Dey *et al.* 2018). In addition, these practices have the potential to enhance C and N cycling, as well as soil aggregation, which is one of the primary mechanisms through which organic carbon is sequestered in soil (Panettieri *et al.* 2015; Mandal *et al.* 2019).

Potassium permanganate oxidizable, hot-water extractable and soil microbial biomass carbon

The influence of tillage and residue management practices on permanganate oxidizable C (KMnO $_4$ -C) showed significant difference in both soil depths (Table 4). It was highest (1.70 and 0.77 g/kg) under PB+RR+GR and lowest (0.58 and 0.66 g/kg) under CT-RR in 0-5 and 5-15 cm depths, respectively. A similar trend for KMnO $_4$ -C content was observed in both soil depths which was influenced by different tillage practices and followed the order

Table 3 Effect of tillage, residue and nutrient management practices on different SOC fractions (g/kg)

Treatment _	Very la	bile C	Labi	Labile C		Less labile C		Non-labile C	
	0-5 cm	5-15 cm	0-5 cm	5-15 cm	0-5 cm	5-15 cm	0-5 cm	5-15 cm	
Tillage and residue	management (T))							
CT-RR	2.26 ^b	2.41	1.28	0.43 ^b	1.62 ^b	1.47	1.15 ^c	2.34	
CT+RI+GI	2.82 ^a	2.73	1.50	0.61a	1.58 ^b	1.54	2.35 ^b	2.43	
PB+RR	2.65 ^a	2.65	1.62	0.62a	1.92 ^a	1.58	2.93a	2.34	
PB+RR+GR	2.84 ^a	2.43	1.74	0.66a	1.87 ^a	1.69	2.97 ^a	2.53	
LSD (P=0.05)	0.33	NS	NS	0.10	0.19	NS	0.54	NS	
Nutrient managemen	nt practices (N)								
FFP	2.61 ^b	2.49	1.52	0.57	1.74 ^b	1.57	2.04 ^b	2.31	
RDF	2.81 ^a	2.67	1.53	0.59	1.63 ^b	1.56	2.49 ^a	2.39	
SSNM	2.52 ^b	2.50	1.55	0.59	1.87 ^a	1.58	2.52 ^a	2.51	
LSD (P=0.05)	0.13	NS	NS	NS	0.11	NS	0.34	NS	
$T \times N$									
LSD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS	

Table 4 Effect of tillage, residue and nutrient management practices on different forms of soil organic carbon

Treatment	KMnO ₄ -C (g/ kg)		HWE-C (g/kg)		MBC (mg/ kg)					
	0-5	5-15	0-5	5-15	0-5	5-15				
	cm	cm	cm	cm	cm	cm				
Tillage and residue management (T)										
CT-RR	0.58^{c}	0.66^{b}	0.23^{b}	0.18^{a}	183 ^c	136 ^b				
CT+RI+GI	1.29 ^b	0.68^{b}	0.26^{b}	0.18^{a}	235 ^b	155 ^b				
PB+RR	1.61 ^a	0.75 ^a	0.30^{a}	0.25a	258 ^b	172 ^{ab}				
PB+RR+GR	1.70a	0.77^{a}	0.32^{a}	0.26^{a}	310 ^a	211 ^a				
LSD (P=0.05)	0.12	0.05	0.03	0.03	36.9	39.3				
Nutrient manager	ment pr	actices	(N)							
FFP	1.07 ^b	0.66^{b}	0.27	0.21	234 ^b	167				
RDF	1.40a	0.80^{a}	0.28	0.23	229 ^b	166				
SSNM	1.41 ^a	0.69^{b}	0.28	0.23	276a	172				
LSD (P=0.05)	0.13	0.07	NS	NS	31.2	NS				
$T \times N$										
LSD (p=0.05)	0.24	NS	NS	NS	NS	NS				

PB+RR+GR>PB+RR>CT+RI+GI>CT-RR. In surface soil, SSNM and RDF were statistically at par with each other whereas in second 5-15 cm depth, RDF performed better as compared to SSNM. The interaction effect of tillage and nutrient management option on KMnO₄-C was found significant only in surface (0-5 cm) soil. The PB+RR+GR in combination with RDF was recorded highest KMnO₄-C (1.86 g/kg) content among different treatment combinations (Fig 1). Also, the impact of different tillage practices on HWE-C was found similar to that of KMnO₄-C for both 0-5 and 5-15 cm depths (Table 4). Nutrient management options did not showed significant effect on HWE-C in soil depths. Labile pools of SOC (KMnO₄-C and HWE-C) were found sensitive to the alteration in tillage, residue retention as well as nutrient management options. Although the quantities of HWE-C and KMnO₄-C are similar and both fractions probably comprised of C derived from dissolved organic matter and microbial biomass, they are most likely derived from different organic matter fractions. HWE-C largely (45-60%) comprised carbohydrates and amides derived from soil microorganisms, enzymes, root exudates




Fig 1 Percent distribution of different SOC fractions in 0-5 and 5-15 cm soil depth.

and lysates, while KMnO₄-C contains also compounds like lignin and complex polysaccharides (Ghani *et al.* 2003, Mandal *et al.* 2013). None the less, determination of such labile SOC fractions may help to elucidate changes in the SOC pool at early stages of changes under various land use or management system.

Soil MBC, a labile form of organic C, showed a significantly measurable response as a result of no-tillage, residue retention as well as nutrient management options in the present study. In the surface soil, the MBC content was 69.4% higher under PB+RR+GR (310 mg/kg) as compared to CT-RR (183 mg/kg). The individual impact of different nutrient options was found significant in 0-5 cm soil depth, while it showed non-significant results for soil depth of 5-15 cm (Table 4). Among all nutrient options, SSNM registered highest MBC (276 mg/kg), followed by FFP (234 mg/kg) and RDF (229 mg/kg) in surface soil. The interaction of tillage and nutrient options showed no effects in both 0-5 and 5-15 cm soil depths. A favorable moisture content and soil temperature of top soil layer during crop growing season encourages faster residue decomposition with microbial growth and more microbial biomass production (Miltner et al. 2012). Also, the decomposition of residues provides enough energy and C source to soil microorganisms, which might cause an increased microbial activity in soil under PB+ residue plots. Similar findings were reported by other researchers (Nath et al. 2017). Also, in the present study the MBC declined sharply in consecutive depths. Such variability in MBC across soil depths is possible due to lesser source of SOM inputs and consequently lowers microbial activity (Mandal et al. 2013).

Mineral N in soil

A significant increase in mineral-N contents (NH₄⁺ and NO₃⁻) was observed under PB as compared to CT in both soil depths; the values were invariably greater under residue recycling treatments. The NH₄-N under PB+RR+GR was 41.9% and 52.7% higher than CT-RR in 0-5 and 5-15 cm depths, respectively (Table 5). Under the fertilizer management treatments, SSNM and RDF contained comparable NH₄-N and NO₃-N in both depths. The PB+RR+GR in combination with RDF showed significantly highest NH₄-N (13.2 mg/kg) as well as NO₃-N (26.1 mg/kg) compared to other treatment combination, hence showed

more favorable results (Fig 2). As the SOC and N are intrinsically linked with each other through various nutrient cycling processes, the increment in mineral-N may be attributed to higher SOC in soil. Similar results were reported under pigeonpea-wheat system with higher mineral-N contents under PB in sandy loam soils of the IGP (Singh *et al.* 2010). Another possible reason for higher mineral N in PB treatments may be associated with the preparation of raised beds owing to

Table 5 Depth-wise distribution of mineral-N (mg/kg soil) as affected by different tillage, residue and nutrient management practices in soil

Treatment	NI	H ₄ -N	NO ₃ -N		
	0-5 cm	5-15 cm	0-5 cm	5-15 cm	
Tillage and residu	e managei	ment (T)			
CT-RR	7.40^{c}	9.09 ^c	16.5 ^c	13.5°	
CT+RI+GI	9.63 ^b	11.0 ^b	22.5ab	20.2 ^b	
PB+RR	9.92^{ab}	11.2 ^b	21.3 ^b	20.6 ^b	
PB+RR+GR	11.3a	12.9a	24.8a	25.7a	
LSD (P=0.05)	1.39	1.10	2.45	1.27	
Nutrient managen	ient practi	ices (N)			
FFP	8.54 ^b	9.51 ^b	19.9	19.5 ^b	
RDF	10.2a	11.9a	20.4	21.5a	
SSNM	9.97^{a}	11.8a	19.6	22.8a	
LSD (P=0.05)	1.04	1.06	NS	1.78	
$T \times N$					
LSD (P=0.05)	2.19	NS	3.79	NS	

greater accumulation of fertile top soil compared with the conventional (flat beds) crop establishment.

Soil aggregation indices

The impact of different tillage and residue management on soil aggregation stability can be distinguished through




Fig 2 Interaction effect of tillage, residue and nutrient management options on permanganate oxidizable C (KMnO $_4$ -C) in soil depth of 0-5 cm.

aggregation indices, *viz*. mean weight diameter (MWD) and geometric mean diameter (GMD). The MWD was 12.9, 22.8 and 27.3% higher in PB+RR+GR compared with PB+RR, CT+RI+GI and CT-RR, respectively in 0-5 cm soil depth (Fig 3a). Similarly, for 5-15 cm depth MWD was highest under PB + RR + GR (0.70 mm) and lowest in CT-RR (0.55 mm). The SSNM showed maximum MWD (0.63 mm) in 0-5 cm soil, followed by RDF (0.61 mm) and FFP (0.59 mm). The GMD was highest (0.58 mm) in surface soil under PB+RR+GR which was 5.45% greater than CT-RR. On the other hand, result was not significant in other 5-15 cm soil depth. The individual effect of nutrient management options as well as their interaction with tillage and residue

Table 6 Simple correlation (r) between yield, mineral N, SOC pools and microbial biomass C in 0-5 cm soil depth

	Yield	C_{VL}	C_{L}	C_{LL}	C _{NL}	KMnO ₄ -C	HWEC
NH ₄ -N	0.54**						
NO ₃ -N	0.51**						
C_{VL}	0.12^{NS}						
C_{L}	0.45**	0.50**					
C_{LL}	0.47**	0.44**	0.98**				
C _{NL}	0.49**	0.12^{NS}	0.52**	0.46**			
KMnO ₄ -C	0.61**	0.41*	0.77**	0.72**	0.80^{**}		
HWEC	0.36*	0.12^{NS}	0.51**	0.48**	0.57**	0.62**	
MBC	0.43**	0.13^{NS}	0.67**	0.61**	0.55**	0.66**	0.54**

Where * and ** indicate the value of 'r' is significant at 5% and 1% probability levels, respectively.




Fig 3 Interaction effect of tillage, residue and nutrient management options on NH₄-N and NO3-N (mg/kg) in 0-5 cm soil depth.

Table 7 Simple correlation (r) between yield, mineral N, SOC pools and microbial biomass C in 5-15 cm soil depth

	Yield	C _{VL}	C_{L}	C_{LL}	C _{NL}	KMnO ₄ -C	HWEC
NH ₄ -N	0.51**						
NO ₃ -N	0.43**						
C_{VL}	0.42*						
C_{L}	0.43**	0.50**					
C_{LL}	0.43**	$0.29^{\rm NS}$	0.46**				
C_{NL}	0.52**	$-0.03^{ m NS}$	$0.32^{\rm NS}$	0.11^{NS}			
KMnO ₄ -C	$0.20^{ m NS}$	0.34*	0.11^{NS}	0.45**	$0.07^{\rm NS}$		
HWEC	0.40^{*}	$0.26^{\rm NS}$	$0.25^{\rm NS}$	0.69**	0.18^{NS}	0.60^{**}	
MBC	0.40^{*}	0.31^{NS}	0.25^{NS}	0.57**	0.18^{NS}	$0.32^{ m NS}$	0.35^{*}

Where * and ** indicate the value of 'r' is significant at 5% and 1% probability levels, respectively.




Fig 4 Effect of tillage and nutrient management options on soil aggregation indices viz. (a) mean weight diameter (MWD) and (b) geometric mean diameter (GMD).

management showed no significant effects on GMD in either soil depths (Fig 3b). Mandal *et al.* (2019) reported that, long-term application of organic residues improved SOC and soil aggregation such as mean weight diameter (MWD) and geometric mean diameter (GMD). Zhang *et al.* (2013) reported that, the average MWD across all depths of NT and CT were 47% and 20% higher than CT without residue under maize-wheat cropping system.

Relationship of wheat grain yield with mineral N and various SOC pools

Among various pools of SOC measured, KMnO₄-C showed apparently highest correlation with yield than others, in surface soil (Table 6). Different parameters such as KMnO₄-C (r = 0.89; p = 0.01), HWEC (r = 0.56; p = 0.05), C_L(r = 0.91; p = 0.01), MBC (r = 0.64; p = 0.01) as well as mineral-N indicated positive correlation with wheat grain yield and similar relationship between different parameter

was observed for 5-15 cm soil depth (Table 7).

Conclusion

From the present study, it may be concluded that permanent bed with crop residues retention as well as balanced fertilization (RDF and SSNM) under CA improved SOC, mineral N and soil aggregation stability which can lead to increased sustainability under cereal-based intensive cropping systems.

ACKNOWLEDGEMENTS

The authors acknowledge Indian Council of Agricultural Research for financial support provided to conduct this research work. The authors are also grateful to Division of Soil Science and Agricultural Chemistry; Division of Agricultural Physics, ICAR-IARI, New Delhi and International Maize and Wheat Improvement Centre (CIMMYT), Central Soil Salinity Research Institute, Karnal

for supporting this work.

REFERENCES

- Awale R, Emeson M A and Machado S. 2017. Soil organic carbon pools as early indicators for soil organic matter stock changes under different tillage practices in Inland Pacific Northwest. *Frontiers in Ecology and Evolution* **5**: 96.
- Bhattacharyya R, Das T K, Das S, Dey A, Patra A K, Agnihotri R, Ghosh A and Sharma A R. 2019. Four years of conservation agriculture affects topsoil aggregate-associated 15nitrogen but not the 15nitrogen use efficiency by wheat in a semi-arid climate. *Geoderma* 337: 333–340.
- Bhattacharyya R, Tuti M D, Kundu S, Bisht J K and Bhatt J C. 2012. Conservation tillage impacts on soil aggregation and carbon pools in a sandy clay loam soil of the Indian Himalayas. *Soil Science Society of America Journal* **76**:617–627.
- Bolan N S, Adriano D C, Kunhikrishnan A, James T, McDowell R and Senesi N. 2011. *Dissol. Org. Matter* **110**: 1–75.
- Chan K Y, Bowman A and Oates A. 2001. Oxidizible organic carbon fractions and soil quality changes in an Oxic Paleustalf under different pasture leys. *Soil Science* **166**: 61–67.
- Chen H, Hou R, Gong Y, Li H, Fan M and Kuzyakov Y. 2009. Effects of 11 years of conservation tillage on soil organic matter fractions in wheat monoculture in Loess Plateau of China. *Soil and Tillage Research* **106**: 85–94.
- Cooper J, Baranski M, Stewart G, Nobel-de Lange M, Bàrberi P, Fliebach A, Peigné J, Berner A, Brock C, Casagrande M, Crowley O, David C, De VliegherA, Döring T F, Dupont A, Entz M, Grosse M. Haase T, Halde C, Hammerl V, Huiting H, Leithold G, Messmer M, Schloter M, Sukkel W, van der HeijdenM G A, Willekens K, Wittwer R andMäder P. 2016. Shallow non inversion tillage in organic farming maintains crop yields and increases soil C stocks: a meta-analysis. *Agronomy for Sustainable Development* 36: 22.
- Das D, Dwivedi BS, Singh VK, Datta SP and Meena MC. 2017. Long-term impact of fertilizers and organic sources on soil organic carbon fractions under a rice-wheat system in the Indo-Gangetic Plains of northwest India. *Soil Research* **55:** 296–308.
- Dey A, Dwivedi B S, Meena M C and Datta S P. 2018. Dynamics of soil carbon and nitrogen under conservation agriculture in rice-wheat cropping system. *Indian Journal of Fertilisers* 14: 12–26
- Ghani A, Dexter M and Perrott K W. 2003. Hot-water extractable carbon in soils: a sensitive measurement for determining impacts of fertilisation, grazing and cultivation. *Soil Biology and Biochemistry* **35**: 1231–1243.
- Gomez K A and Gomez A A. 1984. *Statistical Procedures for Agricultural Research*, Second Edition. An International Rice Research Institute Book. A Wiley-Inter-science Publication, John Wiley and Sons, New York.
- Jenkinson D S and Ladd J N. 1981. Microbial biomass in soil: measurement and turnover. (In) Pavie E A, Ladd J N (Eds). Soil Biochemistry, Vol 5, pp 415-471. Marcel Dekker, New York.

- Kemper W D and Rosenau R C. 1986. Aggregate stability and size distribution, pp 425–442.
- Mandal N, Dwivedi B S, Datta S P, Meena M C and Tomar R K. 2019. Soil hydro-physical properties under different nutrient management practices, their relationship with soil organic carbon fractions and crop yield under pigeonpea-wheat sequence. *Journal of Plant Nutrition* **42:** 384–400.
- Mandal N, Dwivedi B S, Meena M C, Dhyan-Singh, Datta S P, Tomar R K and Sharma B M. 2013. Effect of induced defoliation in pigeonpea, farmyard manure and sulphitation pressmud on soil organic carbon fractions, mineral nitrogen and crop yields in a pigeonpea-wheat cropping system. *Field Crops Research* **154:** 178–187.
- Meena MC, Dwivedi B S, Dhyan-Singh, Sharma B M, Krishan-Kumar, Singh R V, Radeep Kumar and Rana, D S. 2012. Effect of integrated nutrient management on productivity and soil health in pigeonpea (*Cajanus cajan*)-wheat (*Triticum aestivum*) cropping system. *Indian Journal of Agronomy* 57: 333–337.
- Miltner A., Bombach P, Schmidt-Brucken B and Kastner M. 2012. SOM genesis: microbial biomass as a significant source. *Biogeochemistry* 111: 41–55.
- Nath C P, Das T K, Rana K S, Bhattacharyya R, Pathak H, Paul S, Meena M C and Singh S B. 2017. Greenhouse gases emission, soil organic carbon and wheat yield as affected by tillage systems and nitrogen management practices. *Archives of Agronomy and Soil Science* **63:** 1644–1660.
- Rowell D L. 1994. *Soil Science: Methods and Applications*. Department of Soil Science, University of Reading.
- Singh V K, Dwivedi B S, Shukla A K and Mishra R P. 2010. Permanent raised bed planting of the pigeonpea-wheat system on a TypicUstochrept: Effects on soil fertility, yield, and water and nutrient use efficiencies. *Field Crops Research* 116: 127–139.
- Thomas P, Meena M C, Aggarwal B K, Shri-Ram, Mondal S, Mishra A K and Chakraborty D. 2018. Strength and stability of aggregates as the key indicators for evaluating soil physical conditions. *Journal of the Indian Society of Soil Science* **66**: 268–274
- Tirol-Padre A and Ladha J K. 2004. Assessing the reliability of permanganate-oxcvidizable carbon as an index of soil labile carbon. *Soil Science Society of America Journal* **68**: 969–978.
- vanBavel C H M. 1950. Mean weight-diameter of soil aggregates as a statistical index of aggregation. Soil Science Society of America Journal 14: 20–23.
- Vanlauwe B, Wendt J, Giller K E, Corbeels M, Gerard B and Nolte C. 2014. A fourth principle is required to define conservation agriculture in sub-Saharan Africa: the appropriate use of fertilizer to enhance crop productivity. *Field Crops Research* **155**: 10–13.
- Zhang M Y, Wang F J, Chen F, Malemela M P and Zhang H L. 2013. Comparison of three tillage systems in the wheat-maize system on carbon sequestration in the North China Plain. *Journal of Cleaner Production* 54: 101–107.