

Supplementary Table 1 Per cent disease incidence on wheat crop in different treatments

Treatment	2020–21		2021–22	
	Karnal Bunt (%)	Black Point disease (%)	Karnal Bunt (%)	Black Point disease (%)
T ₁	0.11 (1.655 ^a)	0.91 (5.448 ^a)	0.10 (1.605 ^a)	0.87 (5.329 ^a)
T ₂	0.12 (1.793 ^a)	1.06 (5.895 ^a)	0.11 (1.469 ^a)	1.00 (5.895 ^a)
T ₃	0.11 (1.677 ^a)	0.97 (5.611 ^a)	0.09 (1.334 ^a)	0.93 (5.497 ^a)
Control (C)	0.18 (2.152 ^a)	1.11 (6.029 ^a)	0.16 (2.202 ^a)	1.03 (5.785 ^a)
P-value	NS	NS	NS	NS
CD	-	-	-	-
SE(m)	0.521	0.279	0.481	0.303
SE(d)	0.737	0.395	0.68	0.429
CV	64.024	10.871	65.067	12.05

T₁, Seed treatment (Bavistin® + *T. harzianum*), Foliar spray *T. harzianum*; T₂, Foliar spray Neem Baan®, Confidor®; T₃, Seed treatment Bavistin®, Foliar spray of Tilt®, foliar spray Confidor®; NS, Non-Significant; Figures in parentheses are arc-sine transformed values. Letters in superscript indicate the statistical difference between treatments.