

Supplementary Table 1 Effect of foliar application of nano urea on nutrient content in seed and stover of Indian mustard during 2021–22 and 2022–23

Treatment	Seed concentration (%)			Stover concentration (%)		
	N	P	K	N	P	K
Control	1.70	0.30	0.56	0.41	0.09	0.98
RDF	2.18	0.38	0.65	0.53	0.12	1.14
RDN ₂₅ + NU- FS	1.98	0.34	0.59	0.48	0.10	1.04
RDN ₂₅ + NU-2FS	2.23	0.37	0.64	0.54	0.11	1.12
RDN ₅₀ + NU-1FS	2.32	0.35	0.63	0.52	0.11	1.13
RDN ₅₀ + NU-2FS	2.54	0.40	0.67	0.57	0.13	1.16
RDN ₁₀₀ + NU-1FS	2.74	0.43	0.74	0.66	0.14	1.19
RDN ₇₅ + NU-1FS	2.37	0.38	0.67	0.57	0.12	1.13
RDN ₇₅ + NU-2FS	2.64	0.41	0.70	0.60	0.13	1.17
SEm±	0.11	0.01	0.02	0.03	0.0	0.02
CD (<i>P</i> =0.05)	0.34	0.04	0.05	0.07	0.01	0.06

RDF, Recommended dose of fertilizer (N:P:K=80:40:40); RDN, Recommended dose of nitrogen; NU, Nano urea; FS, Foliar spray.

