



## Management of bud worm (*Helicoverpa armigera*) in Virginia tobacco (*Nicotiana tabacum*) through pest diversionary approach\*

U SREEDHAR

Central Tobacco Research Institute, Rajahmundry 533 105

Received: 13 April 2011; Revised accepted: 24 February 2012

**Keywords:** *Azadirachta indica*, *Helicoverpa armigera*, Neem, *Nicotiana tabacum*, Tobacco, Trap crops

Bud worm (*Helicoverpa armigera* Hub.) is one of the major insect pests of tobacco. It infests the crop during the grand growth period, feeds voraciously on the apical bud and bud leaves, adversely affecting the growth of the plant resulting in considerable yield loss. In Virginia tobacco, *Nicotiana tabacum* L. the loss in green leaf and cured leaf was recorded to be 2 891 and 426 kg/ha, respectively (Sreedhar *et al.* 2005). Besides other adverse effects due to sole dependence on insecticides, the problem of insecticide residues is the major cause of concern in tobacco. Trap cropping is a useful strategy to manage the insect pests, as it can be integrated with other cultural, biological and chemical control methods. Marigold, *Tagetes erecta* L. and rustica tobacco, *Nicotiana rustica* L. planted in perimeter pattern (all around the plot) or as strips in east-west direction were found effective as trap crops for management of budworm, *H. armigera* in FCV tobacco (Sreedhar *et al.* 2007, Sreedhar, 2011). Benefits of trap crop can be greatly improved through various manipulations including a skill full combination with bio-pesticides. Hence, a field experiment was conducted for two seasons, integrating trap crop and bio-pesticide, neem, *Azadirachta indica* A. Juss seed kernel extract (NSKE) to develop an eco-friendly IPM strategy for management of budworm, *H. armigera* in Virginia tobacco.

The experiment was laid out in randomized block design with three replications using cv. *Hema* of FCV tobacco with a plot size of 7 m × 7 m for two seasons during 2006–08. Trap crops, viz. marigold (flowers with single whorl ray florets-sw), marigold with multi whorl ray florets (mw) and rustica tobacco were planted around Virginia tobacco plots. Virginia tobacco was sprayed with NSKE @ 0.5% at 30 and 40 days after planting. These treatments were compared with tobacco plots without trap crop but sprayed with NSKE 0.5% and tobacco plots without trap crop and without NSKE

spray in a replicated field experiment. Observations on budworm infestation on tobacco, population of budworm on trap crops, population of natural enemies on trap crops as well as tobacco were recorded periodically. Yield characters, viz green leaf, cured leaf and bright leaf were recorded, and grade index was calculated and subjected to analysis of variance.

The infestation of *H. armigera* was least (2.7%) in tobacco with marigold-sw + NSKE followed by tobacco with rustica tobacco + NSKE (3.0) which were on par with marigold – mw + NSKE (4.3%) at 40, 50 and 60 days after planting during 2006–07 season (Table 1). Similar trend was observed at 50 and 60 DAP. The infestation (12.7–16.3%) in tobacco plots sprayed with NSKE without trap crop was significantly higher than all other treatments and was found to be on a par with control (15.7–18.0%) in all the observations. The infestation of *H. armigera* was least (2.7%) in tobacco with rustica tobacco + NSKE, followed by tobacco with marigold-sw + NSKE (3.00%) which were on par with marigold –mw + NSKE at 40 days after planting during 2007–08 season. Similar trend was observed at 50 DAP. The infestation in tobacco plots sprayed with NSKE without trap crop was significantly higher (12.0–16.7%) than all other treatments and was found to be at par with control (14.7–18.3%) in all the observations. A Significant reduction in budworm infestation was observed in tobacco plots with trap crops + NSKE spray as compared to plots with trap crop and without NSKE spray during both the seasons.

Virginia tobacco plots with rustica tobacco as trap crop + NSKE spray on Virginia tobacco recorded the highest number of eggs than all other treatments at all the observations i.e. 40, 50 and 60 DAP (15.26, 15.83 and 14.66/plant). All the trap crops planted around Virginia tobacco plots sprayed with NSKE recorded significantly higher number of eggs as compared to trap crops planted around tobacco without NSKE spray (Table 2). Marigold –sw with NSKE spray on tobacco recorded significantly higher number of eggs than all other treatments except rustica tobacco as trap crop + NSKE spray

\*Short note

Principal Scientist (e mail:usreedharctri@yahoo.com), Division of Crop Protection

Table 1 Bud worm infestation in Virginia tobacco

Trap crop	Per cent tobacco plants infested					
	40 DAP		50 DAP		60 DAP	
	2006-07	2007-08	2006-07	2007-08	2006-07	2007-08
Tagetes -mw	7.0 (15.3)	6.3 (14.5)	9.0 (17.3)	8.7 (17.1)	9.7 (18.0)	9.0 (17.4)
Tagetes - sw	5.0 (12.9)	5.7 (13.7)	6.0 (14.1)	6.7 (14.9)	7.0 (15.3)	7.3 (15.7)
Rustica tobacco	5.7 (13.7)	5.3 (13.3)	6.7 (14.9)	6.3 (14.5)	7.3 (15.6)	7.7 (16.0)
Tagetes -mw + NSKE 0.5%	4.3 (11.9)	4.0 (11.5)	5.3 (13.3)	5.0 (12.8)	5.7 (13.7)	5.3 (13.3)
Tagetes -sw + NSKE 0.5%	2.7 (9.3)	3.0 (9.9)	3.0 (9.9)	3.3 (10.3)	3.0 (9.9)	3.3 (10.3)
Rustica tobacco+ NSKE 0.5%	3.0 (9.9)	2.7 (9.3)	3.3 (10.3)	3.0 (9.9)	3.3 (10.5)	3.7 (10.9)
NSKE 0.5%	12.7 (20.78)	12.0 (20.2)	14.0 (21.9)	14.3 (22.29)	16.3 (23.8)	16.7 (24.0)
Control	15.7 (23.2)	14.7 (22.4)	17.7 (24.8)	17.0 (24.3)	18.0 (25.1)	18.3 (25.3)
(Virginia tobacco without trap crop and without NSKE)						
SEm ±	1.11	0.84	1.26	1.03	1.03	1.25
CD at 5%	3.35	2.56	3.82	3.12	3.11	3.81

Figures in parentheses are arc sine transformed values

Table 2 Population of *H.armigera* on trap crops with and without NSKE spray on Virginia tobacco

Trap crop	Mean No. of eggs/plant			Mean No. of larvae/plant		
	40 DAP	50 DAP	60 DAP	40 DAP	50 DAP	60 DAP
	2006-08	2006-08	2006-08	2006-08	2006-08	2006-08
Tagetes -mw	4.82	5.43	4.94	3.63	4.07	3.73
Tagetes - sw	7.00	7.70	6.70	4.63	5.33	4.93
Rustica tobacco	8.20	9.36	8.18	4.93	5.57	5.49
Tagetes -mw + NSKE 0.5%	10.16	11.83	10.01	5.43	6.00	5.93
Tagetes -sw + NSKE 0.5%	13.30	14.86	13.16	6.50	7.66	7.50
Rustica tobacco+ NSKE 0.5%	15.26	15.83	14.66	6.76	7.86	7.83
SEm ±	0.31	0.35	0.31	0.30	0.34	0.32
CD at 5%	0.90	0.50	0.88	0.87	1.00	0.92

on Virginia tobacco at all the observations (13.30, 14.86 and 13.16/plant). Marigold-mw without NSKE spray on tobacco recorded least number of eggs (4.82, 5.43 and 4.94) among all the treatments.

As regards larval population, rustica tobacco with NSKE spray on Virginia tobacco (6.76, 7.86 and 7.83) and marigold-sw + NSKE (6.50, 7.66 and 7.50) recorded significantly higher larval population than all other treatments and were on par with each other. Marigold-mw + NSKE recorded significantly less population (5.43, 6.00 & 5.93) as compared to rustica + NSKE and marigold-sw + NSKE at all the observations. All the trap crops with NSKE spray on tobacco recorded significantly more larval population as compared to trap crops without NSKE spray on tobacco.

On tobacco *Nesidiocoris* sp., coccinellids, spiders and syrphids were recorded of which *Nesidiocoris* sp. was predominant. More coccinellids and syrphids were recorded on marigold-mw and their activity was also found to be more on tobacco with marigold-mw as trap crop. The population

of syrphids was least on rustica tobacco trap crop. More spiders were recorded on marigold-mw followed by marigold-sw. Similarly population of other predators and parasitoids were more on marigold-mw, followed by marigold-sw. No parasitoids were recorded on rustica tobacco. As such the parasitoids on tobacco with or without trap crops were negligible. NSKS spray did not adversely affect the activity of natural enemies on tobacco.

Tobacco plots with marigold-sw + NSKE (2 000 kg/ha) and rustica + NSKE (1 974 kg/ha) recorded significantly higher cured leaf yield than all other treatments. The cured leaf yield in marigold mw + NSKE (1 770 kg/ha) was found to be at par with all the tarp crops without NSKE as well as NSKE alone. Among the treatments significantly higher bright leaf yield was recorded in rustica tobacco and marigold-sw with and without NSKS spray than marigold-mw with and without NSKE, NSKE alone and control. As regards grade index, marigold-sw + NSKE recorded the highest (1 428), followed by rustica tobacco + NSKE (1 426), which were

significantly superior to all other treatments. The grade index in marigold-mw + NSKE (1 323) was at par with all the trap crops without NSKE. From the experimental results based on the pest infestation on tobacco, eggs and larvae of budworm trapped on the trap crops and yield data it can be inferred that conjunctive use of tarp crops, marigold-SW and rustica tobacco with NSKE spray is effective for management of budworm, *H.armigera* in Virginia tobacco. Trap cropping was found effective in several crops in reducing the infestation, enhancing natural enemy activity, management of insecticide resistance in insect pests (Srinivasan *et al.* 2004, Hath and Das 2005, Shelton and Badenes-Perez 2006, Van Den Berg 2006, Khan *et al.* 2006). More trapping of *H.armigera* population on trap crops where the main crop was treated with NSKE may be due to the anti-feedent, repellent and anti ovipositional effects of neem which are well documented (Saxena and Vyas 2005). It can be inferred that use of marigold-sw or rustica tobacco as trap crops along with application of NSKE on tobacco diverts *H.armigera* away from the main crop and can be used successfully for management of the pest in FCV tobacco. The benefits of trap crop can be enhanced when the constraints and risks are fully understood and the strategy is used in conjunction with other methods of control in an integrated pest management system.

#### SUMMARY

Two seasons experimentation for management of *H. armigera* in Virginia tobacco, by integrating trap crop and bio-pesticide, neem seed kernel extract (NSKE) revealed a significant reduction in budworm infestation in Virginia tobacco plots with trap crops + NSKE spray as compared to plots with trap crop and without NSKE spray. All the trap crops planted around Virginia tobacco plots sprayed with NSKE recorded significantly higher number of eggs and larvae as compared to trap crops planted around tobacco without NSKE spray. The natural enemy activity was not affected and was found to be more on tobacco with marigold-mw as trap crop. Conjunctive use of trap crops, marigold-SW or rustica tobacco with NSKE spray was effective for

management of budworm, *H.armigera* in Virginia tobacco and also resulted in better grade index.

#### REFERENCES

- Hath T K and Das B R. 2005. Introduction of trap crops for the management of *Helicoverpa armigera* Hubner in late planted tomato under *terai* agro-ecology of West Bengal. *Research on Crops* **6** : 145–7.
- Khan Z R, Midega C A O, Hutter N J, Wilkins R M, Wadhams L J. 2006. Assessment of the potential of Napier grass (*Pennisetum purpureum*) varieties as trap plants for management of *Chilo partellus*. *Entomologia Experimentalis Et Applicata* **119**: 15–22.
- Saxena R C and Vyas B N. 2005. Role of neem and its products in management of *Helicoverpa armigera* : Recent Advances in *Helicoverpa* management. *Proceedings of the National Symposium on Helicoverpa Management : A National Challenge*. pp. 232–40. Hem Saxena, Rai A B, Ahmed R and Sanjeev Gupta (Eds). IIPR. Kanpur.
- Shelton A M and Badenes-perez, F.R. 2006. Concepts and application of trap cropping in pest management. *Annual Review of Entomology* **51**: 285–308.
- Sreedhar U, Sitaramaiah S and Deo Singh K. 2005. *Helicoverpa* management in tobacco – Present status and Future strategies. Recent Advances in *helicoverpa* management. *Proceedings of the National Symposium on Helicoverpa Management : A National Challenge*. pp 295–303. Hem Saxena, Rai A B, Ahmed R and Sanjeev Gupta (Eds). IIPR. Kanpur.
- Sreedhar U, Sitaramaiah S and Padmaja K. 2007. Evaluation of trap crops for management of budworm, *Helicoverpa armigera* in flue cured Virginia tobacco. *Indian Journal of Plant Protection* **35**: 213–7.
- Sreedhar U. 2011. Influence of trap crop planting geometry on budworm, *Helicoverpa armigera* in flue cured Virginia tobacco. *Indian Journal of Plant Protection* **39**: 196–9.
- Srinivasan K, Krishna Moorthy P N and Raviprasad T N. 2004. African marigold as a trap crop for the management of the fruit borer, *Helicoverpa armigera* on tomato. *International Journal of Pest Management* **40**: 56–63.
- Van Den Berg J. 2006. Ovipositional preference and larval survival of *Chilo partellus* (Lepidoptera:Pyralidae) on Napier grass (*Pennisetum pupureum*) trap crops. *International Journal of Pest Management* **52**: 39–44.

## The Indian Journal of Agricultural Sciences

### ARTICLE CERTIFICATE

F.No.

Article Entitled : .....

Title changed to : .....

Authors (names) : .....

It is certified that:

1. The article has been seen by all the authors (signatures given below), who are satisfied with its form and content.
2. The sequence of names of authors in the by-line is as per their relative contribution to this experiment, giving due credit to all scientists who made notable contribution to it.
3. The address of the organization where the research was conducted is given in the by-line (changes of author's address is given the footnote).
4. The experiment was carried out during 20..... and the article is submitted soon after completion of the experiment.
5. The article is exclusive for this journal, and the results reported here have not been sent (and not be sent during its consideration by this journal) for publication in any other journal.
6. The article has not been rejected for publication in any other journal/rejected in ..... whose comments are attached (such a rejection does not disqualify the article for consideration in this journal).
7. It is based/not based on a part/complete M.Sc/Ph.D thesis of the first author, who submitted the thesis in 20..... to the ..... university, place).
8. Correct data and facts are presented in the article.
9. I/we agree to abide by the objective comments of referees and do agree to modify the article into a shortnote as per the recommendation, for publication in *The Indian Journal of Agricultural Sciences*.
10. If published in *The Indian Journal of Agricultural Sciences*, the copyright of this article would vest with the Indian Council of Agricultural Research who will have the right to enter into any agreement with any organization in India or abroad engaged in reprography, photocopying, storage and dissemination of information contained in it, and neither we nor our legal heirs will have any claims on royalty.

Signature with date	Name in full and designation	Present official address	Subscription number
1. ....	.....	.....	.....
2. ....	.....	.....	.....
3. ....	.....	.....	.....
4. ....	.....	.....	.....
5. ....	.....	.....	.....

#### OPTIONAL

The above certificate is correct to the best of my knowledge and I have no objection to the publication of the article cited above in *The Indian Journal of Agricultural Sciences*

**Signature and address of  
Head of the department /Head of the Institute**