



## Effect of different floral preservatives on reducing foliage discoloration and increasing vase life of chrysanthemum (*Dendranthema × grandiflora*) cv White Reagan

RITU JAIN<sup>1</sup>, T JANAKIRAM<sup>2</sup>, KANWAR PAL SINGH<sup>3</sup> and G L KUMAWAT<sup>4</sup>

Indian Agricultural Research Institute, New Delhi 110 012

Received: 12 August 2013; Revised accepted: 13 June 2014

### ABSTRACT

The experiment was conducted at the Division of Floriculture and Landscaping, Indian Agricultural Research Institute, New Delhi. To enhance the vase life and to reduce the foliage discoloration of chrysanthemum (*Dendranthema × grandiflora* Ramat.) cv. White Reagan, different floral preservative treatments consisting of cobalt chloride, salicylic acid, sodium nitro prusside, aluminium sulphate, 8-HQC and sucrose were used. Experiment was laid out in completely randomized block design with eleven treatment combinations and each treatment was replicated thrice. It was observed that maximum vase life, solution uptake, increase in flower size and minimum foliage discoloration was observed in flowers held in preservative containing 400 ppm 8-HQC and 1.5% sucrose, which was significantly superior over all other treatments.

**Key words:** Chrysanthemum, Cut flowers, Floral preservatives, Foliage discoloration, Salicylic acid, Sodium nitro prusside, Vase life, White Reagan

Chrysanthemum (*Dendranthema × grandiflora* Ramat.) belongs to family Asteraceae and ranks second in world cut flower trade. In India chrysanthemum is gaining popularity on cut flower in the urban floriculture trade. Fresh flower loses their freshness and quality both during handling and transportation. Due to high perishability, flower and foliage parts are vulnerable to large postharvest losses. The main postharvest problems for chrysanthemums are premature foliage yellowing, wilting and the failure of the flowers to fully open. The flowers of chrysanthemum have long vase life but its foliage often becomes yellow, brown and wilted during storage and transportation. Foliage yellowing is cultivar specific problem and is caused by poor production, excessive or improper storage and preservative solutions used at higher than recommended concentrations. Leaves of cut chrysanthemum frequently become yellow spontaneously, sometimes prior to the onset of flower senescence, which makes the flowers unsightly, lowers their quality and shortens vase life (Doi *et al.* 2003, 2004). To preserve the best quality of flowers after harvest and to make them tolerant to fluctuations in environmental conditions treatment with floral preservatives is recommended. Influence of different holding solutions on

chrysanthemum has been reported earlier (Kofranek and Halevy 1972, Marousky 1969, 1971, Talukdar *et al.* 2004) but still the information on controlling leaf discoloration is not available. Therefore, the present studies were carried out to study the effect of various floral preservatives on reducing foliage discoloration and improving the vase life of chrysanthemum cut flower cv. White Reagan.

### MATERIALS AND METHODS

The present studies were carried out in the Division of Floriculture and Landscaping, IARI, New Delhi during 2011-12. For this experiment, flowers of cv. White Reagan were cut to a uniform length of 45 cm and were dressed by removing lower 1/3<sup>rd</sup> leaves. The cut stems were then placed in various holding solutions like cobalt chloride 50 ppm, cobalt chloride 100 ppm, cobalt chloride 150 ppm, salicylic acid 50 ppm, salicylic acid 100 ppm, sodium nitroprusside 50 ppm, sodium nitroprusside 100 ppm, aluminium sulphate 300 ppm+ sucrose 1.5 per cent, 8-HQC 300 ppm+ sucrose 1.5 per cent, 8-HQC 400 ppm+ sucrose 1.5 per cent, control (distilled water). The experiment was laid out in completely randomized design, replicated thrice with three stems per replication. Observations on vase life, per cent weight loss, percent increase in flower size (cm), solution uptake (ml), leaf yellowing (%), leaf wilting (%) and leaf browning (%) were recorded and data was subjected to analysis of variance (Panse and Sukhatme 1967).

<sup>1</sup> e mail: ritujain.uhf@gmail.com, <sup>2</sup> e mail: tolety07@gmail.com, <sup>3</sup> e mail: kanwar\_flspusa@yahoo.com.in, Division of Floriculture and Landscaping

Table 1 Effect of floral preservatives on post harvest handling of chrysanthemum cv. White Reagan

Treatment	Leaf discoloration after 15 days (%)	Leaf discoloration at vase life termination	Vase life (Days)	Solution uptake (ml)	Flower size (cm)	% Physiological weight loss
50 ppm cobalt chloride (T <sub>1</sub> )	18.52 (25.10)	61.54 (51.72)	29.00	74.33	3.97	29.98 (32.55)
100 ppm cobalt chloride (T <sub>2</sub> )	59.53 (50.50)	80.79 (64.17)	26.36	50.00	3.87	44.96 (42.08)
150 ppm cobalt chloride (T <sub>3</sub> )	47.67 (43.32)	55.64 (48.34)	22.43	58.00	3.35	33.75 (35.47)
50 ppm salicylic Acid (T <sub>4</sub> )	36.77 (36.54)	88.16 (73.92)	27.00	98.17	3.92	43.46 (41.22)
100 ppm salicylic Acid (T <sub>5</sub> )	59.61 (50.57)	100.00 (90.00)	30.57	97.33	4.58	58.62 (49.96)
50 ppm sodium nitroprusside (T <sub>6</sub> )	53.20 (46.96)	90.75(72.85)	27.90	75.33	3.78	40.15 (39.26)
100 ppm sodium nitroprusside (T <sub>7</sub> )	100.00 (90.00)	100.00 (90.00)	19.00	87.33	2.92	38.38 (38.25)
300ppm aluminium sulphate + 1.5% sucrose (T <sub>8</sub> )	42.67 (40.76)	96.00 (78.44)	28.77	85.67	3.57	48.89 (44.35)
300ppm 8-HQC + 1.5% sucrose (T <sub>9</sub> )	35.32 (35.70)	68.17 (56.56)	31.00	93.67	4.75	31.54 (34.09)
400ppm 8HQC + 1.5% sucrose (T <sub>10</sub> )	17.09 (24.18)	45.96 (42.64)	33.00	103.67	4.75	37.00 (37.43)
Distilled water (T <sub>11</sub> )	94.68 (79.34)	94.68 (79.34)	23.63	87.00	3.10	49.85 (44.89)
CD (P = 0.05)	24.86 (16.10)	16.9 5(13.71)	3.87	17.20	0.84	11.03 (6.97)

Figures within parentheses are transformed values

## RESULTS AND DISCUSSIONS

It is clear from the Table 1 that flowers held in 100 ppm sodium nitroprusside showed 100 per cent foliage discoloration after 15 days, while minimum leaf discoloration (17.09%) was observed in a solution containing 400 ppm 8-HQC +1.5 % sucrose (T<sub>10</sub>) and was statistically at par with T<sub>4</sub> and T<sub>9</sub>. At the time of termination of vase life minimum leaf discoloration (45.96%) was observed when flowers were held in a solution containing 400 ppm 8-HQC+ 1.5 percent sucrose (T<sub>10</sub>) and was at par with T<sub>1</sub> and T<sub>3</sub>, however, 100 percent leaf discoloration was observed in flowers held in 100 ppm salicylic acid (T<sub>5</sub>) and 100 ppm sodium nitroprusside (T<sub>7</sub>) and it was statistically at par with T<sub>8</sub> and T<sub>11</sub>. This may be due to the reason that exogenous application of sucrose during postharvest handling may preserve chlorophyll loss and hence, prevented yellowing. Our findings are in corroboration with the findings of Reyes-Arribas *et al.* (2000), who observed that floating the leaves of chrysanthemum cv. Tara in one per cent sucrose solution preserved the chlorophyll content and extended the yellowing time from 3- 9 days. Moreover, it also reported earlier that the carbohydrate deprivation commonly occurs in higher plants during senescence (Peoples and Dalling 1988), in darkness (Elmarani *et al.* 1994) and under postharvest conditions (King *et al.* 1990).

Data presented in Table 1 indicates that maximum vase life (33.0 days) was recorded in flowers held in a solution containing 400 ppm 8HQC + 1.5 percent sucrose (T<sub>10</sub>) and it was statistically at par with T<sub>9</sub> and T<sub>5</sub>, while minimum vase life (19.0 days) was observed under treatment T<sub>7</sub>, i.e 100ppm sodium nitroprusside which was statistically at par with T<sub>3</sub>. Among the preservative solutions which are used to increase cut flowers longevity, 8-HQ is more effective to control microbial agents (Kader 2002, Reddy *et al.* 1995, Khalighi and Shafiei 2000). The most probable reason of extending vaselife is that 8- HQ has bacteriostatic properties and along with citric acid it reduces the pH of vase solution.

Reduction of pH in preservative solution controls the growth of bacteria at the cut ends of flower stem and reduces the vascular plugging thereby, increases water uptake and hence, enhanced the vaselife. Hussain *et al.* (2001) reported that sucrose in combination with citric acid or aluminium sulphate maintains endogenous levels of soluble sugars and soluble proteins which in turn provide energy as well as required osmoticum for floret development and longevity. Similar were the findings of Yuniarti *et al.* (2007) who reported that use of holding solution containing citric acid increased the vase life of chrysanthemum cv. Cat Eye's by 9.00 days compared to control. It also corroborates with the studies of Jain *et al.* (2009) who reported the maximum vase life of chrysanthemum cv. Shyamal with 150 ppm citric acid +1000 ppm Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> +2% sucrose. Similarly, Wiraatmaja *et al.*, (2007) also reported that longer vase life (6.02 days over control) of chrysanthemum cut flower with 2.70% of sucrose +400 ppm of citric acid. Similarly, Bartoli *et al.* (1997) and Weiming *et al.* (1997) reported that, the vase life of chrysanthemum cut flowers was significantly increased when treated with 8-HQS +sucrose which was attributed to the inhibition of ethylene action by 8-HQS.

The maximum solution uptake (103.67 ml) was observed when the flowers were kept in a preservative solution containing 400 ppm 8-HQC + 1.5% sucrose (T<sub>10</sub>) and it was statistically at par with T<sub>4</sub>, T<sub>5</sub>, T<sub>7</sub>, T<sub>9</sub> and T<sub>11</sub>. However, minimum solution uptake (50.00ml) was observed when flowers were kept in a solution containing 100 ppm cobalt chloride (T<sub>2</sub>) and was statistically at par with T<sub>3</sub>. The possible reason for maximum uptake includes reduced water stress due to the presence of citric acid and decreased microbial population in vase solution due to hydroxyl quinoline. Sucrose in holding solution acts as osmotically active molecule, leading to the promotion of subsequent water relations. Moreover, vase solution enriched with 8-HQ encourage free flow of the water uptake by elimination of the microbial growth and physiological blockage

(Marousky 1972).

It is evident from the Table 1 that the maximum flower opening (4.75cm) was observed in flowers held in a preservative solution containing 300 ppm 8-HQC +1.5% sucrose and it was statistically at par with T<sub>1</sub>, T<sub>4</sub> and T<sub>5</sub>, however minimum flower size (2.92 cm) was observed with 100 ppm sodium nitroprusside solution (T<sub>7</sub>) and was at par with T<sub>3</sub>, T<sub>8</sub> and T<sub>11</sub>. The flowers kept in preservative solution containing citric acid, biocide and sucrose have maximum solution uptake and exogenous supply of respirable substrate which helped in the maximum opening of flower, because addition of sucrose allow the flower to develop fully which is not possible under control. The data presented in Table 1 shows that minimum physiological loss in weight (29.98%) was observed when flowers were held in a solution containing 50 ppm cobalt chloride (T<sub>1</sub>) and was at par with T<sub>3</sub>, T<sub>9</sub> and T<sub>10</sub>, however, maximum weight loss (58.62%) was observed in flowers held in 100 ppm salicylic acid (T<sub>5</sub>) and was statistically at par with T<sub>8</sub> and T<sub>11</sub>.

### CONCLUSIONS

Comparison of all the treatments reveals that the cut flowers of cv. White Reagan held in preservatives containing 400ppm 8-HQC and 1.5 % sucrose (T<sub>10</sub>) showed maximum vase life, volume of solution uptake, minimum weight loss, minimum foliage discoloration.

### REFERENCES

- Bartoli C G, Juan G and Edgrdo M. 1997. Ethylene production and responses to exogenous ethylene in senescing petals of *Chrysanthemum morifolium* cv. RAM. *Plant Science* **124**: 15–21.
- Doi M, Aoe K, Watabe S, Inamoto K and Imanishi H. 2004. Leaf yellowing of cut standard chrysanthemum (*Dendranthema grandiflora* Kitamura) 'Shuho-no-chikara' induced by ethylene and postharvest increase in ethylene sensitivity. *Journal of Japan Society of Horticultural Science* **73**: 229–34.
- Doi M, Nakagawa Y, Watabe S, Aoe K, Inamoto K and Imanishi H. 2003. Ethylene-induced leaf yellowing in cut chrysanthemum (*Dendranthema grandiflora* Kitamura). *Journal of Japan Society of Horticultural Science* **72**(6): 533–5.
- Elmrani A, Gaudillere J O and Raymond P. 1994. Carbohydrate starvation is a major determinant of the loss of greening capacity in cotyledons of dark green sugar beet seedlings. *Physiologia Plantarum* **91**: 56–64.
- Hussain C T S, Misra RnL, Bhattacharjee S K, and Voleti S R. 2001. Changes in soluble carbohydrates and proteins in cut gladiolus. *Journal of Ornamental Horticulture* **4**(2): 83–6.
- Jain Ritu, Prasad K V and Singh O P. 2009. Effect of different floral preservatives on vase life of few chrysanthemum spray cultivars. *Indian Journal of Ornamental Horticulture* **12**(4): 245–50.
- Kader A A. 2002. Postharvest technology of horticulture crops, Third edition. University of California Publication, p 3311.
- Khalighi A and Shafiei M. 2000. Study on effect of chemical, treatments and stage of harvest on vase life and some qualifical characters of cut Carnation. *Iran Agricultural Sciences*. **31**: 119–25.
- King G A, Woollard D C, Irving D E and Borst W M. 1990. Physiological changes in asparagus spear tips after harvest. *Physiologia Plantarum* **80**: 393–400.
- Kofranek A M and Halevy A M. 1972. Conditions for opening cut chrysanthemum flower buds. *Journal of American Society for Horticulture Science* **97**(5): 578–84.
- Marousky F J. 1969. Influence of various commercial floral preservatives and 8- hydroxy quinoline citrate plus sucrose on development and lasting ability of flower buds of several chrysanthemum cultivars. *Proceedings of Florida State Horticultural Society* **82**: 381–5.
- Marousky F J. 1971. Handling and opening bud cut chrysanthemum flowers with 8- hydroxy quinoline citrate and sucrose. Marketing Research Report, United States Development of Agriculture, No. 905, p 14.
- Marousky F J. 1972. Water relations and effects of floral preservatives on bud opening and keeping quality of cut flowers. *Hort science* **7**: 114–7.
- Panse V G and Sukhatme P V. 1967. *Statistical Methods for Agricultural Workers*. ICAR, New Delhi.
- Peoples M B and Dalling J. 1988. The interplay between proteolysis and amino acid metabolism during senescence and nitrogen reallocation. (In) *Senescence and Aging*, pp 182–217. Nooden L D and Leopold A C (Eds). Academic Press, San Dieago.
- Reyes-Arribas T, Barrett J E, Nell T A and Clark D G. 2000. Effect of ethylene, sucrose and benzyl adenine on leaf senescence of two chrysanthemum cultivars Tara and Boladi. *Acta Horticulturae* **518**:125–9.
- Talukdar M C, Sharma B J, Das S and Mahanta S J. 2004. Effect of pulsing on post harvest quality of chrysanthemum (*Dendranthema grandiflora*, Tzevelv.) under different storage conditions. *Journal of Ornamental Horticulture* **7**(2): 133–40.
- Reddy B S, Singh K and Singh A. 1995. Effect of sucrose, citric acid and hydroxyl quinoline sulfate on postharvest physiology of tuberose 'single'. *Advances in Agricultural Research in India* **3**: 161–7.
- Weiming G, Zeng W and Chen F. 1997. Regulation of ethylene on senescence of cut chrysanthemum. *Journal of Nanjing Agric. University* **20**(4): 24–9.
- Wiraatmaja I W, Astawa I N G and Deviantri D. 2007. Prolonging vase life of chrysanthemum by sucrose and citric acid. *Agritrop* **26** (3): 129–35.
- Yuniarti P E, Prahardini R and Santoso P. 2007. Research on lengthening the shelf life of Cat Eye's chrysanthemum cut flower. *Acta Horticulturae* **755**:109–12.