



## Effect of canopy and nutrient management on senile orchards of Indian gooseberry (*Emblica officinalis*) cv Francis

DIVYA TIWARI<sup>1</sup>, H K SINGH<sup>2</sup> and A K SHUKLA<sup>3</sup>

Narendra Deva University of Agriculture & Technology, Faizabad, Uttar Pradesh

Received: 25 May 2014; Revised accepted: 8 December 2014

### ABSTRACT

A study was conducted on 15 years old senile orchard of Indian gooseberry (*Emblica officinalis* G.) cv Francis to convert into productive through canopy and nutrient management. The results revealed that 50% pruning of previous season growth (P<sub>2</sub>) was statistically superior which resulted in maximum duration of flowering, fruit set and retention, yield, physical and chemical qualities of fruit and leaf as well as soil nutrient status. Further, nutrient management revealed that application of 40 kg FYM (M<sub>1</sub>) resulted maximum fruit set, retention, yield and phosphorus content in leaves and NPK in soil and minimum fruit drop, days taken to maturity and fibre content. Similarly 100 per cent RDF (N<sub>2</sub>) resulted in maximum fruit set (64.90 %), retention (20.51 %), yield (118 kg/tree), nitrogen and phosphorus content in leaves (2.46 and 0.44 %) as well as in soil (238.6 and 17.75 kg/ha). A critical observation of the results showed that interactions had been most effective to improve yield and quality parameters of fruits and nutrient status in leaves of aonla as well as soil than the individual application of different levels of pruning, organic manure and nutrient. Among flowering, fruiting and yield parameters, maximum duration of flowering (29 days), fruit set (74.61 %), retention (28.33 %) and yield (145 kg/tree) and minimum fruit drop (71.67 %) and days taken to fruit maturity (216 days) was found under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. Interaction P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> has been noticed statistically superior to enhance physical and chemical characteristics of fruit followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. This treatment has not only enhanced bearing and quality parameters in aonla but also increased the leaf nutrient status as well as soil fertility status. A significant increased yield of Indian gooseberry with application of pruning intensity, organic and inorganic nutrients was found due to vigorous vegetative growth and increased chlorophyll content, which together accelerated the photosynthesis that increased the supply of carbohydrates to plants. There is a positive correlation among nutrient doses with leaf nutrient content yield and quality of fruits. The supplemented organic manures have improved the physical, chemical and biological activities of soil which helped in better nutrient absorption by plants, resulting higher yield.

**Key words:** Indian gooseberry, Nutrient, Pruning, Sodic

Pruning is very effective to induce healthy current season shoot from older branches causes invigorating effect on Indian gooseberry (*Emblica officinalis* G.) plants and also increase the level of available soil nutrients that might be due to recycling of pruned debris and decrease in utilization of excessive nutrients by undesirable branches. Further, proper nutrient management is also essential to maintain soil fertility and plant nutrient supply to an optimum level for sustaining the desired crop productivity through optimization of the benefits from all possible sources of plant nutrients in an integrated manner. Although, it is highly remunerative crop, yet the productivity and quality of fruits of senile orchards are being declining due to poor management practices as most of the cultivators do not

apply manures and fertilizers that resulted in diminishing the fruit size and quality. Moreover, Francis and Banarasi are the most popular variety among aonla growers of Eastern Uttar Pradesh. Yet the old plantations of these varieties possess shy bearing tendency and produce poor fruit quality hence, they require application of rejuvenation technology to improve the productivity. Due to lack of studies on standardized practices of pruning intensity in relation to improving the bearing potential of aonla trees, So that, there is dire need to improve the health and productivity of such unproductive orchards through application of rejuvenation technology. With the foresaid objectives, an investigation on response of pruning and nutrient management on fruit yield, quality, nutrient status of soil and leaf, has been conducted during the two consecutive years.

### MATERIALS AND METHODS

An experiment was conducted at Main Experiment Station, Department of Horticulture, Narendra Deva

<sup>1</sup>Research Scholar (dtiwari1912@gmail.com), NDUAT, Faizabad. Present address: College of Horticulture, Bihar Agricultural University, Sabour, Noorsarai, Nalanda; <sup>2</sup>Professor and Head, Narendra Deva University of Agriculture & Technology, Faizabad; <sup>3</sup>Principal Scientist (anilciah@gmail.com), DRWA, Bhubaneswar, Odisha 751 003

University of Agriculture and Technology, Kumarganj, Faizabad which is situated at 26°47' N latitude and 82° 12' E longitude at an elevation of 113 m above mean sea level. In order to analyse the leaf nutrient status and fertility levels of the soil, samples were collected up to the depth of 45 cm before initiation and after the termination of the experiment. Before experimentation, the status of N, P and K of leaf and soil were analyzed as 1.84-2.42, 0.30-0.34 and 1.00-1.50% and 229.68-233.98, 15.40-15.61 and 230.01-233.63 kg/ha, respectively. .

The experiment was laid out in a Factorial Randomized Block Design with 3 replications having 12 treatment combinations including 3 levels of pruning (un-pruned, 25% and 50% removal of one year old shoot), 2 levels of organic manure (20 kg/tree bio-pressmud and 40 kg/tree FYM) and inorganic nutrients (50% and 100% Recommended Dose of Fertilizer).

The full dose of phosphorus, potassium, FYM and bio-pressmud were applied as a basal dose in the month of February, while nitrogen was applied in two split doses, one with basal dose and another after fruit bud bursting in the month of September. Plants were pruned in the month of February at an intensity of 25 % and 50 %. Plants were left un-pruned under control. The observations were recorded on per cent fruit set, days taken to fruit maturity and yield, physico-chemical characteristics of fruits, leaf nutrient status (%) and available N, P and K (kg/ha) in soil. The physico-chemical properties of fruits were analysed by the standard methods of AOAC (1990), N, P, K status in leaves were analyzed by Linder (1944) and Richards (1968), organic carbon and available soil N, P and K was analyzed by Baruah and Borthakur (1998), and soil pH is analyzed by Singh *et al.* (1999) whereas soil calcium and magnesium is calculated by the method described by Chopra and Kanwar (1999). The data recorded in both the years, i.e. 2007 and 2008 were analyzed in factorial RBD and pooled data are presented in respective tables.

## RESULTS AND DISCUSSION

### *Yield parameters*

The perusal of data presented in Table 1 and 2 related to response of pruning intensity on duration of flowering, % fruit set, fruit retention, fruit drop and days taken to fruit maturity showed significant effect. The maximum duration of flowering, fruit set and fruit retention was recorded in P<sub>2</sub> (26.0, 70.54 and 23.38 days) and minimum in P<sub>0</sub> (20.70, 57.72 and 16.24 days). Minimum fruit drop occurred in P<sub>2</sub> (76.62 %) followed by P<sub>1</sub> (81.65 %) whereas, maximum fruit drop occurred in un-pruned trees (83.76 %). Significantly maximum yield was found under P<sub>2</sub> (130 kg/tree) and minimum under P<sub>0</sub> (100 kg/tree). The individual response of manures and nutrients on flowering and fruiting characteristics was significant. Similarly, significantly maximum fruit set was noticed in M<sub>2</sub> (65.95%) and N<sub>2</sub> (64.50 %). Due to application of organic manures maximum fruit retention was noticed in M<sub>2</sub> (21.38 %) as compared to M<sub>1</sub> (17.26 %). Fruit drop was significantly reduced by

application of organic manure and inorganic fertilizers. Fruit yield was found to be maximum (119 and 118 kg/tree) under M<sub>2</sub> and N<sub>2</sub> and minimum in M<sub>1</sub> (110 kg/tree) and N<sub>1</sub> (111 kg/tree).

Interaction effect of P×M, P×N, M×N and P×M×N was also significantly affected the flowering and fruiting behavior. The maximum fruit retention was noticed in P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> (28.33 %) followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (24.35 %) and minimum in P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (13.70 %). The minimum fruit drop (73.66, 75.25 and 76.98 %) under P<sub>2</sub>M<sub>2</sub>, P<sub>2</sub>N<sub>2</sub> and M<sub>2</sub>N<sub>2</sub> and maximum (85.40, 84.78, 83.48 %) under P<sub>0</sub>M<sub>1</sub>, P<sub>0</sub>N<sub>1</sub> and M<sub>1</sub>N<sub>1</sub> has been recorded, respectively. It was found minimum in P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> (71.67 %) followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (75.65 %) as compared to other treatments. The fruits matured in minimum 217 days under P<sub>2</sub>M<sub>2</sub>, 216 days under P<sub>2</sub>N<sub>2</sub> and 222 days under M<sub>2</sub>N<sub>2</sub> and M<sub>1</sub>N<sub>2</sub> whereas, it took maximum 231 days under P<sub>0</sub>M<sub>1</sub>, 232 days under P<sub>0</sub>N<sub>1</sub> and 229 days under M<sub>1</sub>N<sub>1</sub>, respectively. It was minimum under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> (216 days) followed by P<sub>2</sub>M<sub>1</sub>N<sub>2</sub> (216 days) and P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (218 days) and maximum under P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (237 days). Maximum (145 kg/tree) has been obtained in P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> which was at par with P<sub>2</sub>M<sub>1</sub>N<sub>2</sub> and P<sub>2</sub>M<sub>1</sub>N<sub>1</sub> (127 and 125 kg/tree) and minimum yield (89 kg/tree) was obtained under P<sub>0</sub>M<sub>1</sub>N<sub>1</sub>. Since flowering in Indian gooseberry occurs on new determinate shoots therefore, pruning helps in getting new fruiting units and thus increases the number of flowers per shoot. Similar findings were also observed by Dhaliwal *et al.* (2000), Dalal *et al.* (2000) and Singh (2005).

A significant increase in yield and yield parameters in Indian gooseberry with nutrient application would have been due to vigorous vegetative growth and increased chlorophyll content, which together accelerated the photosynthesis that increased the supply of carbohydrates to plants. In addition, it may also be due to more number of female flowers/shoot and higher fruit retention and reduced fruit drop. The beneficial role of supplemented organic manures in improving soil physical, chemical and biological activities is well known, that helped in better nutrient absorption by plants, resulting higher yield (Prabu *et al.*, 2002). However, other factors like cultural practices and nutrition of the plants also influenced flowering characters to an appreciable extent (Tripathi and Maity 2007). The significant interactive effect as a consequence of organic sources and fertilizers are attributed to the favourable nutritional status of the soil resulting into increased biomass production of the crop. The results of the present study are in conformity with Totawat *et al.* (2001), Singh and Swaroop (2000) and Swarup and Wanjari (2000).

### *Physical characteristics of the fruits*

Different level of pruning intensity significantly influenced equatorial diameter, polar length, weight, volume, pulp stone ratio and fibre content of Indian gooseberry fruit. Effect of P×M×N shows that the application of P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> produced significantly larger equatorial diameter (4.69 cm) of fruit followed by P<sub>2</sub>M<sub>1</sub>N<sub>1</sub> (4.54 cm) whereas, the minimum equatorial fruit diameter was recorded under

Table 1 Effect of the pruning intensity and nutrient management on duration of flowering, fruit set and fruit retention

Treatment	Duration of flowering (days)						Fruit Set (%)						Fruit Retention (%)														
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean									
P <sub>0</sub>	20.4	21.0	18.9	22.5	18.7	22.0	19.0	23.0	20.7	53.79	61.65	56.43	59.01	53.18	54.39	59.67	63.63	57.72	14.60	17.88	15.23	17.25	13.70	15.50	16.75	19.00	16.24
P <sub>1</sub>	23.5	28.0	25.0	26.5	24.0	23.0	26.0	30.0	25.8	58.65	62.90	58.14	63.41	55.81	61.48	60.46	65.34	60.78	16.78	19.93	17.17	19.53	16.20	17.33	18.13	21.73	18.35
P <sub>2</sub>	26.0	26.0	24.0	28.0	25.0	27.0	23.0	29.0	26.0	67.77	73.31	69.99	71.09	67.98	67.56	72.00	74.61	70.54	20.42	26.34	22.01	24.75	19.67	21.17	24.35	28.33	23.38
Mean	23.3	25.0	22.6	25.7	22.6	24.0	22.7	27.3	24.2	60.07	65.95	61.52	64.50	58.99	61.14	64.04	67.86	63.01	17.26	21.38	18.13	20.51	16.52	18.00	19.74	23.02	19.32
	<i>SEM</i> ±						<i>SEM</i> ±						<i>SEM</i> ±														
	0.49						0.850						2.487														
Pruning intensity (P)	0.41						1.42						1.048														
Manure (M)	0.41						1.16						0.293														
Nutrients (N)	0.69						1.16						0.856														
P×M	0.69						2.02						1.482														
P×N	0.69						2.02						1.482														
M×N	0.56						1.65						1.210														
P×M×N	0.97						2.85						2.096														
	<i>CD (P = 0.05)</i>						<i>CD (P = 0.05)</i>						<i>CD (P = 0.05)</i>														

Table 2 Effect of the pruning intensity and nutrient management on fruit drop, days taken to fruit maturity and fruit yield

Treatment	Fruit drop (%)						Days taken to fruit maturity (%)						Fruit yield (kg/tree)														
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean									
P <sub>0</sub>	85.40	82.13	84.78	82.75	86.30	84.50	83.25	81.00	83.76	231*	228	232	227	237	225	227	228	229	94	107	97	103	89	98	105	108	100
P <sub>1</sub>	83.24	82.07	82.84	80.47	83.80	82.67	81.87	78.27	81.65	226	223	225	223	227	224	223	222	224	113	115	112	116	110	115	113	116	114
P <sub>2</sub>	79.58	73.66	77.99	75.25	80.33	78.83	75.65	71.67	76.62	220	217	221	216	224	216	218	216	218	124	124	135	136	121	127	125	145	130
Mean	82.74	78.62	81.87	79.49	83.48	82.00	80.26	76.98	80.68	226	222	226	222	229	222	223	222	224	110	119	111	118	107	113	114	123	115
	<i>SEM</i> ±						<i>SEM</i> ±						<i>SEM</i> ±														
	0.358						0.399						1.384														
Pruning intensity (P)	0.293						0.325						1.130														
Manure (M)	0.293						0.325						1.130														
Nutrients (N)	0.507						0.564						1.958														
P×M	0.507						0.564						1.958														
P×N	0.414						0.460						1.598														
M×N	0.717						0.797						2.768														
P×M×N	1.048						1.166						4.050														
	<i>CD (P = 0.05)</i>						<i>CD (P = 0.05)</i>						<i>CD (P = 0.05)</i>														

Table 3 Effect of the pruning intensity and nutrient management on equatorial diameter, polar length and fruit weight

Treatment	Equatorial diameter of fruit (cm)						Polar length of fruit (cm)						Fruit weight (g)												
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean							
P <sub>0</sub>	3.07	4.01	3.39	3.68	3.00	3.13	3.77	4.24	3.54	3.24	3.22	3.52	3.20	3.28	3.24	3.75	41.74	41.67	40.93	42.48	40.33	43.15	41.53	41.80	41.70
P <sub>1</sub>	4.35	3.91	4.28	3.97	4.31	4.38	4.25	3.57	4.13	3.83	3.48	3.82	3.49	3.70	3.95	3.94	3.66	45.13	45.01	44.54	45.60	46.33	43.93	42.75	47.26
P <sub>2</sub>	4.34	4.50	4.43	4.41	4.54	4.14	4.31	4.69	4.42	4.08	4.38	4.15	4.31	4.12	4.04	4.17	4.58	47.14	58.59	53.34	52.39	51.67	42.60	55.00	62.18
Mean	3.92	4.14	4.03	4.02	3.95	3.88	4.11	4.17	4.03	3.72	3.79	3.73	3.77	3.67	3.76	3.78	3.75	44.67	48.42	46.27	46.82	46.11	43.23	46.43	50.41
	<i>SEm</i> ±						<i>CD</i> ( <i>P</i> = 0.05)						<i>SEm</i> ±						<i>CD</i> ( <i>P</i> = 0.05)						
Pruning intensity (P)	0.144						0.422						1.177						3.443						
Manure (M)	0.118						NS						0.961						2.811						
Nutrients (N)	0.118						NS						0.961						NS						
P×M	0.204						0.596						1.665						4.871						
P×N	0.204						0.596						1.665						4.871						
M×N	0.166						NS						1.359						NS						
P×M×N	0.288						0.843						2.354						6.887						

Table 4 Effect of the pruning intensity and nutrient management on fruit volume, pulp stone ratio and fibre content

Treatment	Fruit volume (cc)						Pulp stone ratio						Fibre content (%)															
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean										
P <sub>0</sub>	40.04	39.27	39.12	40.19	38.50	41.57	39.73	38.8	39.65	17.4:1	17.3:1	17.2:1	17.5:1	17.0:1	17.7:1	17.3:1	17.2:1	17.3:1	17.3:1	0.035	0.031	0.034	0.033	0.036	0.034	0.031	0.031	0.033
P <sub>1</sub>	43.31	43.38	43.10	43.60	44.76	41.86	41.43	45.33	43.35	18.5:1	18.3:1	18.2:1	18.6:1	18.9:1	18.2:1	17.6:1	19.0:1	18.4:1	18.4:1	0.026	0.022	0.025	0.023	0.027	0.024	0.022	0.021	0.024
P <sub>2</sub>	45.19	56.56	51.50	50.25	49.79	40.59	53.20	59.91	50.88	23.1:1	23.1:1	21.3:1	21.2:1	20.9:1	18.0:1	21.7:1	24.4:1	21.3:1	21.3:1	0.019	0.011	0.015	0.015	0.018	0.019	0.011	0.011	0.015
Mean	42.85	46.41	44.57	44.68	44.35	41.34	44.79	48.01	44.63	18.5:1	19.5:1	18.9:1	19.1:1	18.9:1	18.0:1	18.9:1	20.2:1	19.0:1	19.0:1	0.026	0.021	0.025	0.023	0.027	0.026	0.021	0.021	0.024
	<i>SEm</i> ±						<i>CD</i> ( <i>P</i> = 0.05)						<i>SEm</i> ±						<i>CD</i> ( <i>P</i> = 0.05)									
Pruning intensity (P)	1.220						3.569						0.373						0.003									
Manure (M)	0.996						2.914						0.304						0.003									
Nutrients (N)	0.996						NS						0.304						0.003									
P×M	1.725						5.048						0.527						0.004									
P×N	1.725						5.048						0.527						0.004									
M×N	1.409						4.121						0.430						0.003									
P×M×N	2.440						7.139						0.746						0.006									

Table 5 Effect of the pruning intensity and nutrient management on TSS and Vitamin C

Treatment	Total soluble solids ( <sup>o</sup> Brix)									Vitamin C (mg/100 g pulp)								
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean
P <sub>0</sub>	11.28	11.62	11.42	11.48	11.83	10.73	11.00	12.23	11.45	443.2	461.0	446.7	457.5	433.3	453.0	460.0	462.0	452.1
P <sub>1</sub>	12.44	12.43	12.40	12.47	12.47	12.40	12.33	12.53	12.43	481.8	489.0	485.0	485.8	488.0	475.7	482.0	496.0	485.4
P <sub>2</sub>	13.32	13.85	13.37	13.82	13.00	13.67	13.74	13.97	13.59	478.7	486.8	477.5	487.9	478.7	478.7	476.3	497.2	482.7
Mean	12.35	12.63	12.41	12.59	12.43	12.27	12.36	12.91	12.49	467.9	478.9	469.7	477.1	466.7	469.1	472.8	485.1	473.4
	SEm ±			CD (P = 0.05)						SEm ±			CD (P = 0.05)					
Pruning intensity (P)	0.213			0.623						1.776			5.195					
Manure (M)	0.174			NS						1.451			4.241					
Nutrients (N)	0.174			NS						1.451			4.241					
P×M	0.301			0.881						2.511			7.346					
P×N	0.301			0.881						2.511			7.346					
M×N	0.246			NS						2.050			5.998					
P×M×N	0.426			1.245						3.551			10.389					

Table 6 Effect of the pruning intensity and nutrient management on total sugars and reducing sugar

Treatment	Total sugars (%)									Reducing sugar (%)								
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean
P <sub>0</sub>	4.53	4.63	4.60	4.55	4.50	4.55	4.60	4.65	4.58	2.36	2.34	2.36	2.34	2.33	2.39	2.39	2.28	2.35
P <sub>1</sub>	4.72	4.82	4.74	4.79	4.75	4.69	4.84	4.79	4.77	2.41	2.40	2.39	2.42	2.40	2.41	2.38	2.42	2.40
P <sub>2</sub>	4.83	4.91	4.86	4.87	4.85	4.80	4.90	4.92	4.87	2.47	2.47	2.51	2.44	2.51	2.45	2.51	2.42	2.48
Mean	4.69	4.78	4.73	4.74	4.70	4.68	4.78	4.79	4.74	2.41	2.41	2.42	2.40	2.41	2.41	2.43	2.37	2.41
	SEm ±			CD (P = 0.05)						SEm ±			CD (P = 0.05)					
Pruning intensity (P)	0.006			0.016						0.024			0.071					
Manure (M)	0.005			0.013						0.021			NS					
Nutrients (N)	0.005			NS						0.021			NS					
P×M	0.008			0.023						0.034			0.101					
P×N	0.008			0.023						0.034			0.101					
M×N	0.006			0.019						0.028			NS					
P×M×N	0.011			0.033						0.049			0.143					

Po, No pruning, P1, 25% removal of one year old shoot, P2, 50% removal of one year old shoot, M1, 20 kg/tree bio-pressmud, M2, 40 kg FYM /tree, N1, 50% of RDF, N2, 100% of RDF

treatment P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (3.0 cm). The maximum polar fruit length (4.58 cm) was recorded under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (4.17 cm). While, minimum polar length (3.20 cm) was found under P<sub>1</sub>M<sub>1</sub>N<sub>1</sub>. Under interaction of P×M×N maximum fruit weight (62.18 g) in P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (55.00 g) and minimum (40.33 g) under P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> was recorded. The effect of P×M, P×N, M×N and P×M×N on fruit volume, pulp stone ratio and fibre content was found significant. The maximum fruit volume (59.91 cc) was observed under the treatment P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (53.20 cc) as compared to other treatments and minimum fruit volume was obtained under the treatment P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (38.50 cc). Similarly, it was found maximum (24.4:1) under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> and minimum (17.0:1) under P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (Table 4). The results are in accordance with findings of Dhaliwal *et al.* (2000) and Shaaban *et al.* (2006). Use of organic sources leaves better effect on physico-chemical properties of the soil that helps in improvement of soil structure for favorable root growth which caused better plant growth and fruit bearing as compared to chemical fertilizers. These results are in conformity with findings of

Ghugre *et al.* (2007).

A synergistic interaction between organic manures and inorganic fertilizers has resulted in enhanced production of growth promoting substances like gibberellic acid, indole acetic acid and dihydrozeatin. They have positive influence on the physiological activity of the plants that enhanced fruit length and diameter, which ultimately increased the average fruit weight. Combined application of organic manures along with fertilizers might have helped in strengthening process in the nucleic acid and metabolism. Obviously, it has increased utilization of native N and P due to organic acids produced during decomposition and induced chelating effects on micronutrients which probably enhanced the availability of N, P and K along with solubilizing the availability of micronutrients. Thus, application of organic manures and fertilizers resulted in an overall significant increase in N, P and K in plants at lesser cost but longer in durability. The combined use of pruning, organic manures and fertilizers has been found effective not only in maintaining higher productivity but also in providing stable crop yield for sustainable crop production through integrated

Table 7 Effect of the pruning intensity and nutrient management on nitrogen, phosphorus and potassium content in branchlet

Treatment	Nitrogen content in branchlet (%)						Phosphorus content in branchlet (%)						Potassium content in branchlet (%)															
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean										
P <sub>0</sub>	2.18	2.14	1.93	2.20	2.46	2.18	2.45	2.22	2.47	2.33	0.41	0.43	0.40	0.44	0.38	0.43	0.41	0.45	0.42	1.14	1.33	1.24	1.23	1.07	1.20	1.41	1.25	1.23
P <sub>1</sub>	2.34	2.43	2.29	2.47	2.24	2.43	2.34	2.51	2.38	0.42	0.43	0.39	0.46	0.39	0.44	0.39	0.47	0.42	1.37	1.49	1.34	1.52	1.24	1.50	1.44	1.54	1.43	
P <sub>2</sub>	2.43	2.47	2.38	2.52	2.35	2.41	2.53	2.45	0.43	0.48	0.42	0.49	0.38	0.47	0.45	0.50	0.45	1.82	1.56	1.58	1.80	1.77	1.87	1.39	1.72	1.69		
Mean	2.31	2.34	2.20	2.46	2.18	2.45	2.22	2.47	2.33	0.41	0.43	0.40	0.44	0.38	0.43	0.41	0.45	0.42	1.44	1.46	1.38	1.51	1.36	1.52	1.41	1.50	1.45	
	<i>SEm</i> ±																											
Pruning intensity (P)	0.019						0.054						0.054						<i>SEm</i> ±									
Manure (M)	0.015						NS						0.015						0.099									
Nutrients (N)	0.015						0.044						0.015						0.081									
P×M	0.026						0.077						0.026						0.081									
P×N	0.026						0.077						0.026						0.141									
M×N	0.021						0.063						0.021						0.141									
P×M×N	0.037						0.109						0.037						0.198									
	<i>CD (P = 0.05)</i>																											

Table 8 Effect of the pruning intensity and nutrient management on nitrogen and phosphorus content in soil (kg/ha)

Treatment	Nitrogen content in soil (kg/ha)						Phosphorus content in soil (kg/ha)											
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean
P <sub>0</sub>	234.9	237.5	235.4	237.0	233.9	235.9	236.9	238.2	236.2	16.82	17.34	16.97	17.19	16.72	16.92	17.22	17.45	17.08
P <sub>1</sub>	235.0	238.4	235.6	237.9	234.1	235.9	237.1	239.8	236.7	17.02	17.39	17.10	17.31	16.95	17.09	17.25	17.52	17.20
P <sub>2</sub>	240.0	240.8	239.8	241.0	239.1	240.9	240.5	241.1	240.4	18.21	18.96	18.41	18.76	17.85	17.56	18.96	18.96	18.58
Mean	236.6	238.9	236.9	238.6	235.7	237.6	238.2	239.7	237.8	17.35	17.89	17.49	17.75	17.17	17.19	17.81	17.98	17.62
	<i>SEm</i> ±																	
Pruning intensity (P)	0.463						1.355						0.075					
Manure (M)	0.378						1.106						0.061					
Nutrients (N)	0.378						1.106						0.061					
P×M	0.655						1.916						0.106					
P×N	0.655						1.916						0.106					
M×N	0.535						1.564						0.087					
P×M×N	0.926						2.709						0.150					
	<i>CD (P = 0.05)</i>																	

P<sub>0</sub>, No pruning, P<sub>1</sub>, 25% removal of one year old shoot, P<sub>2</sub>, 50% removal of one year old shoot, M<sub>1</sub>, 20 kg/tree bio-pressmud, M<sub>2</sub>, 40 kg FYM /tree, N<sub>1</sub>, 50% of RDF, N<sub>2</sub>, 100% of RDF

Table 9 Effect of the pruning intensity and nutrient management on potassium content in soil and organic carbon in soil

Treat- ment	Potassium content in soil (kg/ha)									Organic carbon in soil (%).								
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean
P <sub>0</sub>	237.8	240.8	238.6	240.1	237.5	238.2	239.7	241.9	239.3	0.38	0.39	0.38	0.38	0.38	0.37	0.38	0.39	0.38
P <sub>1</sub>	240.7	243.4	241.1	243.0	239.6	241.9	242.5	244.2	242.0	0.40	0.42	0.39	0.43	0.39	0.41	0.39	0.44	0.41
P <sub>2</sub>	243.8	246.0	244.4	245.5	242.9	244.8	245.9	246.2	244.9	0.43	0.47	0.45	0.45	0.44	0.42	0.47	0.47	0.45
Mean	240.8	243.4	241.3	242.8	240.0	241.6	242.7	244.1	242.1	0.40	0.42	0.40	0.42	0.40	0.40	0.41	0.43	0.41
	<i>SEm</i> ±			<i>CD</i> ( <i>P</i> = 0.05)						<i>SEm</i> ±			<i>CD</i> ( <i>P</i> = 0.05)					
Pruning intensity (P)	0.563			1.647						0.012			0.035					
Manure (M)	0.461			1.345						0.011			NS					
Nutrients (N)	0.461			1.345						0.011			NS					
P×M	0.796			2.331						0.017			0.050					
P×N	0.796			2.331						0.017			0.050					
M×N	0.650			1.902						0.014			NS					
P×M×N	1.126			3.295						0.024			0.071					

nutrient use. The improvement in physical characteristics of the fruit would have been due to remarkable improvement in N, P and K content of plant under the influence of combined application of nutrients through integrated use resulting greater availability of these nutrients in the soil. Besides, greater availability of nutrients, better development of fruit was due to more absorption of nutrients from soil and their efficient translocation in plant system (Jaggi 2007).

#### Chemical characteristics of fruits

The data on total soluble solids and vitamin C are presented in Table 5 showed significant effect on enhancement of TSS and vitamin C. The highest value (13.59° Brix) under P<sub>2</sub> and lowest (11.45° Brix) under P<sub>0</sub> was recorded. The vitamin C was found maximum (485.4 mg/100g pulp) under P<sub>1</sub> and minimum (452.1 mg/100g pulp) under P<sub>0</sub>. The observations on total sugars and reducing sugar content of Indian gooseberry are presented in Table 6. Significant influence is seen by different levels of pruning. Highest total sugar and (4.87 and 2.48%) in P<sub>2</sub> and lowest (4.58 and 2.35%) in P<sub>0</sub> has been recorded. However, organic manures and nutrients significantly influenced the vitamin C content of fruit. However, application of fertilizers alone has non-significant effect on total sugars. Interaction effect of P×M, P×N and P×M×N on TSS was found to be significant but effect of M×N was non-significant. Maximum TSS was found under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> (13.97° Brix) and minimum under P<sub>0</sub>M<sub>1</sub>N<sub>2</sub> (10.73° Brix). Effect of P×M, P×N, M×N and P×M×N on ascorbic acid was found to be significant. Maximum value (489.0, 487.9, 485.1 and 497.2 mg/100g pulp) was recorded with P<sub>1</sub>M<sub>2</sub>, P<sub>2</sub>N<sub>2</sub>, M<sub>2</sub>N<sub>2</sub> and P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> while, minimum vitamin C content was recorded with P<sub>0</sub>M<sub>1</sub>, P<sub>0</sub>N<sub>1</sub>, M<sub>1</sub>N<sub>1</sub> and P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> (443.2, 446.7, 446.7 and 433.3 mg/100g pulp), respectively. In the same way, maximum total sugar (4.92 %) was obtained under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by 4.90 % under P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. Beneficial effect of pruning on quality parameters of Indian gooseberry are in conformity with findings of Dhaliwal *et al.* (2000).

#### Nutrient status of branchlet

A perusal of pooled data presented in Table 7 reveal that application of pruning intensity has significant influence on nitrogen, phosphorus and potassium content in branchlet of Indian gooseberry during experimentation. Due to interactions P×M, P×N, M×N and P×M×N maximum nitrogen (2.47, 2.52 and 2.47 %) under P<sub>2</sub>M<sub>2</sub>, P<sub>2</sub>N<sub>2</sub>, M<sub>2</sub>N<sub>2</sub> and minimum (2.14, 1.93 and 2.18 %) under P<sub>0</sub>M<sub>2</sub>, P<sub>0</sub>N<sub>1</sub> and M<sub>2</sub>N<sub>2</sub> is seen. Further, it is maximum (2.50 %) under P<sub>2</sub>M<sub>1</sub>N<sub>2</sub>.

The effect of P×M, P×N, M×N and P×M×N was found significant where maximum phosphorus (0.48, 0.49, 0.45 and 0.50 %) was found due to P<sub>2</sub>M<sub>2</sub>, P<sub>2</sub>N<sub>2</sub>, M<sub>2</sub>N<sub>2</sub> and P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> and minimum (0.38, 0.39, 0.38 and 0.38 %) due to P<sub>0</sub>M<sub>1</sub>, (P<sub>0</sub>N<sub>1</sub>, P<sub>0</sub>N<sub>2</sub> and P<sub>1</sub>N<sub>1</sub>), M<sub>1</sub>N<sub>1</sub> and P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> and P<sub>0</sub>M<sub>1</sub>N<sub>2</sub> respectively.

Maximum potassium content in branchlet (1.82, 1.80 and 1.87 %) under P<sub>2</sub>M<sub>1</sub>, P<sub>2</sub>N<sub>2</sub> and P<sub>2</sub>M<sub>1</sub>N<sub>2</sub> and minimum (1.14, 1.23 and 1.07 %) under P<sub>0</sub>M<sub>1</sub>, P<sub>0</sub>N<sub>2</sub> and P<sub>0</sub>M<sub>1</sub>N<sub>1</sub> was recorded. Increase in the nutrient status of the leaves would have been due to enhanced invigorating effect of pruning severity on the absorption of mineral content by the plants which could have increased the leaf nutrient status of the plant. Results are in line with Naik and Babu (2005).

#### Soil nutrient status of the orchard

Post analysis of soil samples were done to compute the status of N, P and K to verify the effect of treatments. The values recorded under Table 8 reveal that pruning levels significantly improved available nitrogen (240.4 kg/ha) in soil under P<sub>2</sub> with respect to P<sub>0</sub> (236.2 kg/ha). Again significantly highest phosphorus value (18.58 kg/ha) in P<sub>2</sub> and minimum (17.08 kg/ha) in P<sub>0</sub> was found. As evident from the pooled data on available potassium and organic carbon in soil presented in Table 9 wherein, maximum potassium value (244.9 kg/ha) and organic carbon (0.45 %) under P<sub>2</sub> and minimum K value (239.3 kg/ha) and organic carbon (0.38 %) under P<sub>0</sub>, respectively was recorded. However, pruning effect enhanced calcium and magnesium

Table 10 Effect of the pruning intensity and nutrient management on calcium and magnesium content in soil

Treatment	Calcium content in soil (m.e./100 g)									Magnesium content in soil (m.e./100 g)								
	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean	M <sub>1</sub>	M <sub>2</sub>	N <sub>1</sub>	N <sub>2</sub>	M <sub>1</sub> N <sub>1</sub>	M <sub>1</sub> N <sub>2</sub>	M <sub>2</sub> N <sub>1</sub>	M <sub>2</sub> N <sub>2</sub>	Mean
P <sub>0</sub>	3.88	3.90	3.88	3.90	3.87	3.88	3.89	3.91	3.89	5.80	5.87	5.80	5.86	5.74	5.85	5.87	5.88	5.83
P <sub>1</sub>	3.94	3.97	3.94	3.97	3.92	3.96	3.95	3.98	3.95	5.89	5.91	5.90	5.90	5.89	5.89	5.90	5.91	5.90
P <sub>2</sub>	3.98	3.99	3.98	3.99	3.98	3.98	3.98	3.99	3.98	5.94	5.97	5.95	5.97	5.93	5.95	5.96	5.98	5.96
Mean	3.93	3.95	3.93	3.95	3.92	3.94	3.94	3.96	3.94	5.87	5.92	5.88	5.91	5.85	5.91	5.91	5.92	5.91
	<i>SEm</i> ±				<i>CD</i> ( <i>P</i> = 0.05)					<i>SEm</i> ±				<i>CD</i> ( <i>P</i> = 0.05)				
Pruning intensity (P)	0.008				0.023					0.017				0.051				
Manure (M)	0.007				0.019					0.014				0.041				
Nutrients (N)	0.007				0.019					0.014				NS				
P×M	0.011				0.033					0.024				0.071				
P×N	0.011				0.033					0.024				0.071				
M×N	0.009				0.027					0.021				0.058				
P×M×N	0.016				0.047					0.035				0.101				

availability in soil significantly. Under pruning maximum calcium (3.98 m.e./100 g) in P<sub>2</sub> and minimum (3.89 m.e./100 g) in P<sub>0</sub> was found. Similarly, maximum magnesium content (5.96 m.e./100 g) was found under P<sub>2</sub> and minimum (5.83 m.e./100 g) under P<sub>0</sub> was found. Application of organic manure and inorganic fertilizers has also significant effect on improvement of available nitrogen and phosphorus. Data presented in Table 9 showed significant effect due to application of organic manure and fertilizers on potassium and organic carbon where maximum potassium (243.4 kg/ha) and organic carbon (0.42 %) in M<sub>2</sub> and minimum (240.8 kg/ha) and organic carbon (0.40 %) in M<sub>1</sub>, respectively.

A perusal of data presented in Table 10 reveal that under application of organic manure, maximum calcium content (3.95 m.e./100 g) in M<sub>2</sub> and minimum (3.93 m.e./100 g) in M<sub>1</sub> was found due to interactions P×M, P×N, M×N and P×M×N phosphorus content is significant. Due to P×M, P×N, M×N and P×M×N potassium content, organic carbon, Calcium and magnesium content of soil improved significantly. Balanced use of fertilizer and integrated approach would have been attributed in enhanced rhizospheric effect and soil biological activity, providing congenial physical condition to soil that conserved the soil NPK and increased the availability of other nutrients being its constituents. Increase in the level of available soil nutrients might be due to recycling of pruned debris and decrease in utilization of excessive nutrients by undesirable branches. The present findings are in conformity with Ray *et al.* (2005).

### CONCLUSION

The results on present investigation revealed that among different pruning intensity, 50% pruning of previous season growth (P<sub>2</sub>) was statistically superior which resulted maximum plant height, spread, duration of flowering, fruit set and retention, yield, physical and chemical qualities of fruit and leaf as well as soil nutrient status. Further, nutrient management revealed that application of 40 kg FYM (M<sub>1</sub>) resulted maximum fruit set, retention, yield and phosphorus

content in leaves and NPK in soil and minimum fruit drop, days taken to maturity and fibre content. Similarly 100 % Recommended Dose of Fertilizer (N<sub>1</sub>) resulted maximum fruit set, retention, yield, nitrogen and phosphorus content in leaves as well as in soil.

Among the interactions, P<sub>2</sub>M<sub>2</sub>, P<sub>2</sub>N<sub>2</sub> and M<sub>2</sub>N<sub>2</sub> was found superior over others in improving the vegetative and reproductive plant growth beside physical and chemical quality of fruits along with nutrient status of branchlet and soil. A critical observation of the results, it is concluded that interactions had been most effective to improve the vegetative and reproductive characteristics, quality parameters of fruits and nutrient status in leaves of aonla as well as soil than the individual application of different levels of pruning, organic manure and nutrient. Plant height and spread was maximum due to the interaction P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> (50% pruning intensity + 40 kg FYM + 100 % Recommended Dose of Fertilizer) followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. Among flowering, fruiting and yield parameters, maximum duration of flowering, fruit set, retention and yield and minimum fruit drop and days taken to fruit maturity was found under P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. Interaction P<sub>2</sub>M<sub>2</sub>N<sub>2</sub> has been noticed statistically superior to enhance physical and chemical characteristics of fruit followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub>. This treatment has not only enhanced bearing and quality parameters in aonla but also increased the leaf nutrient status as well as soil fertility status.

The analysis indicated that highest net return and cost benefit ratio was estimated in the plants that received 50% pruning intensity + 40 kg FYM + 100 % RDF followed by P<sub>2</sub>M<sub>2</sub>N<sub>1</sub> (50% pruning intensity + 40 kg FYM + 50 % RDF). Therefore, 50% pruning intensity along with nutrient management by use of organic manure (40 kg/tree FYM) and inorganic fertilizers (1000:500:1000 g/plant NPK) is recommended for sustainable higher fruit yield and quality of aonla while the interaction of 50% pruning intensity along with 40 kg/tree FYM and 50 % RDF is recommended for at par net return and cost benefit ratio with best treatment on less cost of production.

## REFERENCES

- Dalal S R, Patil S R and Dalal N R. 2000. Effect of severity of pruning on growth and quality of fruits of 25 years old guava cv. Sardar. *Journal of Soils and Crops* **10**: 298–300.
- Dhaliwal G S, Rattanpal H S and Gill H S. 2000. Effect of time and severity of pruning on cropping and physico-chemical properties of Sardar guava. *Haryana Journal of Horticultural Science* **29**: 27–20.
- Ghuge T D, Gore A K and Jadhav S B. 2007. Effect of organic and inorganic nutrient sources on growth, yield and quality of cabbage (*Brassica oleracea* var. capitata). *Journal of Soils and Crops* **17**: 89–92.
- Jaggi R C. 2007. Nutritional research in vegetables of Himachal Pradesh. *Indian Journal Fertilizer* **2**: 25–35.
- Naik M H and Babu R S H. 2005. Feasibility of organic farming in guava. *1<sup>st</sup> International Guava Symposium*, Central Institute for Subtropical Horticulture, Lucknow, p 69.
- Prabu T, Narwadkar P R, Sajindranath A K and Rathod N G. 2002. Integrated nutrient management in coriander. *South Indian Horticulture* **50**: 680–4.
- Ray R, Patra S K, Ghosh K K and Sahoo S K. 2005. Integrated nutrient management in okra (*Abelmoschus esculentus* L. Moench) in a river basin. *Indian Journal of Horticulture* **62**: 260–4.
- Santhy P, Muthuvel P and Selvi D. 2001. Status and impact of organic matter fractions on yield, uptake and available nutrients in a long term fertilizer experiment. *Journal of Indian Society of Soil Science* **49**: 281–5.
- Shaaban S H A, Soliman S S and Hamouda H A. 2006. Nutrient removal and physical, chemical properties in Samany date palm as influenced by different pruning treatments. *Journal of Applied Sciences Research* **2**(12): 1 142–50.
- Singh G B and Swaroop 2000. Lessons from long-term fertility experiments. *Fertilizer News* **45**: 13–8 and 21–4.
- Singh S K. 2005. Studies on pruning behaviour in Indian gooseberry (*Emblica officinalis* Gaertn.) cv. Narendra Aonla 7. M Sc (Ag) thesis in Horticulture, NDUAT, Faizabad.
- Swarup A and Wanjani R H. 2000. Three decades of all India coordinated research project on long-term fertilizer experiments to study changes in soil quality, crop productivity and sustainability. Indian Institute of Soil Science, Bhopal, p 59.
- Totawat K L, Somani L L, Singh R and Singh G. 2001. Integrated nitrogen management in maize-wheat cropping sequence on topic haplustalfts of western India. *Annals Arid Zone* **40**: 439–44.
- Tripathi P and Maity T K. 2007. Evaluation of summer okra (*Abelmoschus esculentus* L. Moench) hybrids under reduced level of chemical fertilizers. *Orissa Journal of Horticulture* **35**: 6–13.