



Performance evaluation of yield crop forecasting models using weather index regression analysis

SANJEEV PANWAR¹, K N SINGH², ANIL KUMAR³, RANJIT KUMAR PAUL⁴, SUSHEEL KUMAR SARKAR⁵,
BISHAL GURUNG⁶ and ABHISHEK RATHORE

Indian Council of Agricultural Research, Krishi Bhawan, New Delhi 110 012

Received: 8 October 2015; Accepted: 9 August 2016

ABSTRACT

A crop forecast is a statement of the most likely magnitude of yield or production of a crop. It is made on the basis of known facts on a given date and it assumes that the weather conditions and damages during the remainder of the growing season will be about the same as the average of previous year. The present paper deals with use of non-linear regression analysis for developing wheat yield forecast model for Allahabad district (India). A novel statistical approach attempted in this study to use nonlinear models with different weather variables and their indices and compare them to identify a suitable forecasting model. Time series yield data of 40 years (1970-2010) and weather data for the year 1970-71 to 2009-10 have been utilized. The models have been used to forecast yield in the subsequent three years 2008-09 to 2009-10 (which were not included in model development). The approach provided reliable yield forecast about two months before harvest.

Key words: Forecasting, Nonlinear regression model, Weather indices approach, Weather variables

Agricultural production is significantly affected by environmental factors. Weather influences crop growth and development, causing large intra-seasonal yield variability. In addition, spatial variability of soil properties, interacting with the weather, cause spatial yield variability. Crop forecast are the great value to all sections of the community. They serve as indicators of the economic condition of agriculture which is such an important portion of any country's economy.

Crop yield is influenced by technological change and weather variability. Technological factors increase yield smoothly through time and, therefore, years or some other parameters of time can be used to study the overall effect of technology on yield. Thus pre-harvest forecasting of production is required when crop is still standing in the field.

An efficient forecasting is thus a pre-requisite for food supply information system at district and state level. The final crop production estimates, though based on objective crop-cutting experiments, are of limited utility as these become available much later after the crop harvest. It is of high importance that with the advancements in non-linear estimation techniques and high computational ability to try for non-linear modeling of yield forecast using different

weather parameters and using their indices.

MATERIALS AND METHODS

In this paper application of non-linear regression technique has been made for modelling and forecasting yield of wheat crop for Allahabad district of Uttar Pradesh.

District level wheat crop yield data for 40 years (1970-2010) have been collected from the Directorate of Economics and Statistics, Ministry of Agriculture, Government of India, New Delhi and from the Agriculture Directorate, Lucknow (U.P.) and India Metrological Department, Pune and CRIDA.

Weather data on temperature (maximum and minimum), relative humidity and total rainfall from the year 1970-71 to 2009-10 have been utilized for model fitting and two years data 2008-09 and 2009-10 used for validation of the model.

Wheat is generally sown in the month of October when average daily temperature falls around 23°-25°C. The pre-sowing phase of the crop is important because in this phase of two to three weeks, the land is prepared for the cultivation. If the weather condition is adverse during the pre-sowing phase the sowing of the crop is generally delayed. After sowing of the crop, germination takes 6-7 days or near about one week after the pre-sowing phase. After germination phase, crown root initiation occurs after 20-25 days of sowing or in about 3 weeks from germination. Tillering phase starts just after the crown root initiation phase and lasts up to 40-45 days after sowing or nearly about 2-3 weeks after crown root initiation phase. Jointing and reproductive phase is the peak plant growth stage and

¹Senior Scientist (e mail: scientist1775@gmail.com), ICAR, Krishi bhawan, ^{2,3,4,5,6}Senior Scientist, Senior Scientist, Principal Scientist, Scientist, Senior Scientist, Scientist IASRI, New Delhi 110 012. Senior Scientist, ICRIASAT, Hyderabad.

starts after the tillering phase or 45-60 days after sowing. The reproductive phase lasts 60-85 days after sowing.

As weather during pre-sowing period is important for establishment of the crop, data starting from two weeks before sowing have been included in model development. Further, as the forecast is required well in advance of harvest, weather data about two months before harvesting have been considered. Thus data on four variables viz., max. temp., min. temp., RH, rainfall during 16 weeks data from 40th SMW to 3rd SMW (next year).

Crops yield forecast models have been developed for districts of Allahabad of U.P. state using weekly weather data. Residuals obtained from the selected non-linear models and linear models. For that, trend analysis has been done through linear and non-linear approaches. In which for each weather variable two indices were developed, one as simple total of values of weather parameter in different weeks and the other one as weighted total, weights being correlation coefficients between detrended yield and weather variable in respective weeks. Weather indices based regression models were developed using weather indices as independent variables while detrended yield (residuals) was considered as dependent variable.

Weather Indices (WI) were obtained using nonlinear and linear modelling approaches. Weather indices based regression models were developed using weather indices as independent variables while character under study such as crop yield was used as dependent variable for Wheat crop. Step-wise regression technique used for selecting significant variables in all the models.

Statistical approaches have been used for development forecast model based on weather variables, which are as follows:

Following nonlinear growth models will provide a reasonable representation of yields which are expected to provide a reasonable representation of crop yield as compared to linear models will be tried for fitting the yield of selected crop in a selected location using the weather data

Logistic model:

$$Y = \frac{\alpha}{1 + \exp(b - ct)} + e$$

Gompertz Model:

$$Y = a \exp [-\exp (b - c)] + e$$

Richards Model:

$$Y = \frac{\alpha}{1 + \exp(b - ct)^{1/d}} + e$$

Wiebull Model:

$$Y = a - b \exp (-ct^d) + e$$

where, a, b, c and d are the unknown parameters to be estimated.

For nonlinear estimation procedures, Gauss-Newton Method is used which yields efficient estimates with proper convergence was applied in this study.

The residuals from above model will be used in the subsequent linear model for fitting against different forms of the weather variables including their indices. The weather variables within an agricultural year will be aggregated through mean or through indices as mentioned above. Further, indices will be computed based on the influence (positive or negative) of the selected weather variable on the crop yield.

For each weather variable, two indices will be developed, one as simple total of values of weather variables parameter in different weeks and the other one as weighted total, weights being correlation coefficients between detrended yield and weather variable in respective weeks. The first index represents the total amount of weather parameter received by the crop during the period under consideration. While the other one takes care of distribution of weather parameters with special reference to its importance in different weeks in relation to the detrended yield. On similar line, indices were computed with products of weather variables (taken two at a time) for joint effects.

These indices are computed as follows:

$$y = a_0 + \sum_{i=1}^p \sum_{j=0}^1 a_{ij} Z_{ij} + \sum_{i \neq j=1}^p \sum_{j=0}^1 b_{ij} Z_{ij} + \epsilon$$

$$Z_{ij} = \sum_{w=n_1}^{n_2} r_{iw}^j X_{iw}$$

$$Z_{iij} = \sum_{w=n_1}^{n_2} r_{ii}^j X_{iw} X_{i'w}$$

where, y is detrended yield forecast, X_{iw} is value of ith weather variable in wth week, r_{iw} is correlation coefficient between Y and ith weather variable in wth week, $X_{i'w}$ is correlation coefficient between Y and product of X_{iw} and $X_{i'w}$ in wth week, .p is number of weather variables, n_1 is initial week for which weather data are included in the model, n_2 is final week for which weather data are included in the model and a_{ij} & b_{ij} are parameters to be estimated ϵ is the random error

Stepwise regression technique was used for retaining significant variables only in the forecast models in each approach.

Comparison and validation of models

Different regression models were compared on the basis of adjusted efficient of determination (R_{adj}^2) which is as follows:

$$R_{adj}^2 = 1 - \frac{ss_{res} / (n - p)}{ss_t / (n - 1)}$$

where $ss_{res} / (n - p)$ is the residual mean square and $ss_t / (n - 1)$ is the total mean square.

From the fitted models, wheat yield forecasts for the years 2008-09 to 2009-10 were obtained and forecasts were compared with the actual yield on the basis of Root Mean Square Deviation (RMSE).

$$RMSE = \left[\frac{1}{n} \sum_{i=1}^n (O_i - E_i)^2 \right]^{\frac{1}{2}}$$

Table 1 Allahabad wheat yield forecast models (Two Steps non-linear & linear models) based on weather indices approach

District name	Forecasting Model	Goodness of fit	
		AdjR ²	RMSE
Allahabad Non-linear	$Y_t = -7.83 + 0.027 Z_{11} - 1.52 Z_{31} + 0.032 Z_{141} - 0.032 Z_{231}$ (0.014) (0.098) (0.0010) (0.01)	0.93	1.79
Linear	$Y_t = 7.44 + 0.038 Z_{11} - 1.56 Z_{31} + 0.0032 Z_{141} - 0.037 Z_{231}$ + 0.01905 Z ₃₄₁ (0.01) (0.16) (0.001) (0.003) (0.001)	0.92	1.91

Table 2 MAPE of forecast of wheat yield from models developed

District	Two step nonlinear models	Linear models
Allahabad	7.16	8.17

where O_i and the E_i are the observed and forecast value of crop yield respectively and n is the number of years for which forecasting has been done.

RESULTS AND DISCUSSION

The two step nonlinear forecasting model was found to be superior or at far to the linear model as its RMSE value (1.79) is much lower as compared that of linear model (1.91). For fitting this residual model the nonlinear model (Logistic) which was found to have better fit was used to output the residuals. The negative values of the coefficients for variables such as Z₃₁ (rainfall) and Z₂₃₁ (min temp*rainfall), showed that increase in the variables results in decrease in the yield. On the other hand variable such as Z₁₁ (max temp) and Z₁₄₁ (max temp*relative humidity) yielded positive coefficients which means that increase in the variables will increase the yield of Wheat crop in Allahabad district of Uttar Pradesh. Similarly, RMSE values are much lower as compared that of linear model in Allahabad districts of UP, thereby showing that

two step non-linear forecasting models were found to be superior or at par for yield forecasting of wheat crop. The models provides reliable forecast around two months before harvest (Table 2).

REFERENCES

- Agrawal Ranjana, Jain R C, Jha M P and Singh D. 1980. Forecasting of wheat yield using climatic variables. *Ind. J. Agric. Sci.*, **50**(9) pp 680–4.
- Bates D M and Watts D G. 1988. *Nonlinear regression analysis and its App. Met.*, **12**(6), pp 937–47.
- Baier W. 1977. Crop weather models and their use in yield assessments. *Tech. note no. 151*, WMO, Geneva, 48 pp.
- Carter M M and Eisner J B. 1997. A Statistical Method for Forecasting Rainfall over Puerto Rico. Department of Meteorology, The Florida State University, U.S.A.
- Frere M and Popov G F. 1979. Agrometeorological crop monitoring and forecasting. FAO plant production and protection paper, *Plant Production and Protection Division*, FAO.
- Panwar Sanjeev, Kumar Anil, Singh K N, Farooqi Md Samir, Rathore Abhishek. 2012. Using Nonlinear Regression Techniques for analysis of Onion Production in India. *Indian Journal of Agriculture Science* Vol **82** (12) pp 43–6, 2012.
- Seber G A F and Wild C J. 1989. Non-linear regression. John Wiley and Sons, New York.
- Panwar Sanjeev, Sharma S C and Kumar Anil. 2009. A critical study of Onion (*Allium cepa*) export of India: Non linear Approach” *Indian Journal of Agriculture Science*. Vol **78**, no.2, pp 178–80.