



Effect of girdling on flowering and fruiting of litchi (*Litchi chinensis*) in subtropics of Bihar

AMRENDRA KUMAR¹, S D PANDEY², R K PATEL³, KULDEEP SRIVASTAVA⁴, MRITUNJAY TRIPATHI⁵ and VISHAL NATH⁶

ICAR-National Research Centre on Litchi, Muzaffarpur, Bihar 842 002

Received: 22 July 2016; Accepted: 12 September 2016

ABSTRACT

An experiment was conducted during 2014 and 2015 to study the effect of varied extent of girdling for induction of flowering in litchi (*Litchi chinensis* sonn.) under Bihar condition. Results indicated that circular girdling of 2mm on 50% primary branches produced more flowering panicle, fruit yield, fruit size, TSS, and TSS/acid ratio in Shahi litchi trees compared to ungirdle (control) trees. Findings also revealed that the wound healing duration of girdled portion increases with increasing the size of girdling notch. The average yields from all girdled treatments were about three to five times higher than control. Gradual decrease in fruit weight was noticed with increase of girdling size in both 25% PB and 50% PB. TSS and TSS/acid ratio showed decreasing trend while acidity and ascorbic acid content of the fruit showed increasing trend with increase in level of girdling width. All girdled shoots showed more accumulation of carbohydrate in leaves than control. Among the girdled branches, gradual increase in leaf carbohydrate and C/N ratio content and decrease in nitrogen content observed with increase in level of girdling but the nitrogen content of bark showed the reverse trend.

Key words: Biochemical parameters, Flowering, Girdling, Litchi, Primary branches

Dormancy in litchi (*Litchi chinensis* Sonn.) is induced by dry weather or recurring periods of relatively cold weather or both, during late fall and early winter (Young 1956, 1977, Menzel and Simpson 1987). Litchi requires dry and cold weather in winter to induce flowering and hot and humid weather during rest of the year. Alternate bearing is a characteristic of some litchi varieties, which is more pronounced when grown in warm climates. According to Garcia-Perez and Martins (2006), the litchi requires seasonal temperature variations for best flowering and fruiting. Most varieties need a cold period of 100 to 200 hours between 0° and 7°C, preferably with low rainfall. There is evidence that lower night temperature below 15°C during autumn months favour floral induction, and high day temperature at the same period reduce low night temperature effectiveness (Menzel and Simpson 1995). Litchi production in hot and subtropical climates is hampered by low flowering. There are technologies which has shown as promise to induce flowering amongst branch girdling is one. Worldwide

irregular production in litchi is associated with sparse flowering and low fruit fixation. Production alternation characteristics in different cultivars and climate order restriction are the main causes related to flowering problems in litchi (Ghosh 2001). An alternative approach to mitigate the low flowering problem in litchi on primary or secondary branch. It has been used in several litchi producing countries. The technique involves making an incision from 1.6 to 4.0 mm width, usually with the help of a pruning saw, in whole branch circumference (Carvalho and Salomao 2000). In this operation, a portion of bark (phloem) is removed from the wood by a technique known as girdling. Since the woody xylem part remains intact, water and nutrient reaches the leaves. After the preparation of photosynthate, it is not transported to other parts below the girdle and accumulation of photosynthate just above the girdle region takes place which in turn improves C:N ratio and there by flowering. A number of studies have been conducted to determine the influence of carbohydrate availability on the development of inflorescence and they have been found to alter carbohydrate availability by reducing the leaf area by introducing a girdle on a developing shoot before the onset of dormancy (Candolfi-vasconcelos and Koblet 1990, Caspari *et al.* 1998, Bennett *et al.* 2005, Sanchez and Dokoozlian 2008, Smith and Holzapfel 2009, Vrsic *et al.* 2009). In litchi, efforts have been made to develop an understanding of the assimilate partitioning behavior with

^{1,2}Principal Scientist, Horticulture (e mail: amrendra14d@gmail.com, pandeynrbc@yahoo.com), ³Senior Scientist, Horticulture (e mail: rkpatelicar@gmail.com), ⁴Senior Scientist, Entomology (e mail: kuldeep.ipm@gmail.com), ⁵Young Professional (e mail: mirtunjay.biochem@gmail.com), ⁶Director, (e mail: vishalnath66@rediffmail.com).

respect to fruit growth and yield. In such studies, girdling used as an effective method for improving the yield by proper source-sink utilization in perennial woody plants (Kumar *et al.* 2015). In view of above facts present investigation was conducted with objective to suppress vegetative growth prior to flowering in litchi cv. Shahi under subtropical climatic condition of Bihar (India).

MATERIALS AND METHODS

A field experiment was conducted during 2014 and 2015 for two consecutive years at ICAR- National Research Centre on Litchi, Mushahari, Muzaffarpur (Bihar) located at an altitude of 210 m asl. The litchi cv. Shahi planted in the year 2005 at a spacing of 8.25 × 8.25 m were used as a study material. The soil of experimental field was sandy loam in texture, alkaline in reaction with medium in fertility status. The experiment was laid out in randomized block design (RBD) with three replication comprising of 7 treatments, viz T1: Girdling 2mm 25% Primary branches (PB), T2: 4mm 25% PB, T3: 6mm 25% PB, T4: 2mm 50% PB, T5: 4mm 50% PB, T6: 6mm 50% PB, T7: ungirdle (control). The girdling was performed during last week of August after appearance of second vegetative flushes. A strip of bark was removed as per treatment in a circular fashion around the selected primary branches. Subsequent girdling during next year was performed 4-5 cm above the previous girdled. The standard cultural practices were followed as per the recommendation of Kumar *et al.* (2014) and Srivastava *et al.* (2015).

The observation on per cent flowering shoot was recorded by tagging the 100 shoots in each direction of the trees. The duration of flowering was counted from the date of girdling operation performed. Wound healing period of girdled portion was recorded from the date of girdling and when phloem of wounded portion joined completely. Fruit maturity (harvest date) was determined on the basis of standard maturity indices, i.e. fruit colour (bright pinkish red blush) tubercles (flattened tubercles) and pulp TSS (18°Brix). Fruit length and diameter were measured from 10 randomly selected fruits with digital vernier caliper. The weight of the fruits was calculated on the basis of 10 representative fruits and the mean was expressed in gram (g). The total soluble solids (TSS) content was determined with Erma Hand Refractometer (0-32°Brix). The titratable acidity and ascorbic acid content of juice were estimated as per AOAC (1995). TSS/acid ratio was derived from the TSS and acidity values. Nitrogen content in plant samples was estimated according to modified Kjeldahl procedure (Jackson 1973) using Pelican Kel Plus Auto analyzer. Carbohydrate content of leaf and bark was analyzed with Anthrone reagent method (Hedge and Hofreiter 1962). Chlorophyll was extracted in 80% acetone and the absorption at 663, 652 and 645 nm were recorded with spectrophotometer (Witham *et al.* 1971). The data were subjected to statistical analysis as per the method of Gomez and Gomez (1984). Least significant of difference at 5% level was used for finding the significance of differences among the treatment means.

Table 1 Effect of girdling on flower induction, wound healing and fruit retention in litchi cv Shahi

Treatment	Flowering induced (%)			Days to flowering after girdling			Wound healing (days)			Fruits/panicle (Early stage)			Fruit retention (%)			Appearance of girdle portion
	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled	
	T ₁	76.67	80.60	78.64	158	161	159.50	99.67	118.07	108.87	38.38	34.53	36.46	34.24	44.62	
T ₂	83.33	82.45	82.89	164	168	166.00	153.67	167.33	160.50	27.60	24.60	26.10	44.35	49.23	46.79	Less swelling
T ₃	73.33	78.50	75.92	171	176	173.50	176.33	201.67	189.00	21.20	21.73	21.47	57.07	56.74	56.91	More swelling
T ₄	76.67	73.33	75.00	160	162	161.00	102.67	136.33	119.50	49.43	44.93	47.18	28.69	35.92	32.31	Smooth
T ₅	83.33	86.33	84.83	165	169	167.00	161.00	180.00	170.50	28.07	31.07	29.57	39.82	42.61	41.22	Less swelling
T ₆	74.50	78.33	76.42	170	176	173.00	179.67	211.67	195.67	27.67	27.53	27.60	40.26	44.93	42.60	More swelling
T ₇	41.00	30.33	35.67	153	155	154.00				28.33	26.67	27.50	34.06	42.48	38.27	
CD (P=0.05)	9.83	18.09	10.65	8.32	8.68	8.41				9.51	NS	8.67	3.43	3.65	3.48	

RESULTS AND DISCUSSION

The data (Table 1) revealed that girdling treatments significantly induced the flowering in the litchi during both the years as compared to control. Within treatments no significant differences were observed. Pooled means over two years showed that the highest flowering (84.83%) was observed in T₅ (4 mm and 50% PB) followed by T₂ (4 mm and 25% PB) 82.89% and minimum in un-girdled tree (35.67%) (Table 1). Girdled branches showed delay to bloom during both the years and panicles remains shorter than control branches. The substantial increase in flowering by girdling was also reported in mango (Gaskins 1963), apple (Samad 1998), jamun (Haldalkar *et al.* 2014). More flowering in girdled litchi branch might be due to suppression of growth flushes which induced the shoot maturity and accumulation of more photosynthates on girdled portion by disruption in sugar transport in phloem, stimulated flowering. Earliest flowering was noticed in ungirdled tree (control) and took 154 days to flower after girdling however T₃ and T₆ (6 mm groove) plants flowered after 173 days. The wound healing duration increases with increase in size of girdling during both the years. The early healing of wound was observed in 2 mm size (108.87 and 119.50 days) and branch above as well as below the girdled portion become uniform. While, girdling with 6 mm size took maximum duration (195.67 and 189.0 days) followed by 4mm size (170.50 and 165.50 days) for healing of wound resulted in

unequal growth of the primary limbs became more thicker branch size above the girdled portion.

Pooled analysis on number of fruit/panicle at initial and harvest stage revealed that girdling of branches do not have any significant difference except that of T₄ (2 mm and 50%PB) where highest fruit per panicle at initial stage was 47.18. Fruit retention at harvest stage showed the increasing trend with increase in girdling size during both the years. The highest (56.91%) fruit retention was recorded in T₃ (6 mm) and lowest (32.31%) in T₂ (2 mm). The fruit retention capacity in a panicle is a function of the strength of source tissue to support the carbohydrate demand of the growing sink (fruits). More fruit retention due to girdling can be attributed to retention of translocation of current photosynthates to other plant parts and its distribution to the developing fruits.

The average yield during both the year from all girdled treatments was about three to five times higher than that from control. The significantly higher fruit yield was noticed in T₄ (33.68 kg/plant) followed by T₅ (30.41 kg) and T₆ (26.34 kg) where 50% primary branches were girdled in comparison to 25% girdled primary branches (Table 3). Among 25% girdled PB treatment, T₁ (2 mm) was found better in respect of fruit yield. The girdling was found to increase fruit yield in wax apple (Minch and Chung 2012), apple (Samad 1998) and litchi (Li and Xiao 2001).

Significantly higher fruit weight and fruit size were

Table 2 Effect of girdling on yield and yield attributes in litchi cv. Shahi

Treatments	Yield (kg/plant)			Fruit weight (g)			Fruit length (mm)			Fruit breadth (mm)		
	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled
T ₁	18.40	19.20	18.80	20.45	21.05	20.75	35.10	34.33	34.72	30.00	29.67	29.84
T ₂	17.90	17.40	17.65	18.91	19.58	19.25	33.00	32.67	32.84	29.20	28.67	28.94
T ₃	16.25	17.37	16.81	16.80	19.16	17.98	31.60	32.00	31.80	27.70	27.67	27.69
T ₄	32.66	34.70	33.68	19.46	20.04	19.75	34.50	34.00	34.25	29.80	30.33	30.07
T ₅	29.65	31.17	30.41	18.05	18.87	18.46	32.80	33.00	32.90	28.40	28.33	28.37
T ₆	27.64	25.03	26.34	15.63	18.20	16.92	31.20	31.00	31.10	26.90	27.00	26.95
T ₇	9.21	4.23	6.72	15.09	17.81	16.45	26.70	28.00	27.35	25.70	27.67	26.69
CD(P=0.05)	2.92	2.72	3.12	1.17	0.37	1.09	0.23	0.38	0.34	0.21	0.47	0.39

Table 3 Effect of girdling on quality parameter in litchi cv. Shahi

Treatment	TSS (°brix)			Acidity (%)			TSS/acid ratio			Ascorbic acid (mg/100gpulp)		
	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled	2014	2015	Pooled
T ₁	18.84	19.23	19.04	0.68	0.66	0.67	27.71	29.14	28.43	28.35	27.63	27.99
T ₂	18.80	19.20	19.00	0.76	0.74	0.75	24.74	25.95	25.35	32.55	30.50	31.53
T ₃	18.61	18.50	18.56	0.77	0.81	0.79	24.17	22.84	23.51	37.61	34.62	36.12
T ₄	20.50	20.17	20.34	0.64	0.73	0.69	32.03	27.63	29.83	30.33	28.80	29.57
T ₅	19.69	19.30	19.50	0.66	0.78	0.72	29.83	24.74	27.29	37.24	34.46	35.85
T ₆	19.27	19.27	19.27	0.72	0.80	0.76	26.76	24.09	25.43	39.15	36.42	37.79
T ₇	17.67	18.83	18.25	0.84	0.82	0.83	21.04	22.96	22.00	30.50	27.58	29.04
CD(P=0.05)	0.82	0.19	0.62	0.19	0.14	0.16	7.22	6.98	7.12	6.36	4.59	5.45

observed in girdled tree than control during both the years (Table 3). It appears that the girdled branches supplied more carbohydrate reserve and necessary hormones which were essential to increase the fruit weight. Gradual decrease in fruit weight was noticed with increase of girdling size in both 25% PB and 50% PB. Girdling of 2mm size produced extra class fruit (fruit length > 33 mm) with fruit weight (20.75g in 25% PB and 19.75g in 50% PB) and fruit size (34.72 mm and 34.25 mm) than other girdling treatments. Carbohydrate accumulation and canopy photosynthesis is essential for fruit developments. Partitioning of available assimilates to total number of fruits have resulted into increase in fruit weight even under the condition of source limitation. Lower fruit weight in ungirdled tree might be due to less accumulation of carbohydrate reserves for developing fruit which might have diverted to other plant parts like shoots, roots and non-fruiting terminals that subsequently lead to new flushing during the active phase of fruit growth. Production of extra class fruit in 2mm girdling size might be due to proper utilization of photosynthates and necessary hormones reserve besides complete wound healing of girdled portion during fruit growth period resulted into re-establishment of sap flow. Gradual decrease in fruit weight with increase in girdling size might be associated with delay in wound healing resulting into improper establishment of sap flow which hampered the proper regulation of carbohydrate and nitrogen reserve during fruit growth (Khalkho *et al.* 2015)

TSS and TSS/ acid ratio showed decreasing trend while acidity and ascorbic acid content of the fruit showed increasing trend with increase level of girdling width. TSS ranged from 18.25 to 20.34 °brix and TSS/ acid ratio from 22.96 to 29.14 among treatments. The lowest acidity (0.67%) was also noticed in T₁ and highest under control (0.83%). The highest ascorbic acid content in fruit was found in T₇ (37.79 mg/100g) followed by T₃ (36.12mg/100g), T₅ (35.85mg/100g) and lowest in T₁ (27.99 mg/100g). All the girdled treatment showed more accumulation of carbohydrate in leaves than control (Fig 1). Among the girdled treatment, gradual increase in leaf carbohydrate and C/N ratio and decrease in nitrogen content was observed with increased level of girdling. However, the nitrogen content of bark showed the reverse trend. The highest carbohydrate

content in leaves was recorded in T₃ (5.36%) which is closely followed by T₆ (5.31%) and lowest in control (3.53%). The carbohydrate content in bark of girdled branches varied from 4.21% (T₁) to 5.07% (T₄). Girdling stops the basipetal movement of assimilates through phloem which resulted in accumulation of carbohydrates and plant growth hormones above the girdle (Urban and Alphonsout 2006). Transitory block phloem flux apparently increases sugar availability in the aerial part of tree which induced flowering (Wallerstein *et al.* 1978 and Yamanishi 1995). Leaf nitrogen content (1.38%) and C/N ratio (2.56) in ungirdled tree apparently found to be lower than girdled treatment (Fig 2 and 3). The highest leaf N was observed in T₁ (1.86%) followed by T₂ (1.84%) and T₄ (1.81%) whereas the C/N ratio was found highest in T₆ (4.15) and T₃ (4.12). Chlorophyll (chl) content (a, b and total) of matured leaves showed decreasing pattern with level of increasing in girdle size. The highest chl a, chl b, chl total content was recorded in treatment with 2mm girdling in 25% PB (1.84, 0.78 and 2.66 mg/g) followed by 2mm girdling in 50% PB (1.73 0.77 and 2.59 mg/g) (Fig 4). All the girdled treatment recorded lower value of chl a/b ratio (2.00 to 2.36) than ungirdled treatment (4.06). Higher chlorophyll content in thin groove size girdling branches might be due early healing of injured portion which regulated the leaf photosynthetic activity and resulted in retardation of the leaf senescence by restricting/modifying the changes in chlorophyll.

On the basis of present finding it may be concluded that circular girdling of 2mm on 50% primary branches of




Fig 1 Effect of girdling on carbohydrate content in leaf and bark of litchi cv. Shahi




Fig 2 Effect of girdling on nitrogen content in leaf and bark of litchi cv. Shahi




Fig 3 Effect of girdling on C/N ratio in leaf and bark of litchi cv. Shahi

Shahi litchi produced more flowering panicle, fruit yield, fruit size, TSS, and TSS/acid ratio.

REFERENCES

- AOAC. 1995. *Official Methods of Analysis*, 16th edn. The Association of Official Analytical Chemists, Washington DC.
- Bennett J, Jarvis P, Creasy G L and Trought M C T. 2005. Influence of defoliation on overwintering carbohydrate reserves, return bloom, and yield of mature Chardonnay grapevines. *American Journal of Enology and Viticulture* **56**: 386–93.
- Candolfi-vasconcelos C C and Koblet W. 1990. Yield, fruit quality, bud fertility and starch reserves of the wood as a function of leaf removal in *Vitis vinifera*- Evidence of compensation and stress recovering. *Vitis* **29**: 199–21.
- Carvalho C M and Salomao CC. 2000. Culture of litchi. Federal University of Viçosa, Viçosa; 38 p.
- Caspari H W, Lang A and Alspach P. 1998. Effects of girdling and leaf removal on fruit set and vegetative growth in grape. *American Journal of Enology and Viticulture* **49**: 359–66.
- García-Pérez E and Martins A B G. 2006. Flowering and fruiting lychee trees due to the girdling branches. *Journal of Fruit Growing* **28**(1): 14–7.
- Gaskin M H. 1963. Girdling in mango seedlings for inducing early fruit bearing. *Florida State Horticultural Society*: 360–3.
- Ghosh S P. 2001. World trade in litchi: past, present and future. *Acta Horticulturae* **558**(34): 233–5.
- Gomez A A and Gomez K A. 1984. *Statistical Procedures for Agricultural Research*, p 680. John Wiley and Sons Inc, New York.
- Haldalkar P M, Dalvi, N V, Parulekar Y R and Lawande K E. 2014. Induction of flowering by girdling in Jamun cv. Konkan Bahadoli. *Journal of Plant Studies* **3**(2): 45–9.
- Hedge J E and Hofreiter B T. 1962. *Carbohydrate Chemistry*, 17. Whistler R L and Be Miller J N (Eds), Academic Press, New York.
- Jackson M L R. 1973. Soil chemical analysis - advanced course: A manual of methods useful for instruction and research in soil chemistry, physical chemistry of soils, soil fertility, and soil genesis. Madison, Wis.
- Khalkho N S, Horo P and Jha K K. 2015. Effect of girdling and defoliation on physical properties of litchi fruit. *Journal of Crop and Weed* **11**(1): 200–3.
- Kumar A, Pandey S D, Patel R K, Singh S K, Srivastava K and Nath V. 2015. Induction of flowering by use of chemicals and cincturing in Shahi litchi. *Ecoscan Special issue VII*: 493–96.
- Kumar A, Srivastava K, Patel R K and Nath V. 2014. Management of litchi fruit borer and litchi mite using bio-rational approaches under subtropics of Bihar. *Ecoscan Special issue VI*: 285–9.
- Li C B and Xiao Y. 2001. Girdling increases yield of Nuomici litchi. (In) *International Symposium on litchi, longan and other sapindaceae fruit*. Huang H and Menzel C (Eds). *Acta Horticulturae* **558**: 233.
- Menzel C M and Simpson D R. 1986. Effects of cincturing on growth and flowering of litchi: preliminary observations in subtropical Queensland. *Australian Journal of Experimental Agriculture* **26**: 255–9.
- Menzel C M and Simpson D R. 1987. Effects of cincturing on growth and flowering of litchi over several seasons in subtropical Queensland. *Australian Journal of Experimental Agriculture* **27**: 733–38.
- Menzel C M and Simpson D R. 1995. Temperatures above 20^oC reduce flowering in lychee (*Litchi chinensis* Sonn.). *Journal of Horticultural Science and Biotechnology* **70**(6): 981–8.
- Minch T N and Chung R Y. 2012. Effect of S- girdling on fruit growth and fruit quality of wax apple. *World Academy of Science, Engineering and Technology* **72**: 216.
- Samad A. 1998. Effect of girdling and bridge grafting on apple tree growth, yield, apple quality and biochemical composition for apple trees growing under moderate and ultra high density conditions. Ph D thesis, Lincoln University.
- Sanchez L and Dokoozlian N. 2008. Bud carbohydrate status and fruitfulness of *Vitis vinifera* L. *Hort Science* **43**: 1 145–8.
- Smith J P and Holzapel B P. 2009. Cumulative responses of Semillon grapevines to late season perturbation of carbohydrate reserve status. *American Journal of Enology and Viticulture* **60**: 461–70.
- Srivastava K, Patel R K, Pandey S D, Kumar A and Nath V. 2015. Integrated management of lepidopteran defoliators in litchi under subtropics of Bihar. *Ecoscan VII*: 483–90.
- Urban L and Alphonsout L. 2006. Girdling decreases photosynthetic electron fluxes and induce sustained photoprotection in mango leaves. *Tree Physiology* **27**: 345–52.
- Vrsic S, Pulko B and Valdhuber J. 2009. Influence of defoliation on carbohydrate reserves of young grapevines in the nursery. *European Journal of Horticulture Science* **74**: 218–22.
- Wallerstein I, Goren R and Ben-Tal Y. 1978. Effect of ringing on root starvation in sour orange seedling. *Journal of Horticultural Science* **53**: 109–13.
- Witham F H, Blaydes B F and Devlin R M. 1971. *Experiments in Plant Physiology*, pp 167–200. Van Nostrand Reinhold, New York, USA.
- Yamanishi O K. 1995. Trunk strangulation and winter heating effects on fruit size, internal quality and maturation of Tosa Buntan grown in plastic house. *Journal of Horticultural Science* **69**: 305–8.
- Young T W. 1956. Response of lychees to girdling. (In) *Proceedings of Florida State Horticulture Society* **69**: 305–8.
- Young T W. 1977. Effect of branch girdling on yield of severely pruned 'Brewster lychee trees'. *Proceedings of Florida State Horticulture Society* **90**: 251–3.



Contribution of subterranean clover (*Trifolium subterraneum*) to changes in morphological and physiological parameters raised alone and with birdsfoot trefoil (*Lotus corniculatus*)

VILIANA VASILEVA¹ and ANNA ILIEVA²

Institute of Forage Crops, 89 General Vladimir Vazov Str, Pleven 5800, Bulgaria

Received: 12 August 2016; Accepted: 30 September 2016

ABSTRACT

Changes in some morphological and physiological parameters (leaves/stems ratio, total plastid pigments content, fixed amount nitrogen) of birdsfoot trefoil (*Lotus corniculatus* L.) (cv. Targovishte 1) and three subterranean clover subspecies, i.e. *Trifolium subterraneum* ssp. *brachycalicinum* (cv. Antas), *Trifolium subterraneum* ssp. *yananicum* (cv. Trikkala) and *Trifolium subterraneum* ssp. *subterraneum* (cv. Denmark) were studied in a field trial in the Institute of Forage Crops, Pleven, Bulgaria (2012-15). Crops were grown alone (100%) and in mixtures as follows: birdsfoot trefoil + *Trifolium subterraneum* ssp. *brachycalicinum* (50:50%); birdsfoot trefoil + *Trifolium subterraneum* ssp. *yananicum* (50:50%); birdsfoot trefoil + *Trifolium subterraneum* ssp. *subterraneum* (50:50%). Leaves/stems ratio for alone grown crops was found: birdsfoot trefoil (1.33), *Trifolium subterraneum* ssp. *brachycalicinum* (1.02), *Trifolium subterraneum* ssp. *yananicum* (0.93) and *Trifolium subterraneum* ssp. *subterraneum* (0.93). Leaves/stems ratio for birdsfoot trefoil in all mixtures was found to be higher as compared to alone grown birdsfoot trefoil - by 13.0% for the mixture with *Trifolium subterraneum* ssp. *yananicum*, by 16.3% for that with *Trifolium subterraneum* ssp. *brachycalicinum* and by 20.1% for the mixture with *Trifolium subterraneum* ssp. *subterraneum*. Total plastid pigments in birdsfoot trefoil increased in all mixtures: with *Trifolium subterraneum* ssp. *brachycalicinum* by 7.4%, with *Trifolium subterraneum* ssp. *subterraneum* by 8.1% and with *Trifolium subterraneum* ssp. *yananicum* by 15.6%. From mixtures of birdsfoot trefoil were obtained more fixed amount of nitrogen - from 1.71 kg N/da (with *Trifolium subterraneum* ssp. *subterraneum*) to 4.84 kg N/da (with *Trifolium subterraneum* ssp. *brachycalicinum*) as compared to alone grown birdsfoot trefoil. Birdsfoot trefoil and subterranean clover in mixtures showed good tolerance, morphological and physiological status.

Key words: Birdsfoot trefoil, Leaves stems ratio, Mixtures, Plastid pigments, Subterranean clover

Interest towards intercropped systems is rising in recent years due to their role in formation of a system for sustainable agriculture (Luscher *et al.* 2014, Kusvuran *et al.* 2014). They are more effective than pure grown in using environmental resources, better withstand adverse conditions and are more productive (Porqueddu *et al.* 2003, Chourkova 2014). The components in mixtures go into competitive relations and many morphological and physiological parameters change. The leaves/stems ratio is an important morphological characteristic and indicator for quality and uptake of the forage (Buxton *et al.* 1985). From the physiological status of components depends on the productivity of mixtures.

Due to the nitrogen fixation in mixtures with participation of legumes, the nitrogen content increased and nitrogen obtained in this way is used directly by the plants (Peeters *et al.* 2006). In the present study birdsfoot trefoil is included - traditional nitrogen fixing forage crop that is

accepted by all kinds of animals and does not cause swelling under grazing (Vučković 2004, Stevanovic *et al.* 2015).

Subterranean clover (*Trifolium subterraneum* L.) is a widespread component in the pastures and other grasslands of the temperate areas of Central and Northern Europe and America (Nichols *et al.* 2012). Investigations on subterranean clover as a component of sown pasture in recent years have shown that it has practical application for local climatic conditions in Bulgaria (Vasilev 2006, Vasileva and Vasilev 2012, Ilieva *et al.* 2015).

The aim of this work was to study changes in some morphological and physiological parameters - leaves/stems ratio, total plastid pigments content (chlorophylls and carotenoids) and amount of fixed nitrogen in the plants of birdsfoot trefoil and subterranean clover grown pure and in mixtures with direction of use for forage.

MATERIALS AND METHODS

The trial was conducted at the experimental field (43° 23'N, 24° 34'E, 230 m altitude) of the Institute of Forage Crops, Pleven, Bulgaria (2012-15) on podzolized soil

¹e mail: viliana.vasileva@gmail.com, ²e mail:anna_ibg@yahoo.com.

subtype without irrigation. Long plot method was used with size of plots 70 m² (1.40 × 50 m) four replicated. The object of study were birdsfoot trefoil (*Lotus corniculatus* L.) cv. Targovishte 1 and three subclover subspecies, i.e. *Trifolium subterraneum* ssp. *brachycalycinum* (cv. Antas), *Trifolium subterraneum* ssp. *yananicum* (cv. Trikkala) and *Trifolium subterraneum* ssp. *subterraneum* (cv. Denmark), grown pure (100%) and in mixtures in ratios birdsfoot trefoil: subterranean clover (50:50%), as follows: birdsfoot trefoil + *Trifolium subterraneum* ssp. *brachycalycinum* (50:50%); birdsfoot trefoil + *Trifolium subterraneum* ssp. *yananicum* (50:50%); birdsfoot trefoil + *Trifolium subterraneum* ssp. *subterraneum* (50:50%). The sowing was done in autumn (beginning of October) of 2012 with sowing rates for birdsfoot trefoil 1.5 kg/da and for subterranean clover 2.5 kg/da, between row space was 11.5 cm. Sowing rate in mixtures was 1/2 from this for pure swards. No fertilizers and pesticides were applied during the vegetation.

The swards were harvested for forage. Two cuts were obtained during the first and second year after sowing (2013 and 2014) and one cut during the third (2015). At each harvest, samples were taken from an area of 0.25 m², divided are the leaves and stems, and weighted separately (g fresh weight). The leaves/stems ratio was calculated (individually and in mixtures).

In fresh plant samples during the three years total plastid pigments content (chlorophyll a, chlorophyll b, and carotenoids) (mg/100 g FW) was determined according Zelenskii and Mogileva (1980). For the mixtures samples were taken from every component.

The Carlsson and Huss-Danell (2003) formulae for roughly determine the amount of fixed nitrogen in the field conditions was calculated in the swards of birdsfoot trefoil grown pure and in mixtures with subterranean clover (kg N/da). The data were averaged and processed statistically using SPSS software (2012).

RESULTS AND DISCUSSION

Agrometeorological conditions in the first year after sowing were unfavourable to the development of crops. Small quantities and unevenly distributed rainfall in May initiated the early spring drought lasting one month. The period was accompanied by monthly average temperatures with values above normal for the area. Evenly distributed rainfall are from the second half of May and in June. After the first decade of July comes drought lasted and in August and September (82 days). These adverse conditions troubled the growth and development of plants.

The second and third years of vegetation were more favourable. The average annual temperature in the second year was with higher values (13.2°C), without extremely high temperatures as well prolonged drought periods and in the third year the amount of rainfall (856.8 l/m²) was greatest.

These conditions impacted on the overall development of birdsfoot trefoil and subterranean clover. Important morphological characteristic object in the current study is the

leaves/stems ratio. It is related to the quality and acceptance of the forage (Cohen 2001, Soster *et al.* 2004). Data show that the leaves/stems ratio in birdsfoot trefoil (alone and in mixtures) during the first year after sowing was with the lowest values (1.08) (Table 1). This is understandable, due to adverse agro meteorological conditions, as well as the age of the crop. In mixtures with subterranean clover values were higher than pure grown birdsfoot trefoil - from 13.1% for the mixture with *Trifolium subterraneum* ssp. *yananicum* (1.23) to 34.9% for the mixture with *Trifolium subterraneum* ssp. *subterraneum* (1.46). *Trifolium subterraneum* ssp. *subterraneum* is subspecies clover, which is small in size leaves less competitive compared to the other two subspecies, lower as well, which has an impact on the development of the accompanying component (birdsfoot trefoil). Pecetti and Piano (1998) found that the size of the leaves is essential for the competitiveness of subterranean clover in mixtures. Competition for basic nutrients, as well as factors of the environment, such as light, heat ect. in a mixture with *Trifolium subterraneum* ssp. *subterraneum* was smaller. Because of this the deviation in value was so high.

The leaves/stems ratio in subterranean clover in pure swards varied in narrow limits and the lowest was found in *Trifolium subterraneum* ssp. *subterraneum*. In mixtures the differences were similar for *Trifolium subterraneum* ssp. *brachycalycinum* and *Trifolium subterraneum* ssp. *yananicum*. Cohen (2001) found a directly proportional relationship between the leaves/stems ratio and protein

Table 1 Leaves/stems ratio in birdsfoot trefoil and subterranean clover pure grown and in mixtures

Treatments	2013	2014	2015
Birdsfoot trefoil (100%)	1.08±0.34	1.49±1.63	1.41±1.43
Birdsfoot trefoil +	1.28±1.27	1.74±0.89	1.62±0.81
<i>Tr. subterr.</i> ssp. <i>brachycalycinum</i>	1.10±0.31	1.26±1.86	1.17±1.20
Birdsfoot trefoil +	1.23±0.24	1.67±2.27	1.61±2.14
<i>Tr. subterr.</i> ssp. <i>yananicum</i>	0.99±0.01	1.05±0.26	1.00±0.02
Birdsfoot trefoil +	1.46±1.29	1.77±3.71	1.56±2.78
<i>Tr. subterr.</i> ssp. <i>subterraneum</i>	0.89±0.02	1.09±0.16	1.11±0.21
<i>Tr. subterr.</i> ssp. <i>brachycalycinum</i> (100%)	0.99±0.07	1.07±0.39	1.01±0.08
<i>Tri. subterr.</i> ssp. <i>yananicum</i> (100%)	0.90±0.45	0.93±0.34	0.95±0.25
<i>Tr. subterr.</i> ssp. <i>subterraneum</i> (100%)	0.85±0.54	0.96±0.80	0.99±0.32
SE (P=0.05) for birdsfoot trefoil	0.07	0.06	0.04
SE (P=0.05) for subterranean clover	0.03	0.05	0.04
±SD for birdsfoot trefoil	0.15	0.12	0.09
±SD for subterranean clover	0.08	0.11	0.08

content in subterranean clover and consider this ratio as an important indicator of the forage quality.

In the second year the leaves/stems ratio in birdsfoot trefoil in mixtures was higher than that in pure sward - from 11.6% for the mixture with *Trifolium subterraneum* ssp. *yananicum* (1.67) to 18.6% for mixtures with *Trifolium subterraneum* ssp. *subterraneum* (1.77). Similar were the increases for subterranean clover as well (12.9%-18.5%). The leaves/stems ratio in birdsfoot trefoil in the last experimental year varied in narrow limits (1.56-1.62).

Notice that, in mixtures of birdsfoot trefoil with subterranean clover the leaves/stems ratio values were higher than its value in pure swards. The higher leaves/stems ratio determines a higher quality and intake of forage and this was found by many authors (Frame *et al.* 1998, Smart *et al.* 1998, Cohen 2001). Improving the leaves/stems ratio in birdsfoot trefoil would have practical significance for increasing quality and preference of the animals to forage.

Data for the leaves/stems ratio on average for the period (Fig 1) showed that in pure grown birdsfoot trefoil it was 1.33. Our results are in agreement with these of Soster *et al.* (2004) and Ducati *et al.* (2015), who have determined some agronomical characteristics in this crop and found leaves/stems ratio with values 1.40. Ayres *et al.* (2008) found higher values ranged from 1.47 to 1.68.

Leaves/stems ratio in birdsfoot trefoil in mixtures with subterranean clover varied between 1.50 to 1.60. In the three mixtures with subterranean clover the leaves/stems ratio was higher than in pure birdsfoot trefoil - by 13.0% for mixtures with *Trifolium subterraneum* ssp. *yananicum*, by 16.3% for this with *Trifolium subterraneum* ssp. *brachycalicinum* and by 20.1% for mixtures with *Trifolium subterraneum* ssp. *subterraneum*.

During the first experimental year total plastid pigments content was lowest due to the age of the sward as well agro meteorological conditions (Table 2). In mixtures of birdsfoot trefoil with *Trifolium subterraneum* ssp. *subterraneum*




Fig 1 Leaves/stems ratio in birdsfoot trefoil and subterranean clover pure grown and in mixtures (average for the period). BT - Birdsfoot trefoil; Trs brach - *Tr. subterraneum* ssp. *brachycalicinum*; Trs yanin - *Tr. subterraneum* ssp. *yananicum*; Trs subter - *Tr. subterraneum* ssp. *subterraneum*. (for birdsfoot trefoil, \pm SD=0.11, SE (P=0.05)=0.05; for subterranean clover, \pm SD=0.10, SE (P=0.05)=0.04

Table 2 Total plastid pigments content in birdsfoot trefoil and subterranean clover pure grown and in mixtures (mg/100 g FW)

Treatments	2013	2014	2015
Birdsfoot trefoil (100%)	213.22	279.78	230.16
Birdsfoot trefoil +	285.86	265.14	331.23
<i>Tr. subterr.</i> ssp. <i>brachycalicinum</i>	273.00	317.64	373.35
Birdsfoot trefoil +	267.49	324.96	336.09
<i>Tr. subterr.</i> ssp. <i>yananicum</i>	252.07	226.71	297.93
Birdsfoot trefoil +	263.91	284.43	358.98
<i>Tr. subterr.</i> ssp. <i>subterraneum</i>	298.24	250.51	287.22
<i>Tr. subterr.</i> ssp. <i>brachycalicinum</i> (100%)	294.57	345.03	335.55
<i>Tri. subterr.</i> ssp. <i>yananicum</i> (100%)	251.52	280.99	249.51
<i>Tr. subterr.</i> ssp. <i>subterraneum</i> (100%)	285.43	315.45	465.36
SE (P=0.05) for birdsfoot trefoil	17.39	13.00	15.87
SE (P=0.05) for subterranean clover	4.91	11.82	20.81
\pm SD for birdsfoot trefoil	34.78	26.00	31.75
\pm SD for subterranean clover	12.03	28.96	50.99

total plastid pigments content was found increased in the two components (for birdsfoot trefoil by 18.0% and for subterranean clover by 6.7%, respectively). The tendency was analogue for the mixture with *Trifolium subterraneum* ssp. *yananicum* (for birdsfoot trefoil by 39.9% and for subterranean clover by 7.9%). In mixture of birdsfoot trefoil with *Trifolium subterraneum* ssp. *brachycalicinum*, total plastid pigments content in birdsfoot trefoil increased by 18.2% and in subterranean clover decreased by 7.7%.

Total plastid pigments content in the second experimental year remains almost unchanged in the mixture of birdsfoot trefoil with *Trifolium subterraneum* ssp. *subterraneum* for both components as compared pure grown crops. In the other two mixtures plastid pigments content in birdsfoot trefoil decreased to 19.0% and in subterranean clovers increased to 8.6%.

Significant increases in the values of this indicator in the components of mixtures as compared to these of pure grown crops were reported in the last experimental year. Total plastid pigments content in all mixtures in birdsfoot trefoil increased (from 8.4 to 29.4%). Total plastid pigments content in subterranean clover increased in *Trifolium subterraneum* ssp. *subterraneum* by 38.5% and in *Trifolium subterraneum* ssp. *brachycalicinum* by 8.4%, and decreased in *Trifolium subterraneum* ssp. *yananicum* by 10.1%. Total plastid pigments content in two components increased in mixtures of birdsfoot trefoil with *Trifolium subterraneum* ssp. *brachycalicinum* and with *Trifolium subterraneum* ssp. *subterraneum*.

For the period of the study a greater increase in the values of the total plastid pigments content in mixtures as compared to pure grown crops was recorded in birdsfoot




Fig 2 Total plastid pigments content in birdsfoot trefoil and subterranean clover pure grown and in mixtures (mg/100 g FW) (average for the period of study). BT - Birdsfoot trefoil; Trs brach - *Tr. subterraneum* ssp. *brachycalicinum*; *Trs yanin* - *Tr. subterraneum* ssp. *yaninicum*; *Trs subter* - *Tr. subterraneum* ssp. *subterraneum*. (for birdsfoot trefoil, \pm SD=15.37, SE (P=0.05)=7.68; for subterranean clover, \pm SD=8.84, SE (P=0.05)=21.66)

trefoil. Data on average for the period (Fig 2) show this trend. Total plastid pigments content in birdsfoot trefoil was 241.05 mg/100 g FW, but in subterranean clover varied from 294.08 mg/100 g FW to 321.33 mg/100 g FW. Total plastid pigments content in mixtures changes in both components. So, in birdsfoot trefoil it increased from 7.4 to 15.6%. In subterranean clover total plastid pigments content significantly increased only in *Trifolium subterraneum* ssp. *subterraneum* – by 14.8%.

Total plastid pigments content significantly increased in both components of mixtures of birdsfoot trefoil (by 8.1%) with *Trifolium subterraneum* ssp. *subterraneum* (by 14.8%). These data correspond to the change in the leaves/stems ratio, the coefficient of correlation between leaves/stems ratio and total plastid pigments is $r = 0.60$.

The amount of fixed nitrogen in the mixtures changes and depends on the competition between the components. There were amount of fixed nitrogen from 1.71 N/da (birdsfoot trefoil + *Trifolium subterraneum* ssp. *subterraneum*) to 4.84 kg N/da (birdsfoot trefoil + *Trifolium subterraneum* ssp. *brachycalicinum*) more as compared to the amount obtained by pure grown birdsfoot trefoil (Fig 3). Competition for soil nitrogen mixtures may have a beneficial effect on nitrogen fixing process as in the case birdsfoot trefoil and subterranean clover have the same type of nitrogen exchange. They fix nitrogen from the atmosphere through nitrogenase in the nodules.

Changes in the morphological and physiological parameters of the crops studied are associated with the metabolic role of nodules, which are strong acceptor of carbon assimilates from leaves, but are simultaneously and donor of organic nitrogen for leaves, mainly in the form of transport amino acids and amides (Atkins *et al.* 1986).

Leaves/stems ratio for pure grown birdsfoot trefoil (1.33), *Trifolium subterraneum* ssp. *brachycalicinum* (1.02), *Trifolium subterraneum* ssp. *yaninicum* (0.93) and *Trifolium*




Fig 3 Amount of fixed nitrogen from birdsfoot trefoil pure grown and in mixtures with subterranean clover for three years period. BT - Birdsfoot trefoil; Trs brach - *Tr. subterr.* ssp. *brachycalicinum*; *Trs yanin* - *Tr. subterr.* ssp. *yaninicum*; *Trs subter* - *Tr. subterr.* ssp. *subterraneum* (\pm SD=2.09, SE (P=0.05)=1.04)

subterraneum ssp. *subterraneum* (0.93) was found.

Leaves/stems ratio for birdsfoot trefoil in all mixtures was found be higher as compared to pure grown birdsfoot trefoil - by 13.0% for the mixture with *Trifolium subterraneum* ssp. *yaninicum*, by 16.3% for that with *Trifolium subterraneum* ssp. *brachycalicinum* and by 20.1% for the mixture with *Trifolium subterraneum* ssp. *subterraneum*.

Total plastid pigments content in birdsfoot trefoil increased in all mixtures tested: with *Trifolium subterraneum* ssp. *brachycalicinum* by 7.4%, with *Trifolium subterraneum* ssp. *subterraneum* by 8.1% and with *Trifolium subterraneum* ssp. *yaninicum* by 15.6%. Total plastid pigments content in subterranean clover was found increased only for *Trifolium subterraneum* ssp. *subterraneum* (by 14.8%).

From mixtures of birdsfoot trefoil were obtained more fixed amount of nitrogen - from 1.71 kg N/da (with *Trifolium subterraneum* ssp. *subterraneum*) to 4.84 kg N/da (with *Trifolium subterraneum* ssp. *brachycalicinum*) as compared to pure grown birdsfoot trefoil. Birdsfoot trefoil and subterranean clover in mixtures showed good tolerance, morphological and physiological status.

REFERENCES

Atkins B T, Kegl J and Levin B. 1986. Explicit and implicit information in dictionaries. Lexicon Project Working Papers 12, Center for Cognitive Science, MIT, Cambridge, MA.
 Ayres J F, Kelman W M and Lane L A. 2008. Phoenix, Venture and Matador – new birdsfoot trefoil cultivars for permanent pasture applications in eastern Australia. (In) *Proceedings of the 23rd Annual Conference of the Grassland Society of NSW, Grassland Society of NSW Inc: Orange*, pp 107–9.
 Buxton D R, Hornstein J S, Wedin W F and Marten G C. 1985.

- Forage quality in stratified canopies of alfalfa, birdsfoot trefoil, and red clover. *Crop Science* **25**: 273–79.
- Carlsson G and Huss-Danell K. 2003. Nitrogen fixation in perennial forage legumes in the field. *Plant and Soil* **253**: 353–72.
- Chourkova B. 2014. Productivity and botanical composition of a mixed sward of birdsfoot trefoil and red fescue depending on the term of sowing and proportion of components. *International Journal of Agriculture Innovations and Research* **3** (1): 276–80.
- Cohen D C. 2001. Degradability of crude protein from clover herbage used in irrigated dairy production systems in northern Victoria. *Australian Journal of Agricultural Research* **52**(3): 415–25.
- Ducati C, Neres M A, Castagnara D D, Oliveira P S R , Weirich D, Silva G M, and Sunahara S M M. 2015. Forage potential of oat Iapar 61 alone or in association with birdsfoot trefoil. *Cien. Inv. Agr.* **42**(3): 341–51.
- Frame J, Charlton J F L and Laidlaw A S. 1998. *Temperate Forage Legumes*, p 327. CAB International, Wallingford.
- Ilieva A, Vasileva V and Katova A. 2015. The effect of mixed planting of birdsfoot trefoil, sainfoin, subterranean clover, and tall fescue on nodulation, and nitrate reductase activity in shoots. *Journal of Global Agriculture and Ecology* **3**(4): 222–8.
- Kusvuran A, Ralice Y and Saglamtimur T. 2014. Determining the biomass production capacities of certain forage grasses and legumes and their mixtures under mediterranean regional conditions. *Acta Advances Agricultural Sciences* **2**: 13–24.
- Luscher A, Mueller-Harvey I, Soussana J F, Rees R M and Peyraud J L. 2014. Potential of legume-based grassland-livestock systems in Europe: a review. *Grass and Forage Science* **69**: 206–28.
- Nichiporovich A. 1971. Some principles to optimize photosynthetic activity of plants in crops. L. Nauka 104–8.
- Nichols P G H, Revell C K, Humphries A W, Howie J H, Hall E J, Sandral G A, Ghamkhar K and Harris C A. 2012. Temperate pasture legumes in Australia – their history, current use and future prospects. *Crop and Pasture Science* **63**: 691–725.
- Pecetti L and Piano E. 1998. Leaf size variation in subterranean clover (*Trifolium subterraneum* L. sensu lato). *Genetic Resources and Crop Evolution* **45**(2): 161–5.
- Peeters A, Parente G and Gall A. 2006. Temperate legumes: key-species for sustainable temperate mixtures. *Grassland Science in Europe*, **11**: 205–0.
- Porqueddu C, Parenteand G and Elsaesser M. 2003. Potential of grasslands. *Grassland Science in Europe* **8**: 11–20.
- Smart A J, Schacht W H, Pedersen J F, Undesander D J and Moser L E. 1998. Prediction of leaf:stem ratio in grasses using near infrared reflectance spectroscopy. *Journal of Range Management* **51**(4): 447–9.
- Smirnova E, Reshetnikova V, Makarova T and Karavaeva G. 2013. Features of genotic relations in the one-specy and mixed crops of *Mellilotus officinalis* L. *Proceedings of the Samara Scientific Center of RAS* **15**(3): 793–5.
- Soster M. T B, Scheffer-Basso S M, Dall’Agnol M, Brustolin R and R S Fontaneli. 2004. Caracterização agrônômica de genótipos de cornichão (*Lotus corniculatus* L.). *Revista Brasileira de Zootecnia* **33**: 1 662–71.
- SPSS 2012. SPSS Version 20.0. *SPSS Inc.*, 233 S. Wacker Drive, Chicago, Illinois.
- Stevanovic P, Vuckovic S, Popovic V, Ikanovic J, Zivanovic L, Tabakovic M, Vujic R and Lakic Z. 2015. Influence of the mineral fertilization at morphological and productive characteristics of the *Lotus corniculatus* on pseudogley *Wulfenia*. *Austria* **22**(10): 190–204.
- Vasilev E. 2006. Productivity of subterranean clover (*Tr. subterraneum* L.) in pasture mixtures with some perennial grasses for the conditions of Central North Bulgaria. *Plant Science S* **4**(2): 149–52.
- Vasileva V. and Vasilev E. 2012. Dry mass yield from some pasture mixtures with subterranean clover (*Trifolium subterraneum* L.). *Journal of Mountain Agriculture on the Balkans* **15**: 1 024–33.
- Vuckovic S M. 2004. Travnjaci. Poljoprivredni fakultet Univerzitet, ISBN868073375X, pp 506
- Zelenskii M and Mogileva G. 1980. Comparative evaluation of photosynthetic ability of agricultural crops by photochemical activity of chloroplasts. VIR, Leningrad, p 36 (in Russian).



Yield, economics, nutrient uptake and quality of coriander (*Coriandrum sativum*) under different sowing time, varieties and plant growth regulators

B R KURI¹, N L JAT², A C SHIVRAN³, S L JAT⁴ LALIT KUMAR⁵, S PANWAR⁶ and C M PARIHAR⁷

S. K. N. College of Agriculture Jobner, Rajasthan 303 329

Received: 11 August 2016; Accepted: 14 September 2016

ABSTRACT

A field experiment was carried out in *rabi* seasons of 2011-12 and 2012-13 at Agronomy Farm, S.K.N. College of Agriculture, Jobner (Rajasthan), to assess the effect of sowing time and plant growth regulators on yield, economics, nutrient uptake and quality of coriander (*Coriandrum sativum* L.) varieties grown under irrigated loamy sand soil of semi-arid eastern plain zone of Rajasthan. The experiment consisting of two sowing times, viz. normal (1st week of November) and late (3rd week of November), three varieties, viz. RCr-41, RCr-435 and RCr-480 and five plant growth regulators treatments (PGRs), viz. triacontanol @ 1000 ppm, brassinolide @ 1.0 ppm, thiourea @ 1000 ppm and NAA @ 50 ppm, including one control (water sprayed) treatment were replicated thrice in split-plot design. The 2-year results revealed that the normal sown coriander crop produced significantly higher seed (by 31%) and stover yields, total N and P uptake (by 20 and 25%), per-day productivity (by 17%) and net returns (by 59%), partial factor productivity (PFP), nutrient harvest index (NHI), protein (by 21%) and oil (by 41%) yields compared to late sown crop. However, late sown crop resulted significantly higher nitrogen content in seed and stover and protein content in seed over normal sown. Among 3-varieties, RCr-480 produced significantly higher yields (15%), total N (15%) and P (by 14%) uptake, per-day productivity (by 14%) and net returns (by 34%), essential oil (by 21%) and protein (17%) yields, PFP-NP and NHI-NP during both the years as compared to RCr-41, which were statistically at par with RCr-435, except oil yield, NHI-NP in second year. The foliar spray of PGRs significantly enhanced the yields (by 10-22%), nutrient content and uptake and quality of coriander over water spray. The highest seed and stover yields, N and P uptake, per-day productivity and net returns, protein and oil yield and PFP-NP with 1000 ppm triacontanol, over control, 50 ppm NAA and 1000 ppm thiourea, and remained statistically at par with 1.0 ppm brassinolide, except per-day-net returns and oil yield. Overall, our results suggest that adoption of RCr-480 variety under normal sown condition with 1000 ppm triacontanol spray can enhance crop productivity, profitability, nutrient uptake and quality of coriander in semi-arid region of India and elsewhere under similar agro-climatic conditions.

Key words: Coriander, Nutrient uptake, Partial factor productivity, Plant growth regulators, Profitability

Coriander (*Coriandrum sativum* L.) is one of the oldest seed spices most widely used by the mankind as condiment. The green leaves and fruits of coriander have a pleasant aromatic odour due to presence of 'coriandrol'. Good quality oleoresin has extracted from coriander seed which can be used for flavouring beverages, sweets, pickles, sausages, snacks, etc. The research findings across the country conclusively revealed negative impact of delayed sowing on yield, nutrient content and uptake and quality of seed especially on test weight but the extent of reduction depends

upon the prevailing weather conditions. A late sown crop produces shrivelled grains as well as poor quality compared to early sowing leading to low yield.

Selection of proper improved variety as per the growing conditions is another way to improve the productivity and quality of the crop. All the three (RCr-41, RCr-435 and RCr-480) varieties are suitable for normal conditions and covers a large area of state but yield stagnation has been reported prevailing management practices. Therefore, it was felt necessary to enhance the productivity of coriander through alternative practices like use of PGRs. Use of triacontanol, brassinolide, thiourea and NAA based PGRs has been reported to induce physiological efficiencies including photosynthetic ability of plants (Jain *et al.* 1988, Sarada *et al.* 2008), growth promoting activity (Mandava 1988), also improve the resistance in plant against the water, salinity, low and high temperature stresses (Rao *et al.* 2002), improve the dry matter partitioning and subsequently enhance the

¹Ph D Scholar (e mail: brkuri84@gmail.com), ^{2&3}Professor, S K N College of Agriculture Jobner (e mail: nlsepat@gmail.com & acs_shivran@rediffmail.com), Rajasthan ⁴Scientist (e mail: sliari@gmail.com), ⁵The Fertilizer Association of India, New Delhi (e mail: lalitgoliyan81@gmail.com), ⁶Indian Council of Agricultural Research, New Delhi (e mail: scientist1775@gmail.com), ⁷Scientist (e mail: pariharc@gmail.com), ICAR-IIMR, New Delhi.

productivity of crops and may mitigate the problem of late sowing to bring marked biological activity in plants (Sahu and Solanki 1991). Thus, for achieving the attainable yield potential with good quality of the crop the optimum time of sowing, adoption of most suitable varieties and application of plant growth regulators are essentially required. So keeping in view the above facts, the present investigation has been planned to evaluate the effect of sowing times, varieties and plant growth regulators on yield, economics, nutrient uptake and quality of coriander.

MATERIALS AND METHODS

A field experiment was conducted for two consecutive years in *rabi* seasons of 2011-12 and 2012-13 at the Agronomy Farm, S K N College of Agriculture, Jobner, situated at 27° 05' N latitude, 75° 28' E longitude and at an altitude of 427 AMSL in Jaipur district of Rajasthan. This region is categorized as semi-arid eastern plain zone of Rajasthan. The soil of the experimental site was loamy sand in texture with low in organic carbon (0.16%) and available nitrogen (126.5 kg/ha), medium in available phosphorus (16.7 kg/ha) and potassium (171.9 kg/ha) and alkaline in reaction with pH 8.3. The experiment consisting of two sowing times viz., normal (1st week of November) and late (3rd week of November), 3-varieties, viz. RCr-41, RCr-435 and RCr-480 as main plot treatments and five treatments of plant growth regulators viz., triacontanol @ 1 000 ppm, brassinolide @ 1.0 ppm, thiourea @ 1 000 ppm and NAA @ 50 ppm including control (water sprayed) as sub-plot treatment, total 30 treatment combinations were replicated thrice in split plot design. Before sowing coriander varieties seed was treated with bavistin @ 2 g/kg of seed. The crop was sown in rows at 30 cm apart and 2-3 cm depth with a seed rate of 12 kg/ha as per treatments under normal and late sown conditions. A uniform dose of 60 kg N, 40 kg P₂O₅ and 20 kg K₂O/ha was applied in all plots. Entire quantity of P and K and half dose of N were applied at the time of sowing and the remaining nitrogen was applied in two equal splits with third and fourth irrigations. Two foliar spray of triacontanol @ 1 000 ppm, brassinolide @ 1.0 ppm, thiourea @ 1 000 ppm and NAA @ 50 ppm were done at 40 and 70 days after sowing. The spray volume was 400 and 600 litres/ha for application at 40 and 70 DAS, respectively.

Coriander was sown on 4th and 1st November (under normal condition) and 19th and 16th November (under late sown condition) in 2011 and 2012, respectively, and the crop was harvested on 22 and 21 March during 2012 and 2013, respectively. The plant samples were collected at harvest and dried in oven ($\pm 70^{\circ}\text{C}$), processed and thoroughly mixed and analyzed for estimation of nitrogen, phosphorus and essential oil content was by Kjeldahl, Vanado-molybdate yellow colour and Clevenger apparatus, respectively. N and P uptake were calculated by formula (Nutrient content/100 multiplied by seed/stover yield in kg/ha). The estimated values of different nutrient use indices viz; partial factor productivity (PFP), and harvest index (HI) of applied N and P were computed by using the formulas as suggested by

Fageria and Baligar (2003) and Dobermann (2005) which are as follows:

$$\text{Partial factor productivity of applied nutrient (PFP)} = Y_t / N_a$$

$$\text{Nutrient harvest index (\%)} = N_s / N_t \times 100$$

where, Y_t = Yield under treatment plots (kg/ha); N_a = Amount of nutrient added (kg/ha); N_s = Nutrient uptake by seed at harvest; N_t = Nutrient uptake by whole plant (seed + stover) at harvest.

Yield, nutrient uptake and quality parameters were measured as per standard procedures. Economics was calculated based on prevailing cost of input and output in local market. All the data were statistically analyzed using the analysis of variance (ANOVA) techniques (Gomez and Gomez 1984). The critical difference at 0.05% level probability were calculated to assess the significance between treatments.

RESULTS AND DISCUSSION

Effect of sowing time

The sowing times significantly influenced the yield, economics, nutrient content and uptake and quality parameters of coriander. Normal sown crop resulted significantly higher seed (1456 and 1501 kg/ha) and stover (2150 and 2221 kg/ha) yields (Table 1), productivity (10.40 and 10.65 kg/day/ha) and net returns (₹ 228 and 239/day/ha) in 2011-12 and 2012-13 over late sown crop, respectively (Table 2).

It is an established fact that sinks capacity largely governs crop productivity (Evans 1975). This might be due to favourable environmental conditions available to the crop during its initial growth, flowering and fruiting stages. The possible reason for low yield in delayed sowing might be due to availability of insufficient time for vegetative growth as the plant entered the reproductive phase at a faster rate. These results are in conformity with the findings of Meena *et al.* (2006) and Bhadkariya *et al.* (2007).

The similar results were observed in total (seed + stover) N (62.5 and 66.5 kg/ha) and P (13.2 and 14.3) uptake (Fig 1 and 2) as well as uptake in seed and stover individually, total NP-PFP (14.56 and 15.01) and NHI of (65.0 and 64.6%) as well as individual nutrient PFP and NHI was significantly higher with normal sown crop compared to late sown crop in 2011-12 and 2012-13, except NHI-N and NHI-NP in 2011-12, respectively (Table 1, 2 and 3). However, N content in seed and N and P content in stover were significantly higher with late sown crop during both the years. Therefore, the nitrogen content in both seed and stover followed a reverse trend to that of seed and stover yield. A decreased nitrogen content under normal sown crop could be due to the dilution effect by higher yield. The increased nitrogen and phosphorus uptake under normal sown crop could be ascribed due to higher total biomass production. Thus, significant improvement in seed and stover yield by virtue of higher photosynthesis under normal sown crop ultimately resulted in higher uptake of nitrogen and

Table 1 Effect of sowing time, varieties and plant growth regulators on yields, nitrogen content and uptake in seed and stover of coriander

Treatment	Seed yield (kg/ha)		Stover yield (kg/ha)		N content (%)				N uptake (kg/ha)			
					Seed		Stover		Seed		Stover	
	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13
<i>Sowing time</i>												
Normal	1456	1501	2150	2221	2.84	2.92	0.969	1.005	41.5	44.0	21.0	22.5
Late	1106	1143	1735	1807	3.09	3.16	1.026	1.065	34.3	36.2	17.6	19.0
SEM±	22	21	33	32	0.04	0.03	0.012	0.011	0.44	0.42	0.22	0.22
CD (P=0.05)	69	66	104	100	0.11	0.10	0.037	0.035	1.37	1.31	0.70	0.68
<i>Varieties</i>												
RCr-41	1179	1217	1853	1921	2.90	2.98	0.979	1.017	34.1	36.2	17.8	19.2
RCr-435	1316	1356	1998	2069	3.02	3.08	1.012	1.050	39.6	41.7	20.1	21.6
RCr-480	1347	1393	1976	2053	2.98	3.06	1.003	1.038	40.0	42.5	19.9	21.4
SEM±	27	26	40	39	0.04	0.04	0.014	0.014	0.53	0.51	0.27	0.27
CD (P=0.05)	85	81	127	123	NS	NS	NS	NS	1.68	1.61	0.86	0.83
<i>Plant growth regulators</i>												
Control	1128	1178	1733	1825	2.66	2.72	0.918	0.952	29.9	31.9	15.8	17.3
Tricantanol @ 1000 ppm	1380	1421	2061	2152	2.97	3.06	1.009	1.048	40.8	43.3	21.2	22.4
Brassinolide @ 1 ppm	1365	1398	2109	2107	3.06	3.13	1.020	1.058	41.6	43.6	20.7	22.1
Thiourea @ 1000 ppm	1273	1315	1949	2052	3.11	3.19	1.029	1.067	39.4	41.8	20.1	21.8
NAA @ 50 ppm	1258	1298	1860	1935	3.03	3.10	1.012	1.050	37.9	40.1	18.7	20.2
SEM±	24	25	37	38	0.05	0.04	0.015	0.015	0.56	0.54	0.29	0.28
CD (P=0.05)	69	70	104	106	0.13	0.12	0.043	0.042	1.60	1.54	0.81	0.79

Table 2 Effect of sowing time, varieties and plant growth regulators on phosphorus content and uptake in seed and stover, per day productivity and net returns of coriander

Treatments	P content (%)				P uptake (kg/ha)				Productivity/day (kg/ha)		Net returns/day (Rs/ha)	
	Seed		Stover		Seed		Stover					
	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13
<i>Sowing time</i>												
Normal	0.530	0.550	0.253	0.269	7.73	8.27	5.48	6.02	10.40	10.65	228	239
Late	0.538	0.560	0.269	0.283	5.96	6.41	4.60	5.03	8.85	9.07	143	154
SEM+	0.007	0.006	0.003	0.003	0.10	0.09	0.06	0.06	0.16	0.15	3	3
CD (P=0.05)	NS	NS	0.010	0.011	0.31	0.28	0.19	0.18	0.50	0.48	11	10
<i>Varieties</i>												
RCr-41	0.525	0.548	0.256	0.273	6.19	6.66	4.66	5.16	8.86	9.08	155	165
RCr-435	0.540	0.560	0.265	0.279	7.11	7.59	5.27	5.75	9.89	10.11	196	207
RCr-480	0.537	0.558	0.261	0.275	7.24	7.77	5.18	5.67	10.12	10.39	206	218
SEM+	0.009	0.007	0.004	0.004	0.12	0.11	0.07	0.07	0.20	0.19	4	4
CD (P=0.05)	NS	NS	NS	NS	0.38	0.35	0.23	0.22	0.62	0.59	13	13
<i>Plant Growth Regulators</i>												
Control	0.513	0.534	0.248	0.261	5.78	6.29	4.27	4.73	8.47	8.78	149	163
Tricantanol @ 1000ppm	0.538	0.559	0.262	0.277	7.42	7.94	5.50	5.92	10.37	10.60	218	228
Brassinolide @ 1 ppm	0.545	0.565	0.268	0.284	7.44	7.89	5.43	5.94	10.26	10.43	186	195
Thiourea @ 1000 ppm	0.540	0.562	0.266	0.281	6.87	7.39	5.20	5.73	9.56	9.81	190	201
NAA @ 50 ppm	0.534	0.555	0.259	0.276	6.72	7.20	4.79	5.31	9.45	9.68	185	196
SEM+	0.008	0.008	0.004	0.004	0.11	0.10	0.07	0.07	0.18	0.18	4	4
CD (P=0.05)	0.024	0.022	0.011	0.011	0.31	0.28	0.21	0.21	0.52	0.52	10	11




Fig 1 Effect of sowing time, varieties and plant growth regulators on total N uptake by coriander




Fig 2 Effect of sowing time, varieties and plant growth regulators on total P uptake by coriander

phosphorus over late sown crop.

Quality parameters (protein and essential oil content in seed) of coriander were also significantly influenced with sowing times (Table 4). The significantly higher protein yield (260 and 275 kg/ha), oil yield (5.58 and 5.96 kg/ha) and essential oil content in seed were observed with normal sown crop as compared to late sown crop, while significantly higher protein content in seed was recorded with late sowing during both years, respectively. The reason could be traced back in temperature mediated effects on plants physiological processes and adequate absorption of nutrients. These findings are in close conformity with the findings of Balai and Keshwa (2011).

Effect of varieties

Coriander varieties performed differentially due to agronomic management and application of PGRs. RCr-480 recorded significantly higher seed (1 347 and 1 393 kg/ha), stover (1 976 and 2 053 kg/ha) yields (Table 1), crop

productivity (10.12 and 10.39 kg/day/ha) and net returns (₹ 206 and 218/day/ha) compared to RCr-41 and being at par with RCr-435 during 2011-12 and 2012-13, respectively. The RCr-480 variety resulted in average (mean of 2-year) increase of 14, 7, 14 and 33% in seed yield, stover yield, per-day-productivity and net returns over RCr-41. Thus, the improvement in above these regulative processes of coriander evidenced from higher biomass and nutrients accumulation in RCr-480 and RCr-435 varieties over RCr-41. The marked variation in yield and economics of these varieties RCr-480, RCr-435 and RCr-41 were also observed under multi-location trials conducted under AICRP on spices (AICRPS, 2011 and Balai and Keshwa 2010).

Nutrient uptake and quality of coriander seed were also significantly differed in all the 3-varieties (Table 1, 2 and 3). However, the N and P content in seed and stover of all the varieties remained statistically at par. Total uptake of N (59.9 and 63.9 kg/ha) and P (12.4 and 13.4 kg/ha) (Fig 1 and 2) as well as individual N and P uptake by seed and stover,

Table 3 Effect of sowing time, varieties and plant growth regulators on partial factor productivity (PFP) and nutrient harvest index (NHI) of nitrogen and phosphorous in coriander

Treatment	PFP-N		PFP-P		PFP-NP		NHI-N		NHI-P		NHI-NP	
	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13
<i>Sowing time</i>												
Normal	24.26	25.02	36.39	37.53	14.56	15.01	66.4	66.0	58.5	57.8	65.0	64.6
Late	18.43	19.05	27.65	28.58	11.06	11.43	66.1	65.6	56.5	56.0	64.4	63.9
SEM±	0.37	0.35	0.55	0.53	0.22	0.21	0.3	0.1	0.4	0.0	0.2	0.1
CD (P=0.05)	1.16	1.11	1.73	1.66	0.69	0.66	NS	0.3	1.4	0.1	NS	0.3
<i>Varieties</i>												
RCr-41	19.65	20.28	29.48	30.43	11.79	12.17	65.6	65.2	57.0	56.3	64.1	63.7
RCr-435	21.94	22.60	32.90	33.90	13.16	13.56	66.3	65.8	57.3	56.8	64.8	64.2
RCr-480	22.45	23.22	33.68	34.83	13.47	13.93	66.7	66.5	58.2	57.7	65.3	64.9
SEM±	0.45	0.43	0.68	0.65	0.27	0.26	0.3	0.1	0.5	0.1	0.3	0.1
CD (P=0.05)	1.42	1.35	2.12	2.03	0.85	0.81	1.0	0.4	NS	0.3	0.9	0.3
<i>Plant growth regulators</i>												
Control	18.80	19.63	28.20	29.45	11.28	11.78	65.4	64.9	57.4	56.9	63.9	63.4
Tricantanol @ 1000 ppm	23.00	23.68	34.50	35.53	13.80	14.21	65.8	65.9	57.3	57.1	64.3	64.3
Brassinolide @ 1 ppm	22.75	23.30	34.13	34.95	13.65	13.98	66.8	66.3	57.7	56.9	65.2	64.7
Thiourea @ 1000 ppm	21.22	21.92	31.83	32.88	12.73	13.15	66.2	65.7	56.8	56.2	64.6	64.1
NAA @ 50 ppm	20.97	21.63	31.45	32.45	12.58	12.98	66.9	66.4	58.2	57.4	65.5	64.9
SEM±	0.40	0.41	0.61	0.62	0.24	0.25	0.2	0.1	0.3	0.3	0.2	0.1
CD (P=0.05)	1.15	1.17	1.72	1.76	0.69	0.70	0.5	0.3	0.9	0.8	0.5	0.3

total PFP (13.47 and 13.93) and NHI (65.3 and 64.9%) as well as individual nutrient PFP and NHI were recorded significantly higher with RCr-480 and remained at par with RCr-435 compared to RCr-41 in both the years. Nutrient uptake is a cumulative effect of content and biomass yield, so the varieties having higher content will not every time uptake the higher nutrients.

During both years essential oil content in RCr-480 variety seed (0.379 and 0.393%), oil yield (5.11 and 5.47 kg/ha) and protein yield (251 and 266 kg/ha) was recorded significantly higher as compared to RCr-41, respectively, being at par with RCr-435 (Table 4). The results of the present investigation indicate differential behavior of coriander varieties with respect to nutrient content, quality and its uptake is in close conformity with findings of other workers (AICRPS 2011 and Balai and Keshwa 2011).

Effects of plant growth regulators

Foliar spray of plant growth regulators (PGRs) significantly increased the yields, per-day-productivity and net returns of coriander as compared to water sprayed (control). Among the PGRs, foliar spray of 1000 ppm triacontanol enhanced significantly seed (1380 and 1421 kg/ha and stover (2061 and 2152 kg/ha) yields, productivity (10.37 and 10.60 kg/day/ha) and net returns (₹ 218 and 228/day/ha) as compared to water sprayed, 50 ppm NAA and 1000 ppm thiourea being at par with 1.0 ppm brassinolide (Table 1 and 2). Under all the PGRs

treatments the increase in seed and stover yields, per-day productivity and net returns was 10-22, 6-22, 10-22 and 20-46% over control, respectively. This might be due to the stimulatory effect of growth regulators which induced large number of reproductive sinks leading to greater activity of carboxylating enzymes (ribose 1,5-di phosphate carboxylase) resulting in higher photosynthetic rates with greater translocation and accumulation of metabolites in sink (Nehara *et al.* 2006) and ultimately higher seed yield. The cost of triacontanol was lower in comparison to added outputs. Thus, the increased seed yield led to higher net returns. The similar response with foliar spray of triacontanol was also reported by Nehara *et al.* (2006) in fenugreek and Sarada *et al.* (2008) in coriander.

Foliar application of plant growth regulators was also brought significant effect on nutrient content, uptake and quality of coriander (Table 1 and 2). Foliar spray of 1000 ppm thiourea recorded significantly higher nitrogen content in both seed (3.11 and 3.19%) and stover (1.029 and 1.067%) as compared to control but found to be statistically at par with other PGRs treatment. Phosphorus content in seed and stover were significantly higher under foliar spray of 1.0 ppm brassinolide as compared to control and statistically at par with other PGRs treatment. However, foliar spray of 1000 ppm triacontanol recorded significantly higher total N (62.0 and 65.7 kg/ha) and P (12.9 and 13.9 kg/ha) uptake (Fig 1 and 2) as well as individual nutrient uptake by seed and stover over control and 50 ppm NAA but

Table 4. Effect of sowing time, varieties and plant growth regulators on protein and essential oil content in seed and yields of coriander

Treatments	Protein content (%)		Protein yield (kg/ha)		Essential oil content (%)		Oil yield (kg/ha)	
	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13	2011-12	2012-13
<i>Sowing time</i>								
Normal	17.8	18.3	260	275	0.383	0.397	5.58	5.96
Late	19.3	19.8	215	227	0.357	0.370	3.95	4.23
SEm±	0.27	0.23	6	6	0.005	0.004	0.05	0.06
CD (P=0.05)	0.84	0.73	20	20	0.015	0.014	0.17	0.19
<i>Varieties</i>								
RCr-41	18.1	18.6	214	227	0.361	0.373	4.26	4.54
RCr-435	18.9	19.3	248	261	0.371	0.385	4.88	5.22
RCr-480	18.6	19.1	251	266	0.379	0.393	5.11	5.47
SEm±	0.33	0.28	8	8	0.006	0.005	0.08	0.09
CD (P=0.05)	NS	NS	24	24	0.018	0.017	0.23	0.24
<i>Plant growth regulators</i>								
Control	16.6	17.0	187	200	0.331	0.344	3.73	4.05
Tricantanol @ 1000 ppm	18.6	19.1	255	271	0.383	0.397	5.29	5.64
Brassinolide @ 1 ppm	19.1	19.6	261	273	0.371	0.384	5.06	5.37
Thiourea @ 1000 ppm	19.4	19.9	247	262	0.391	0.406	4.98	5.34
NAA @ 50 ppm	18.9	19.4	238	251	0.376	0.388	4.73	5.04
SEm±	0.28	0.27	8	8	0.006	0.005	0.06	0.07
CD (P=0.05)	0.80	0.76	22	22	0.016	0.015	0.18	0.20

remained statistically at par with 1.0 ppm brassinolide and 1000 ppm thiourea except in thiourea N uptake by stover during 2011-12, P uptake by seed in both the years and uptake by stover in 2011-12, respectively. The application of PGRs increased 22-36% and 14-28% in total N and P uptake over control. Total PFP (13.80 and 14.21) as well as individual nutrients PFP were recorded significantly higher with 1000 ppm triacontanol, and remained at par with 1.0 ppm brassinolide and total NHI (65.5 and 64.9%) with 50 ppm NAA, and remained at par with 1000 ppm thiourea and control in NHI-P as well as the individual nutrient as compared to other PGRs treatment and control.

However, the significantly higher protein (19.4 and 19.9%) and essential oil content (0.391 and 0.406%) in seed under 1000 ppm thiourea over control and 50 ppm NAA. The significantly higher protein yield (261 and 273 kg/ha) were recorded under 1.0 ppm brassinolide over control and oil yield (5.29 and 5.64 kg/ha) was recorded with 1000 ppm triacontanol than rest of the treatments of PGRs and control. The increase in nitrogen content in seed, stover, protein and essential oil content in seed with thiourea application might be due to improvement of metabolic processes and better growth and development, leading to greater absorption of nutrients from rhizosphere leads to increase yield and quality. Thiourea also creates better microbial population in soil which are responsible to mobilize essential nutrients. The quantum of nutrient uptake and oil yield by the crop is dependent on biomass production and nutrient content at cellular level. Since,

nutrients content are improved by application of PGRs, the total biomass production is primarily responsive for the quantum of nutrient uptake. These results are in close conformity with the findings of Balai and Keshwa (2011) and Nehara *et al.* (2006).

Thus, on the basis of results of 2-years investigation, it could be concluded that sowing of coriander variety RCr-480 in first week of November with foliar application of 1000 ppm triacontanol at 40 and 70 DAS gives higher yield, per-day-productivity and net returns, nutrient uptake and quality of coriander seed in semi-arid eastern plain zone of Rajasthan.

REFERENCES

- AICRPS. 2011. *Annual Report 2010-11*. All India Co-ordinated Research Project on Spices. S K N College of Agriculture, Jobner.
- Balai L R and Keshwa G L. 2010. Effect of thiourea on yield and economics of coriander (*Coriandrum sativum* L.) varieties under normal and late sowing conditions. *Journal of Progressive Agriculture* **1**(1): 52-5.
- Balai L R and Keshwa G L. 2011. Effect of thiourea on yield and nutrient uptake of coriander (*Coriandrum sativum* L.) varieties under normal and late sowing conditions. *Journal of Spices and Aromatic Crops* **20**(1): 34-7.
- Bhadkariya S K, Gupta A, Babade A, Kasana B S and Tomer L S. 2007. Effect of different time of sowing on growth, yield and seed quality of coriander (*Coriandrum sativum* L.) cv. CIMPOS- 33. *Bhartiya Krishi Anusandhan Parika* **22**(3): 229-32.
- Evans L T. 1975. *Crop Physiology*. University Press, Cambridge.

- Gomez K A, and Gomez A A. 1984. *Statistical Procedures for Agricultural Research*, Second Edition. An International Rice Research Institute Book. A Wiley-Inter- science Publication. John Wiley & Sons, New York.
- Jain G L, Sharma G L, Rajput M S and Jain H K. 1988. Effect of some new growth substances on yield attributes and yield of groundnut. (In) Abstracts, *Symposium on Efficient Cropping System Zones of India*, 7-10 January 2009, UAS, Bangalore, pp 75.
- Krishna De A. 1999. Spices: Some known and unknown facts. *Science and Culture* **65**: 220-8.
- Mandava N B. 1988. Plant growth promoting brassinosteroids. *Annals of Review Plant Physiology* **39**: 23-52.
- Meena S S, Sen N L and Malhotra S K. 2006. Influence of sowing date, nitrogen and plant growth regulators on growth and yield of coriander (*Coriandrum sativum* L.). *Journal of Spices and Aromatic Crops* **15**(2): 88-92.
- Nehara K C, Kumawat P D and Singh B P. 2006. Response of fenugreek (*Trigonella foenum-graecum*) to phosphorus, sulphur and plant growth regulators under semi-arid plains zone of Rajasthan. *Indian Journal of Agronomy* **51**(1): 73-6.
- Rao S S R, Vardhini B N, Sujatha E and Anuradha S. 2002. Brassinosteroids: A new class of phytohormones. *Current Science* **81**: 1 339-45.
- Sahu M P and Solanki A S. 1991. Role of sulphydral compounds in improving dry matter accumulation and grain production of maize (*Zea mays* L.). *Journal of Agronomy and Crop Science* **167**: 65-9.
- Sarada C, Girdhar K and Reddy T V. 2008. Effect of bio-regulators and their time of application on growth and yield of coriander (*Coriandrum sativum* L.). *Journal of Spices and Aromatic Crops* **17**(2): 183-6.



Integrated nutrient management induced changes in nutrient uptake, fruit yield and quality of Kinnow mandarin

MANISH BAKSHI¹, V K WALI², PARSHANT BAKSHI³, AKASH SHARMA⁴, DIVYA SHARMA⁵ and RAFIQ AHMAD SHAH⁶

Sher-e-Kashmir University of Agricultural Science and Technology-Jammu, Faculty of Agriculture, Chatha, Jammu, J&K 180 009

Received: 10 June 2016; Accepted: 7 September 2016

ABSTRACT

The present study on the integrated nutrient management in Kinnow mandarin involved application of inorganic fertilizers, vermicompost and *Azotobacter* under different combinations. The pooled data analysis indicated that highest concentration of leaf and fruit N (2.57 and 0.06%), Ca (4.31 and 0.033%) and Mg (0.46 and 0.024%) with the combined application of *Azotobacter* + 25% nitrogen as vermicompost and 75% nitrogen as urea. Maximum leaf and fruit P (0.19 and 0.025%) were recorded with the application of vermicompost (to supply 50% N) and urea (to supply 50% N) augmented with *Azotobacter*. While highest leaf and fruit K (1.56 and 0.092%) were recorded with full dose of nitrogen applied as vermicompost along with *Azotobacter*. Maximum *Azotobacter* counts (31.6×10^6 cfu g soil), bacterial counts (28.9×10^6 cfu g soil) and fungal counts (35.5×10^5 cfu g soil) were recorded with the application of full dose of nitrogen through vermicompost along with *Azotobacter*. The results suggested that 25% nitrogen can be replaced through chemical fertilization along with vermicompost on N- equivalent basis plus *Azotobacter* inoculation.

Keywords: *Azotobacter*, Integrated nutrient management, Kinnow, Leaf nutrients, Quality, Yield

Kinnow mandarin is highly nutrient responsive and a balanced nutrition programme is mandatory to maintain a sustained productive life of orchard in addition to quality production (Ghosh 1990). There has been a surge of interest to adopt certain measures such as Integrated Nutrient Management (INM) for making its culture eco-friendly by reducing the tradition of inorganic fertilizers, pesticides and other synthetic formulations. Use of inorganic fertilizers along with organic manures and biofertilizers is a proven technology to build up the fertility status of the soil (Srivastava and Ngullie 2009a, Srivastava and Singh 2009b). Organic manure like vermicompost is rich in plant nutrients compared to other organic manures in respect of supply of N, P and K (Srivastava *et al.* 2001, Srivastava and Singh 2009c). Vermicompost application is one of the effective methods to rejuvenate the depleted soil fertility and enrich the available pool of nutrients, maintain soil quality and conserve more biological resources (Srivastava *et al.* 2001). On the other hand, biofertilizers are preparations containing living cells

or latent cells of efficient strains of microorganisms that help crop plants uptake of nutrients by their interactions in the rhizosphere when applied through seed or to soil (Srivastava and Ngullie 2009a). *Azotobacter*, a free living microbe, acts as plant growth promoting rhizobacteria (PGPR) in the rhizosphere of almost all crops. (Gomare *et al.* 2013). *Azotobacter* produces many growth regulators such as IAA and GA which positively influence plant growth (Sharma and Kumar 2008). However, soil application of nutrients does not provide a clear picture of nutrient status of plant (Srivastava *et al.* 2001). In fruit crops like citrus, leaves have been found to be practically sensitive and convenient index of the nutrient status of the plant (Srivastava *et al.* 2008). Lot of research is being done on use of integrated supply of nutrients based on soil fertility status but information on leaf nutrient uptake and its correlation with yield and quality parameters is lacking (Marathe *et al.* 2012). Hence, the study was carried out to find the most suitable combination of inorganic fertilizers, organic manure and biofertilizer and its possible effects on nutrient status, yield, quality and soil microbial populations in Kinnow mandarin.

MATERIALS AND METHODS

This study was conducted on six year old Kinnow plants grafted on rough lemon rootstock having uniform shape, size and vigour. The experiment was laid out in the research orchard of Division of Fruit Science, Faculty of

¹Research Associate (e mail: manish.bakshi78@gmail.com), Division of Fruit Science, ²Professor and Head (e mail: vkwali@gmail.com), ³Associate Professor (e mail: bakshi_parshant@rediffmail.com), ⁴Assistant Professor (e mail: akashskuastj@gmail.com) ⁵Research Associate (e mail: divyabakshi300@gmail.com) Division of Vegetable Science, ⁶Research Scholar (e mail: rafiq_masoodi@rediffmail.com).

Agriculture, Udheywalla, SKUAST-Jammu during 2013 and 2014 in the sub-tropical zone at latitude of 32.43° North and longitude of 74.54° East. The experimental soil had pH 7.09, sand 65.5%, silt 18.5%, clay 17.0%, organic carbon 0.53%, electrical conductivity 0.14 ds/m, available nitrogen 222.5 kg/ha, available phosphorus 13.45 kg/ha, available potassium 140.5 kg/ha, available calcium 6.84 meq/100g, available magnesium 2.82 meq/100g. Recommended dose of NPK as per Package of practices for horticulture crops-SKUAST-J for Kinnow was maintained in the experiment, where only nitrogen was manipulated through different sources of fertilization. A total of 12 treatments replicated thrice were evaluated in a randomized block design: T₁: 100% N as urea, T₂: 25% N as vermicompost and 75% N as urea, T₃: 50 % N as vermicompost and 50% N as urea, T₄: 75% N as vermicompost and 25% N as urea, T₅: *Azotobacter* + 100% N as urea, T₆: *Azotobacter* + 25% N as vermicompost and 75% N as urea, T₇: *Azotobacter* + 50% N as vermicompost and 50% N as urea, T₈: *Azotobacter* + 75% N as vermicompost and 25% N as urea, T₉: *Azotobacter* + 100% N as vermicompost, T₁₀: 100% N as vermicompost, T₁₁: *Azotobacter* application and T₁₂: Absolute control.

The treatment wise application of vermicompost was applied to the trees around the trunk in the first week of January along with phosphatic fertilizers. Whole potash and first half dose of urea was applied 15 days before flowering and remaining half dose of urea was applied after fruit set. The biofertilizer *Azotobacter* was supplied with a uniform dose of 100 g / plant mixed with 10% jaggery solution prepared separately for each tree and was fed to the roots after one month of chemical fertilization.

Leaf samples were collected from seven months old spring cycle leaves on non fruiting terminals in the month of July (Chahill *et al.* 1988, Srivastava *et al.* 1994, Srivastava *et al.* 1999), thoroughly washed and ground to have a homogenous sample as per method described by Chapman (1964). Nitrogen was estimated by Micro-Kjeldahl's method as suggested by Jackson (1973). Phosphorus was determined by Vanado-molybdophosphoric acid yellow colour method (Jackson, 1973). Potassium was estimated using flame photometer and Ca, Mg was measured using atomic absorption spectrophotometer. Composite soil sample from the Kinnow rhizosphere of each treatment was used for estimation of soil microbial population following serial dilution method using specific media for each microbe (Black *et al.* 1965). The pooled mean was generated during the course of study were subjected to statistical analysis and were tested on SPSS (Version 16) software using Duncan's multiple-ranged test to identify the homogeneous type of the data sets among different treatments.

RESULTS AND DISCUSSION

Leaf nutrient content

The effect of integrated sources of nitrogen on Kinnow showed significant differences among all the

macro nutrients and are presented in Table 1 where, highest leaf nitrogen content (2.57%) was recorded in T₅ (*Azotobacter* + 100% N as urea) and T₆ (*Azotobacter* + 25% N as vermicompost and 75% N as urea). Leaf phosphorus reached to a highest level of 0.19% under T₇ (*Azotobacter* + 50% N as vermicompost and 50% N as urea) followed by T₈ (*Azotobacter* + 75% N as vermicompost and 25% N as urea) where leaf phosphorus 0.18% was obtained. T₉ (*Azotobacter* + 100% N as vermicompost) registered highest leaf potassium (1.56%) while, lowest leaf potassium content was obtained under control. Maximum leaf calcium (4.31%) was recorded in T₆ (25% nitrogen as vermicompost + 75% nitrogen as urea + *Azotobacter*) followed by plants receiving 50% nitrogen as vermicompost and 50% nitrogen as urea + *Azotobacter* (T₇). Treatment T₆ (25% nitrogen as vermicompost + 75% nitrogen as urea + *Azotobacter*) and T₇ (*Azotobacter* + 50% N as vermicompost and 50% N as urea) registered highest leaf magnesium content (0.46% each) followed by 0.45% under T₅ (100% nitrogen as urea in combination with *Azotobacter*). Minimum values for leaf macro nutrients were recorded under control. This may be due to the production of enzymatic complexes by nitrogen fixers which might have solubilized the unavailable form of nutrient elements and render them available. Increase in nitrogen content of the leaves with urea application has also been reported by Lal *et al.* (2000) in guava. The increase in phosphorus uptake might have been due to the better availability and translocation of phosphorus under *Azotobacter* application. Leaf nitrogen, phosphorus and potassium content of Kinnow mandarin were high when vermicompost was added as reported by Yadav *et al.* (2013) due to addition of organic manure in guava. Increase in foliar potassium content obtained in the present study is in conformity with the findings of Singh and Sharma (1993) in case of sweet orange cv. Mosambi.

Table 1 Leaf macronutrient composition of Kinnow mandarin as affected by different INM treatments (Pooled mean of two years)

Treatment	Nitrogen (%)	Phosphorus (%)	Potassium (%)	Calcium (%)	Magnesium (%)
T ₁	2.51 ^g	0.13 ^c	1.47 ^c	4.25 ^f	0.42 ^d
T ₂	2.51 ^g	0.13 ^c	1.50 ^e	4.28 ^{fg}	0.43 ^{de}
T ₃	2.47 ^f	0.15 ^d	1.51 ^f	4.27 ^f	0.42 ^d
T ₄	2.38 ^d	0.15 ^{de}	1.53 ^g	4.09 ^c	0.39 ^c
T ₅	2.57 ⁱ	0.16 ^f	1.48 ^d	4.29 ^{ghi}	0.45 ^{ef}
T ₆	2.57 ⁱ	0.17 ^g	1.52 ^{fg}	4.31 ⁱ	0.46 ^f
T ₇	2.54 ^h	0.19 ^h	1.54 ^h	4.30 ^{hi}	0.46 ^f
T ₈	2.42 ^e	0.18 ^g	1.55 ^{hi}	4.21 ^e	0.42 ^d
T ₉	2.39 ^d	0.15 ^e	1.56 ⁱ	4.18 ^e	0.42 ^d
T ₁₀	2.35 ^c	0.15 ^d	1.48 ^d	4.12 ^d	0.37 ^c
T ₁₁	2.33 ^b	0.12 ^b	1.45 ^b	3.79 ^b	0.34 ^b
T ₁₂	2.25 ^a	0.09 ^a	1.42 ^a	3.42 ^a	0.24 ^a

Fruit nutrient content

Data regarding effect of different INM treatments on fruit nutrient content is presented in Table 2. Highest fruit nitrogen content (0.06%) was achieved under T₅ (100% nitrogen as urea in combination with *Azotobacter*) and T₆ (25% nitrogen as vermicompost + 75% nitrogen as urea + *Azotobacter*) followed by T₁ (100% nitrogen as urea), T₂ (25% nitrogen as vermicompost + 75% nitrogen as urea) and T₇ (50% nitrogen as vermicompost and 50% nitrogen as urea + *Azotobacter*) where 0.05% fruit nitrogen was recorded. Fruit phosphorus reached to a highest level of 0.025% under T₇. Highest fruit potassium (0.092%) was recorded under T₉ (100% nitrogen as vermicompost + *Azotobacter*). Calcium content (0.033%) was highest under T₆ (25% nitrogen as vermicompost + 75% nitrogen as urea + *Azotobacter*) and T₇ (50% nitrogen as vermicompost and 50% nitrogen as urea + *Azotobacter*). Magnesium content in fruits did not show much significant difference among different treatments and ranged from 0.018 to 0.024%, lowest being observed in control (T₁₂). The above results are in line with the studies of Rana (2001) who found increase in nitrogen, potassium and calcium content of strawberry due to nitrogen and *Azotobacter* application and also reported no effect of various nitrogen fixers and urea on magnesium content of the berries. The findings on leaf and fruit nutrient content with the integrated application of inorganic fertilizers, organic manures and biofertilizers are in conformity with those reported in guava (Sharma *et al.* 2013, 2016) and Nagpur mandarin (Srivastava *et al.* 2008, 2015).

Correlation of macronutrients with yield and quality parameters

The correlation coefficient values (Table 3) indicated that fruit yield was significantly correlated with all the macronutrients in leaf. Highest significant positive correlation was observed with N (r = 0.925) followed by

Table 2 Fruit nutrient composition of Kinnow mandarin as affected by different INM treatments (pooled mean of two years)

Treatment	Nitrogen (%)	Phosphorus (%)	Potassium (%)	Calcium (%)	Magnesium (%)
T ₁	0.05 ^c	0.021 ^c	0.072 ^b	0.029 ^g	0.022 ^c
T ₂	0.05 ^{de}	0.022 ^a	0.074 ^c	0.030 ^h	0.022 ^c
T ₃	0.04 ^c	0.023 ^e	0.078 ^d	0.032 ^{ef}	0.022 ^c
T ₄	0.04 ^c	0.022 ^c	0.079 ^d	0.028 ^{fg}	0.022 ^c
T ₅	0.06 ^g	0.023 ^g	0.083 ^e	0.032 ^g	0.024 ^d
T ₆	0.06 ^f	0.023 ^c	0.086 ^f	0.033 ^{cd}	0.024 ^d
T ₇	0.05 ^e	0.025 ^d	0.088 ^f	0.033 ^h	0.024 ^d
T ₈	0.04 ^d	0.023 ^d	0.091 ^g	0.031 ^{de}	0.021 ^b
T ₉	0.04 ^c	0.022 ^f	0.092 ^g	0.029 ^{fg}	0.024 ^d
T ₁₀	0.03 ^{bc}	0.019 ^g	0.081 ^e	0.028 ^{bc}	0.022 ^c
T ₁₁	0.03 ^b	0.018 ^f	0.070 ^b	0.027 ^b	0.022 ^c
T ₁₂	0.02 ^a	0.016 ^b	0.060 ^a	0.024 ^a	0.018 ^a

Table 3 Correlation of leaf nutrient content with yield and quality of kinnow mandarin

Leaf nutrient	Coefficient of correlation (r) values				
	Yield	Quality parameter			
		Juice content	TSS	Total sugars	Ascorbic acid content
N	0.925**	0.966**	0.679**	0.605*	0.365
P	0.558*	0.559*	0.927**	0.946**	0.727**
K	0.268	0.342	0.793**	0.860**	0.778**
Ca	0.677**	0.837**	0.893**	0.848**	0.740**
Mg	0.759**	0.852**	0.900**	0.855**	0.722**

* and ** Significant at 5% and 1% level, respectively.

Mg (r = 0.759), Ca (r = 0.677) and P (r = 0.558). These findings are in conformity with the works of Marathe *et al.* (2012) in sweet orange. Trivedi *et al.* (2012) observed that incorporation of vermicompost and biofertilizers resulted in the maximum uptake of nitrogen and higher fruit yield in guava cv. Sardar. Highest degree of correlation was observed in case of juice content % with N (r = 0.966) followed by Mg (r = 0.852), Ca (r = 0.837) and P (r = 0.559). Srivastava *et al.* (2001) reported significant correlation of % juice with leaf N levels. Total soluble solids showed a strong positive correlation with P (r = 0.927) followed by Mg (r = 0.900), Ca (r = 0.893), K (r = 0.793) and N (r = 0.679). Total sugar content in fruits had a positive correlation (r = 0.946) with leaf P content followed by K (r = 0.860) and Mg (r = 0.855). Ascorbic acid content was correlated with leaf K (r = 0.778) followed by Ca (r = 0.740), P (r = 0.727) and Mg (r = 0.722). Such information would lead to find out the multiple nutrient deficiencies to manage them through site specific nutrient management (Srivastava and Singh 2015)

Soil microbial population

Soil microbial populations showed significant differences under different integrated nutrient management




Fig 1 Effect of urea, vermicompost and *Azotobacter* treatments on *Azotobacter* population (x 10⁶ cfu per gram soil) in rhizosphere of Kinnow mandarin (pooled mean of two years)




Fig 2 Effect of urea, vermicompost and *Azotobacter* treatments on bacterial population ($\times 10^6$ cfu per gram soil) in rhizosphere of Kinnow mandarin (pooled mean of two years)

treatments (Fig 1, 2 and 3). Maximum *Azotobacter* population (31.6×10^6 cfu per gram soil), bacterial population (28.9×10^6 cfu per gram soil) and fungal population (35.5×10^5 cfu/g soil) was observed in T₉ (100% nitrogen through vermicompost + *Azotobacter*). Panigrahi and Behera (1993) also observed higher *Azotobacter* population at lower level of nitrogen which is in conformity with the present finding. It has been observed that organic matter serves as energy source for growth and multiplication of *Azotobacter*. It was observed that increased nitrogen application reduced microbial population. Results are in agreement with the findings of Sharma *et al.* (2010, 2013) on apricot and Huchche *et al.* (1998).

From the present studies, it is evident that integrated nutrient management system standardized the schedule of manure and fertilizer application in Kinnow mandarin for sustaining the soil fertility to enhance the production potential and reduce the requirement of chemical fertilizers. The findings have clearly indicated that there was a positive effect of fertilizers when N was manipulated through different sources, viz. vermicompost, biofertilizers and inorganic fertilizers. The results also indicated that there was substantial improvement in leaf and fruit nutrient status of Kinnow mandarin through use of integrated nutrient management system comprising inorganic fertilizers, vermicompost and *Azotobacter*. The nutrient content in leaf further supplemented via significant correlation with fruit yield and quality parameters. Also integration of organic manure along with *Azotobacter* increased soil health in terms of microbial populations. Based on the experimental results obtained, it may finally be concluded that application of 25 % N applied through urea could be replaced with vermicompost application of nutrient equivalent basis along with inoculation of *Azotobacter* @100 g/tree for increasing the nutrient status of Kinnow mandarin. At the same time, integrated nutrient management could effectively address the environmental and sustainability concerns without losing out on productivity.

REFERENCES

Anonymous. 2014. *Indian Horticulture Database*. NHB, Ministry of Agriculture, Government. of India, Gurgaon.

- Black C A, Evans D D, Whit J L, Ensminger L E and Clark F E. 1965. *Methods of Soil Analysis*, Part (I & II). Agronomy 9, American Society of Agronomy, Madison, Washington, USA.
- Chahill B S, Dhatt A S and Singh R. 1988. Standardization of foliar sampling technique in kinnow mandarin. *Punjab Journal of Horticulture* **28**: 118–24.
- Chapman H D. 1964. Suggested foliar sampling and handling techniques for determining nutrient status of some field, horticultural and plantation crops. *Indian Journal of Horticulture* **21**: 98–118.
- Ghosh S N. 1990. Nutritional requirement of sweet orange (*Citrus sinensis* Osbeck) cv. Mosambi. *Haryana Journal of Horticultural Sciences* **19**: 39–44.
- Gomare K S, Mese M and Shetkar Y. 2013. Isolation of *Azotobacter* and cost effective production of biofertilizer. *Biotechnology* **3**(5): 54–6.
- Huchche A D, Ladaniya M S, Lallan Ram, Kohli R R and Srivastava A K. 1998. Effect of nitrogenous fertilizer and farmyard manure on yield and shelf-life of Nagpur mandarin. *Indian Journal of Horticulture* **55**(2):108–12.
- Jackson M L. 1973. *Soil Chemical Analysis*, p 498. Prentice Hall of India Pvt Ltd New Delhi. Lal G, Sen N L and Jat R G. 2000. Yield and leaf nutrient composition of guava as influenced by nutrients. *Indian Journal of Horticulture* **57**(2): 130–2.
- Marathe R A, Bharambe P R, Sharma R and Sharma U C. 2012. Leaf nutrient composition, its correlation with yield and quality of sweet orange and soil microbial population as influenced by INM in vertisol of central India. *Indian Journal of Horticulture* **69**(3): 317–21.
- Panigrahi U C and Behera B. 1993. Response of *Azotobacter* inoculants on total nitrogen, organic carbon and microbial population of soil and yield of potato. *Indian Journal of Agricultural Chemistry* **26**(1): 17–23.
- Sharma A, Wali V K, Bakshi P and Jasrotia A. 2013. Effect of integrated nutrient management strategies on nutrient status, yield and quality of guava. *Indian Journal of Horticulture* **70**(3): 333–9
- Sharma A, Wali V K, Bakshi P, Sharma V, Sharma V, Bakshi M and Rani S. 2016. Impact of poultry manure on fruit quality attributes and nutrient status of guava (*Psidium guajava* L.) cv. L-49 plant. *Indian Journal of Agricultural Sciences* **86**(4): 533–40.
- Sharma S D and Kumar P. 2008. Relationship of arbuscularmycorrhizal fungi and *Azotobacter* with plant growth, fruit yield, soil and leaf nutrient status of mango orchards in North-Western Himalayan region of India. *Journal of Applied Horticulture* **10**(2): 172–8.
- Singh C and Sharma B B. 1993. Leaf nutrient composition of sweet orange as affected by combined use of bio and chemical fertilizers. *South Indian Horticulture* **41**(3): 131–4.
- Singh V J, Sharma S D, Kumar P, Bhardwaj S K and Raj H. 2010. Conjoint application of bio-organic and inorganic nutrient sources for improving cropping behavior soil properties and quality attributes of apricot (*Prunus armeniaca*). *Indian Journal of Agricultural Sciences* **80**(11): 981–7.
- Srivastava A K, Kohli R R, Dass H C, Huchche A D and Lallan Ram. 1999. Evaluation of the nutritional status of Nagpur mandarin (*Citrus reticulata* Blanco) by foliar sampling. *Journal of Tropical Agriculture* **72**(6): 93–8.
- Srivastava A K, Lallan Ram, Huchche A D, Kohli R R and H C Dass. 1994. Standardisation of leaf sampling technique in Nagpur mandarin under sub-humid tropical climate. *Indian*

- Journal of Horticulture* **51**(1): 32–6.
- Srivastava A K and Ngullie E. 2009a. Integrated nutrient management: Theory and practice. *Dynamic Soil, Dynamic Plant* **3**(1): 1–30.
- Srivastava A K and Singh Shyam. 2004. Zinc nutrition, a global concern for sustainable citrus production. *Journal of Sustainable Agriculture* **25**(3): 5–42.
- Srivastava A K and Singh Shyam. 2009c. Zinc nutrition in Nagpur mandarin on haplustert. *Journal of Plant Nutrition* **32**(7): 1 065–81.
- Srivastava A K and Singh Shyam. 2015. Site-Specific nutrient management in Nagpur mandarin (*Citrus reticulata* Blanco) raised on contrasting soil types. *Communication in Soil Science and Plant Analysis* **47**(3): 447–56.
- Srivastava A K, Singh Shyam and Albrigo L G. 2008. Diagnosis and remediation of nutrient constraints in Citrus. *Horticultural Reviews* **34**: 277–64.
- Srivastava A K, Singh Shyam and Huchche A D. 2015. Evaluation of INM in citrus on Vertic Ustochrept: Biometric response and soil health. *Journal of Plant Nutrition* **38**(5): 1–15.
- Srivastava A K, Singh Shyam, Huchche A D and Lallan Ram. 2001. Yield based leaf and soil test interpretations for Nagpur mandarin in Central India. *Communication in Soil Science and Plant Analysis USA* **32** (3 and 4): 585–99.
- Trivedi Y V, Patel N L, Ahlawat T R, Gaikwad S S and Bhalerao P P. 2012. Impact of organic manures and inorganic fertilizers on growth, yield, nutrient uptake and soil nutrient status in guava. *Indian Journal of Horticulture* **69**(4): 501–6.
- Yadav R I, Singh R K, Jat A L and Choudhary H R. 2013. Effect of nutrient management through organic sources on productivity and profitability of guava (*Psidium guajava* L.) under Vindhyan region. *Environment and Ecology* **31**(2A): 735–7.



Effect of plant density and foliar nutrient application on seed yield and quality in Asiatic carrot (*Daucus carota*)

C MANIMURUGAN¹, V K PANDITA², B S TOMAR³, N GUPTA⁴ and P M SINGH⁵

ICAR- Indian Agricultural Research Institute, New Delhi 110 012

Received: 22 July 2016; Accepted: 29 August 2016

ABSTRACT

A field experiment was conducted to study effect of plant density and foliar spray of nutrients on carrot (*Daucus carota* L.) seed yield and quality during winter season of 2012-13 and 2013-14. Plant height, days to 50% flowering, number of umbels/plant, number of seeds/umbel, seed yield/umbel, percentage contribution of different order umbel to total seed yield, seed yield/plant, seed yield/ha, 1 000 seed (mericarp) weight, germination (%) and vigour index were recorded. Significant highest average plant height, earliness in days to 50% flowering and seed yield/ha (12.9 q) were observed in high plant density (13 plants/m²). More number of secondary (11.6) and tertiary umbels (30.1) and higher seed yield/plant (25.2g) were recorded in low plant density (4 plants/m²). Per cent contribution of primary umbel to total seed yield was maximum (41.76%) in 13 plants/m² and minimum (16.89%) in 4 plants/m². But an increasing trend in contribution percentage to total seed yield by secondary and tertiary umbels was recorded from 13 plants/m² to 4 plants/m². Seeds from primary and secondary umbels recorded high quality in plant density of 5 plants/m² and 4 plants/m² respectively. Comparative reduction in pooled seed vigour was observed in 7 plants/m² and 13 plants/m². Foliar spray of 0.1% MgSO₄ at 30 and 60 days after transplanting (DAT) increase the seed quality. Both seed yield and quality increase was observed in 0.1% borax spray at 30 and 60 DAT.

Key words: Borax, Carrot, Foliar nutrient spray, MgSO₄, Plant density, Seed quality, Seed yield

Carrot (*Daucus carota* L.) is one of the important root vegetable in umbelliferae family. It covers an area of 62.4 thousand ha in India with the production of 10.7 lakh tonnes (Saxena and Gandhi 2015). Both Asiatic and temperate type carrots are cultivated in India but they require different climatic condition for their seed production. Seeds of Asiatic type can be produced in plains but that of temperate types in hilly region only. Carrot inflorescence consists of many branches and is divided into different order umbels (primary, secondary and tertiary etc.) based on their position and developmental stages. This also leads to production of seeds in different maturity levels. When the primary umbel reaches maturity the secondary or tertiary umbels may be in immature or flowering stage, respectively. This is one of major problem in seed production of carrot crop. Highest contribution by primary umbel to total seed yield in high density planting in temperate carrot was demonstrated (10 to 80 plants/m², 2-84 plants/m² and 2-100 plants/m²) by

many authors (Gray *et al.* 1983, Gray and Steckel 1983 and Merfield *et al.* 2010). High quality seeds are generally produced from primary and secondary umbels in carrot (Pandita *et al.* 2005). This indicates promotion of primary and secondary umbel by increasing plant density may promote uniform maturity and quality of seeds. But there is a need of fine balance between increase in plant density, increase in seed yield and maintenance of seed quality. One should not compromise seed quality for higher seed yield by increasing plant density. The optimum plant density also depends upon environment and cultivars. Apart from plant density, balanced nutrition of macro and micronutrients also play a vital role in increase and enhancement of seed yield and quality. In this study an attempt was made to suppress the side branches of inflorescence by increasing the plant density in carrot and to know the supportive effect of foliar nutrient for obtaining higher seed yield and quality in carrot.

MATERIALS AND METHODS

The experiment was conducted at Seed Production Unit of Indian Agricultural Research Institute, New Delhi during the period of September to May, 2012-13 and 2013-14. The experimental site was situated at 28°35'N latitude & 77°12'E longitude and at an altitude of 228.6 m above mean sea level. Seeds of carrot cv. Pusa Rudhira were sown in the last week of September and transplanting of uniform size seedlings was done in third

^{1,4}Research Scholars, Division of Seed Science and Technology, ³Head (email: bst_spu_iari@rediffmail.com), Division of Vegetable Science, ICAR-Indian Agricultural Research Institute, New Delhi 110 012. ²Head (I/c) (email: vndpandita@yahoo.com), ICAR-IARI, Regional station, Karnal 132 001, ⁵Head (I/c), Division of Crop Improvement, ICAR-Indian Institute of Vegetable Research, Varanasi 221 305.

week of January. Split plot design was adopted to conduct experiments with plant density as main plot treatment and foliar spray of nutrients as sub-plot treatments. Stecklings were transplanted with spacing of 15×50cm, 30×50cm, 40×50cm and 50×50cm which gave plant density of D₁-13 plants/m², D₂-7 plants/m², D₃-5 plants/m² and D₄-4 plants/m² respectively. Sub-plot size was 3m×3m and main plot consists of seven sub-plots for six foliar spray treatments and one control. Main plots were replicated three times with all sub-plot treatments. Foliar spray treatments consist of T₁-0.1% borax at 30 DAT, T₂- 0.1% borax at 60 DAT, T₃- 0.1% borax at 30+60 DAT, T₄- 0.1% Magnesium sulphate (MgSO₄) at 30 DAT, T₅- 0.1% MgSO₄ at 60 DAT, T₆- 0.1% MgSO₄ at 30+60 DAT and T₇- control (sprayed with water). Borax used as boron source. All the package of practices and plant protection measures were carried out as per recommendation. Days to 50% of flowering was counted from initiation of primary umbel to 50% flowering of primary, secondary and tertiary umbels in individual treatment. Plant height of 10 randomly selected plants was measured at the time of 50% of flowering. Number of umbels/plant, percentage contribution of all different order umbel(s) to total seed yield and seed yield/plant was estimated from average of observations taken from ten randomly selected plants at the time of harvest. Number of seeds/umbel and seed yield/umbel was estimated from 10 randomly selected umbels of different order. Seed yield/ha was estimated based on average of main plot and subplot seed yield. Seed quality parameters like 1 000 seed weight, germination percentage (ISTA 2012) and vigour index

(Abdul-Baki and Anderson, 1973) were calculated. Since seeds of primary and secondary umbels were combined together in general seed lot, a pooled vigour index was also calculated by taking average of seed vigour index one of primary and secondary umbels. Statistical analysis was carried out by a software AgRes version 3.01.

RESULTS AND DISCUSSION

Growth attributes

In the present study higher average plant height with respect to primary (118.7cm), secondary (133.6cm) and tertiary (139.4cm) umbels were observed in highest plant density (D₁). This may be due to more apical growth and suppression of side branches. Similar results have also been reported by Ahmad and Tanki (1997) and Kumar (2005) in carrot. In case of other crops, increasing plant density generally reduces the plant height or length due to competition among plants for space, nutrient and sunlight. Bahlgerdi *et al.* (2014) observed highest plant length in medicinal pumpkin in lowest plant density. Highest plant density (D₁) recorded earliness for days to 50% flowering as compared to other plant densities. Decreasing plant density showed delay in days to 50% flowering. Earliness in flowering may be due to competition between plants in D₁. Andriolo (1999) reported that an optimized leaf area index (LAI) can be obtained by ideal population density for better interception of maximum useful radiation for photosynthesis. Before plants start the reproductive phase, an optimized LAI can be reached very early on high density

Table 1 Effect of plant density and foliar spray on growth and flowering in carrot cv. Pusa Rudhira (Pooled data of two season 2012-13 and 2013-14)

Treatment	Plant height (cm)			Days to 50 % flowering			Number of umbels/plant	
	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary	Secondary	Tertiary
<i>plant density</i>								
D ₁	118.7 ^a	133.6 ^a	139.4 ^a	26.5 ^c	33.3 ^d	40.2 ^c	6.2 ^d	17.5 ^d
D ₂	112.7 ^b	127.7 ^b	131.5 ^b	27.3 ^b	34.1 ^c	41.5 ^b	9.3 ^c	23.8 ^c
D ₃	108.9 ^b	123.6 ^c	126.6 ^c	27.8 ^{ab}	34.5 ^b	43.0 ^a	11.3 ^b	27.5 ^b
D ₄	104.1 ^c	121.5 ^c	124.5 ^c	28.3 ^a	35.0 ^a	43.4 ^a	11.6 ^a	30.1 ^a
CD (P=0.05)	3.96	2.97	3.96	0.60	0.34	1.02	0.12	1.10
<i>Foliar nutrient spray</i>								
T ₁	110.0	126.1	129.4	27.1	33.8	41.5	9.6	24.6
T ₂	112.0	127.5	130.6	27.5	34.1	42.0	9.7	25.1
T ₃	113.3	128.6	131.5	27.8	34.6	42.4	9.7	25.7
T ₄	110.4	125.1	130.2	27.1	34.0	41.5	9.5	24.0
T ₅	110.9	126.8	130.9	27.6	34.1	42.0	9.6	24.6
T ₆	111.8	128.1	131.8	28.1	34.8	42.9	9.6	25.3
T ₇	109.2	123.9	129.1	27.1	33.8	41.8	9.5	23.8
CD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	NS
D×T	NS	NS	NS	NS	NS	NS	NS	NS

Means followed by different letters are statistically different. Separation done by Duncan's Multiple Range Test.

plantings. Thus, planting density in any crop influences plant growth, changing the flower initiation and fruit development and interfering on the biomass distribution between source and sink. Number of umbels/plant increased with decrease in plant density and was highest in D_4 for both secondary (11.6) and tertiary umbels (30.1). This may be due to availability of more nutrient and sunlight to individual plants in low plant density. These results are in accordance with Kumar (2005) in carrot. None of the growth parameters were recorded significant difference with respect of foliar nutrient sprays. No significant interaction effect was observed between density and foliar nutrient spray for all growth attributes (Table 1).

Seed yield attributes

Highest number of seeds/primary umbel was observed in D_2 (1742) and D_3 (1725) followed by D_1 (1700) and D_4 (1702). This indicates both reduction and increase in plant density affect the seed number in primary umbel. Lowest plant density (D_4) recorded highest number of seeds in secondary and tertiary umbel. Similar results were observed in onion by Asaduzzaman *et al.*, (2012). Seed yield/primary umbel was recorded higher and on par in D_2 (4.24g), D_3 (4.30g) and D_4 (4.22g) than D_1 (4.03). Seed yield/secondary and tertiary umbels were highest in lowest plant density (D_4).

Percentage contribution by primary umbel (41.73%) to total seed yield was observed highest in high plant density (D_1). But D_3 and D_4 received significantly highest contribution from secondary umbels because increased number of secondary umbel in low plant density. Contribution of tertiary umbels to total seed yield (10.70%) was highest in lowest plant density (D_4). None of the foliar spray recorded significant differences with respect to percentage contribution in primary and secondary umbels. Seed yield/plant was highest (25.2g) in lowest plant density (D_4). But seed yield/ha was significantly higher (12.9q) in highest plant density (D_1). It may be due to increased plant population in high plant density (D_1). These results are similar with results obtained by Guerrero *et al.* (1986), Amjad and Anjum (2001), Lima *et al.* (2003) and Pandita *et al.* (2005) in temperate and tropical carrot cultivars.

Recent studies in watermelon (Edelstein and Nerson 2002), muskmelon (Nerson 2002), onion (Kanwar *et al.* 2000) and medicinal pumpkin (Bahlgerdi *et al.* 2014) showed that relatively high plant densities were required to obtain the highest seed yield. In contrary, there are few reports in seed production of winter squash and squash suggested low plant densities for higher seed yield (Dematte *et al.* 1978, Edelstein *et al.* 1985, Lima *et al.* 2003). Al Mamun *et al.*, (2016) also reported that early planting of potato top shoot cutting with closer spacing (50×10 cm and 50×15cm) was most suitable for breeder seed multiplication. So optimization of plant density for seed production of any crop and their different cultivars are very important to obtain higher seed yield and seed quality.

Among the foliar sprays, T_3 recorded higher number

of seeds, seed yield/umbel, seed yield/plant (20g) and seed yield/ha (12.3q). However, T_6 was on par for number of seeds and seed yield/primary umbel with T_3 . These results are in accordance with results obtained by Sharma *et al.* (1999), Vratarić *et al.* (2006) and Kumar *et al.*, (2012). For all seed yield attribute, significant interaction between density and foliar spray was not observed (Table 2).

Seed quality attributes

Highest 1000 seed (mericarp) weight with respect to primary umbel was recorded in D_3 (2.489g) followed by D_2 (2.446g) and D_4 (2.476g). Many authors reported highest 1000 seed weight of carrot in less plant density (Kumar 2005). In our study maximum seed weight was observed in D_3 instead of D_4 . Higher number secondary and tertiary umbel in D_4 leads to slight quality reduction in seeds form primary umbel. Though, only primary umbels promoted in D_1 , the reduced seed weight indicate the competition among the plants for various resources.

Similar kind of results was observed in *Cucurbita pepo*. L (Loy 1988) and medicinal pumpkin (Bahlgerdi *et al.* 2014). These results indicated requirement of optimum level of plant density for quality seed production instead of high or less plant density. Plant density had no significant effect on germination per cent of seeds from primary umbels. Seeds of secondary and tertiary umbels recorded highest germination at lowest plant density (D_4). Highest significant vigour index with respect to primary umbel was recorded in D_3 (1134) followed by D_2 (1077) and D_4 (1087). An increasing trend of vigour index was observed for seeds of secondary and tertiary umbel from highest (D_1) to lowest plant density (D_4). Foliar spray treatment T_3 (2.472g) had maximum test weight from primary umbels and was on par with T_2 (2.457g) and T_6 (2.458g), 1000 seed weight was observed highest with T_3 in secondary and tertiary umbel. Similar results were also observed in chilli (Dongre *et al.* 2000) and Niger (Paikray, *et al.*, 2001).

Among the foliar treatments, both T_3 and T_6 recorded significantly highest germination percentage in seeds from primary, secondary and tertiary umbel. T_3 followed by T_6 recorded highest vigour index for seeds of all order umbels. The following may be the reason for seed quality enhancement by T_6 . Magnesium is central element in chlorophyll. It participates in activity of many enzymes and involved in phosphorous translocation (Curley 1994). Foliar 'boosting' of Mg through leaves may deliver the Carbon and Nitrogen containing compounds to the seeds which effectively increases the seed quality (Gerendas and Fuhrs 2013). Quality of seeds form primary umbel was always higher than the other order umbel in all the density and foliar treatments. Highest pooled vigour index was observed in D_3 (1082), D_4 (1071), T_3 (1102) and T_6 (1089). This indicates, less plant density and spraying of foliar nutrient was promoting the seed quality in carrot. Interaction effect of density and foliar nutrient spray was not significant for all the seed quality parameters. Germination per cent of seeds from tertiary umbel was always less than seed certification

Table 2 Effect of plant density and foliar spray on seed yield parameters in carrot cv. Pusa Rudhir (Pooled data of two season 2012-13 and 2013-14)

Treatments	Number of seeds/umbel			Seed yield/umbel (g)			Percentage contribution to total seed yield			Seed yield/ plant (g)	Seed yield/ ha (q)
	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary		
<i>Plant density</i>											
D ₁	1700 ^b	389 ^c	23 ^d	4.03 ^b	0.80 ^d	0.039 ^d	41.73 ^a	51.17 ^c	7.11 ^c	9.7 ^d	12.9 ^a
D ₂	1742 ^a	609 ^b	27 ^c	4.24 ^a	1.33 ^c	0.045 ^c	24.05 ^b	69.94 ^b	6.01 ^d	17.7 ^c	11.8 ^b
D ₃	1725 ^a	632 ^b	37 ^b	4.30 ^a	1.39 ^b	0.064 ^b	19.83 ^c	72.10 ^a	8.07 ^b	21.7 ^b	10.9 ^c
D ₄	1702 ^b	692 ^a	48 ^a	4.22 ^a	1.57 ^a	0.089 ^a	16.89 ^d	72.41 ^a	10.70 ^a	25.2 ^a	10.3 ^d
CD (P=0.05)	21.71	26.11	2.74	0.09	0.03	0.002	0.58	0.74	0.34	0.29	0.23
<i>Foliar nutrient spray</i>											
T ₁	1688 ^c	577 ^{bc}	32 ^c	4.14 ^{cd}	1.25 ^{bc}	0.059 ^{bc}	25.78	66.27	7.95 ^{abc}	18.3 ^{cd}	11.3 ^{bce}
T ₂	1730 ^b	592 ^b	36 ^b	4.25 ^b	1.29 ^b	0.062 ^{ab}	25.52	66.18	8.30 ^{ab}	18.9 ^b	11.5 ^b
T ₃	1793 ^a	621 ^a	40 ^a	4.40 ^a	1.36 ^a	0.066 ^a	24.74	66.74	8.52 ^a	20.0 ^a	12.3 ^a
T ₄	1670 ^c	556 ^c	30 ^{cd}	4.07 ^d	1.23 ^{cd}	0.055 ^c	25.98	66.53	7.49 ^c	17.9 ^{de}	11.2 ^{ce}
T ₅	1723 ^b	572 ^{bc}	32 ^c	4.21 ^{bc}	1.27 ^{bc}	0.058 ^{bcd}	25.84	66.36	7.80 ^{bc}	18.5 ^{bc}	11.4 ^{bce}
T ₆	1772 ^a	588 ^b	35 ^b	4.35 ^a	1.30 ^b	0.062 ^{ab}	25.72	65.96	8.32 ^{ab}	19.1 ^b	11.5 ^{bc}
T ₇	1644 ^d	557 ^c	28 ^d	3.95 ^e	1.20 ^d	0.053 ^d	25.78	66.79	7.44 ^c	17.3 ^e	11.1 ^e
CD (P=0.05)	26.53	24.29	2.40	0.09	0.05	0.004	NS	NS	0.71	0.60	0.30
D×T	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Means followed by different letters are statistically different. Separation done by Duncan's Multiple Range Test.

Table 3 Effect of plant density and foliar spray on seed quality of different order umbels in carrot cv. Pusa Rudhira (Pooled data of two season 2012-13 and 2013-14)

Treatment	1000 mericarp weight (g)			Germination (%)			Vigour index			Pooled Vigour index
	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary	Primary	Secondary	Tertiary	
<i>Plant density</i>										
D ₁	2.357 ^c	2.089 ^d	1.778 ^c	71 (57.48)	67 (55.20) ^c	40(39.08) ^c	1024 ^c	933 ^d	507 ^d	978 ^c
D ₂	2.446 ^b	2.162 ^c	1.808 ^b	73 (58.81)	71 (57.12) ^b	42(40.15) ^{bc}	1087 ^b	990 ^c	555 ^c	1039 ^b
D ₃	2.489 ^a	2.240 ^b	1.844 ^a	75 (59.88)	72 (57.99) ^{ab}	43(41.23) ^b	1135 ^a	1030 ^b	593 ^b	1083 ^a
D ₄	2.476 ^{ab}	2.286 ^a	1.858 ^a	74 (59.12)	74 (59.00) ^a	50(44.94) ^a	1077 ^b	1064 ^a	692 ^a	1071 ^a
CD (P=0.05)	0.04	0.04	0.02	NS	1.23	1.25	45.26	22.37	34.38	18.62
<i>Foliar nutrient spray</i>										
T ₁	2.442 ^{ab}	2.188 ^b	1.816 ^{bc}	72 (57.90) ^c	70 (56.55) ^{bc}	42(40.31) ^{bc}	1055 ^{cde}	980 ^{ce}	559 ^{cd}	1018 ^{cd}
T ₂	2.457 ^a	2.211 ^{ab}	1.842 ^{ab}	73 (58.82) ^{bc}	70 (57.05) ^{bc}	44(41.61) ^{ab}	1093 ^{bc}	1014 ^b	601 ^{ab}	1054 ^b
T ₃	2.472 ^a	2.231 ^a	1.857 ^a	75 (60.15) ^{ab}	73 (58.86) ^a	46(42.86) ^a	1140 ^a	1064 ^a	642 ^a	1102 ^a
T ₄	2.425 ^{bc}	2.180 ^{bc}	1.800 ^{cd}	72 (58.27) ^{bc}	70 (56.47) ^{bc}	42(40.40) ^{bc}	1049 ^d	976 ^{ce}	546 ^d	1013 ^d
T ₅	2.445 ^{ab}	2.193 ^b	1.826 ^b	73 (58.84) ^{bc}	72 (57.78) ^{ab}	45(41.87) ^{ab}	1077 ^{cd}	1012 ^{bc}	591 ^{bc}	1045 ^{bc}
T ₆	2.458 ^a	2.210 ^{ab}	1.835 ^{ab}	76 (60.85) ^a	73 (58.62) ^a	46(42.76) ^a	1137 ^{ab}	1041 ^{ab}	630 ^{ab}	1089 ^a
T ₇	2.397 ^c	2.146 ^c	1.779 ^d	70 (56.99) ^c	69 (55.92) ^c	41(39.68) ^c	1015 ^c	944 ^e	538 ^d	980 ^e
CD (P=0.05)	0.03	0.04	0.03	1.93	1.52	1.63	43.63	45.09	41.45	30.62
D×T	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Means followed by different letters are statistically different. Separation done by Duncan's Multiple Range Test. Values in parenthesis are arcsine transformed value.

standard of 60% in all the treatments (Table 3).

Though highest seed yield/ha was observed in 13 plants/m², we recommend 5 plants/m² is ideal for breeder and foundation seed production without quality reduction. Since seeds form primary and secondary umbels of all the plant density recorded germination per cent higher than ISTA seed standard (60%) for carrot, even we can adopt 13 plants/m² for certified seed production of carrot cv. Pusa Rudhira. Harvesting seeds from tertiary umbel must be avoided to maintain higher seed quality.

REFERENCES

- Abdul-Baki A A and Anderson J D. 1973. Vigour determination in soybean seed by multiple criteria. *Crop Science* **13**: 630–3.
- Ahmad N and Tanki M I. 1997. Effect of planting date, spacing and steckling size on growth and seed yield of carrot (*Daucus carota* L.). *Haryana Journal of Horticultural Sciences* **26**: 274–6.
- Al Mamun M A, Al-Mahmud A, Zakaria M, Hosssain M M and Hossain M T. 2016. Effect of planting times and plant densities of top-shoot cuttings on multiplication of breeder seed potato. *Agricultural and Natural Resources* **50**: 26–31.
- Amjad M and Anjum M A. 2001. Effect of root size, plant spacing and umbel order on the quality of carrot seed. *International Journal of Agriculture and Biology* **3**(2): 239–42.
- Andriolo J L. 1999. Fluxo de carbono da planta. (In) *Fisiologia das culturas protegidas*, p. 13–46. Santa Maria, Editora UFSM.
- Asaduzzaman M, Mahmudul H, Mainul H and Moniruzzaman M. 2012. Quality seed production of onion (*Allium cepa* L.): an integrated approach of bulb size and plant spacing. *Jouranal of Agricultural Reserach* **50**(1): 119–28.
- Bahlgerdi M, Aroiee H and Azizi M. 2014. The study of plant density and planting methods on some growth characteristics, seed and oil yield of medicinal pumpkin (*Cucurbita pepo* var. *styriaca*, cv. 'Kaki'). *American Journal of Life Sciences* **2**(5): 319–24.
- Curley S. 1994. *Foliar Nutrition*, p 26. Midwest Laboratories, Omaha.
- Dematte M E S P, Alves S, Camargo L S and Nagai V. 1978. Seed production in *Cucurbita pepo* var. *meloepo* (summer squash) cv. Caserta IAC-1967 under 3 different spacings. *Cientifica* **6**: 165–73 (in Portuguese).
- Dongre S M, Mahorkar V K, Joshi P S and Deo D D. 2000. Effect of micronutrients spray on yield and quality of chilli (*Capsicum annum* L.) cv. Jayanti. *Agricultural Science Digest* **20**: 106–7.
- Edelstein M and Nerson H. 2002. Genotype and plant density affect watermelon grown for seed consumption. *HortScience* **37**: 981–3.
- Edelstein M, Nerson H, Paris H S, Karchi Z, Burger Y and Zohar R. 1985. Hybrid seed production in spaghetti squash (*Cucurbita pepo* L.) using ethephon and honeybees. *Hassadeh* **65**: 2 416–9 (in Hebrew).
- Gerendas J and Fuhrs H. 2013. The significance of magnesium for crop quality. *Plant and Soil* **368**(1): 101–28.
- Gray D, Steckel J R A and Ward I A. 1983. Studies on carrot seed production. Effects of plant density on yield and components of yield. *Journal of Horticultural Science* **58**(1): 83–90.
- Gray D, Steckel J R A. 1983. Seed quality in carrots: the effect of seed crop plant density, harvest date and seed grading on seed and seedling variability. *Journal of Horticultural Science* **58**: 393–401.
- Guerrero M A, Diaz A A, Heredia Z A and Salinas G J. 1986. A study of carrot seed production in El Bajío. *Proceedings of the Triopical Region- American Society of Horticultural Science* **23**: 272–4.
- ISTA. 2012. Seed Testing Rules. International Seed Testing Association, CH- Swizerland.
- Kanwar J S, Gill B S and Bal S S. 2000. Response of planting time and density to onion seed yield and quality. *Seed Research* **28**: 212–4.
- Kumar K .2005. Effect of plant density, steckling size, pruning and growth regulators on seed crop of carrot (*Daucus carota* L.). Ph D thesis, Department of Vegetable Science, College of Agriculture, Chaudhary Charan Singh Haryana Agricultural University, Hisar.
- Kumar S, Yadav M K and Yadav Y C. 2012. Effect of foliar application of boron on growth, yield, quality and seed production of Radish (*Raphanus sativus* L.) cv. Jaunpuri Local. *Progressive Agriculture* **12**(1): 199–202.
- Lima M S, Cardoso A I I and Verdial M F. 2003. Plant spacing and pollen quantity on yield and quality of squash seeds. *Horticultura Brasileira* **21**(3): 443–7.
- Loy J B. 1988. Improving seed yield in hull-less seeded strains of *Cucurbita pepo*. *Cucurbit Genetics Cooperative* **11**: 72–3.
- Merfield C N, Hampton J G, Wratten S D, Prapanoppasin P and Yeeransiri P. 2010. The effect of plant density on seed yield and quality of carrot (*Daucus carota* L.) Agronomy Society of New Zealand Special Publication No. 13, Grassland Research and Practice Series No. **14**, pp 75–83.
- Nerson H. 2002. Relationship between plant density and fruit and seed production in muskmelon. *Journal of American Society of Horticultural Science* **127**: 855–9.
- Nerson H. 2005. Effects of fruit shape and plant density on seed yield and quality of squash. *Scientia Horticulturae* **105**: 293–304.
- Paikray R K, Mishra P J, Mohapatra A K, Haldar J and Panda S. 2001) Response of niger (*Guizotia abyssinica*) to secondary and micronutrients in acid red soil. *Annals of Agricultural Research* **22**: 140–2.
- Pandita V K, Rana S C, Chaudhry D and Kumar V. 2005, Seed productivity and quality in relation to plant spacing in carrot (*Daucus carota*). *Indian Journal of Agricultural Sciences* **75**(11): 722–4.
- Saxena M and Gandhi C P. 2015. Indian horticulture database 2014, National Horticulture Board, Ministry of Agriculture, Government of India.
- Sharma S K, Singh H and Kohli U K 1999. Influence of boron and zinc on seed yield and quality in radish. *Seed Research* **27**: 154–8.
- Vrataric M, Sudaric A, Kovacevic V, Duvnjak T, Krizmanic M, Mijic A. 2006. Response of soybean to foliar fertilization with magnesium sulfate. *Cereal Research Commutation* **34**: 709–12.