



Impact of elevated carbon dioxide on the protective enzymes in brown planthopper (*Nilaparvata lugens*) and infested rice (*Oryza sativa*) plants

VEERANNA DARAVATH¹, SUBHASH CHANDER² and D SAGAR³

ICAR-Indian Agricultural Research Institute, New Delhi 110 012

Received: 17 August 2017; Accepted: 28 May 2018

ABSTRACT

The effect of elevated carbon dioxide (CO₂) on protein content, and catalase and peroxidase activity in rice (*Oryza sativa* L.) plant and brown planthopper (BPH) [*Nilaparvata lugens* (stal)] was studied in open top chamber (OTC) at elevated CO₂ (570±25 µl/l) compared to ambient CO₂ (400±25 µl/l). Uninfested rice plants under elevated CO₂ had lower protein content but increased activity of both catalase and peroxidase compared to ambient CO₂. The BPH infested rice plants had significantly lower protein content (15.89 g/l) and reduced catalase activity (0.78 µmoles of H₂O₂ decomposed/min/mg of protein) under both elevated and ambient CO₂ conditions compared to uninfested rice plant. However, the activity of peroxidase in infested rice plant was non-significantly increased at both elevated and ambient CO₂ compared to uninfested rice plants, indicating perhaps that it is an important enzyme in plant defence against BPH in addition to elevated CO₂. Further, lower protein content in rice plants under elevated CO₂ resulted in lower protein in BPH that fed on them. The activity of catalase and peroxidase in BPH was enhanced when they were fed rice plants under elevated CO₂ compared to ambient CO₂.

Key words: Catalase, Elevated CO₂, *Nilaparvata lugens*, Peroxidase, Protein, Rice

Rice (*Oryza sativa* L.) is indisputably the world's most important staple food that provides nutrition to more than half of the world's population (Khush 2004). It is attacked by several insect pests, which cause severe yield loss in one or the other region of India from nursery to harvest (Asghar *et al.* 2009). Among the sap sucking pests of rice, the planthoppers have become economically important all over the world (Magnumder *et al.* 2013). Brown planthopper (BPH) [*Nilaparvata lugens* (Stal)] (Hemiptera: Delphacidae) is one of the most destructive insect pests of rice. It causes direct damage by sucking phloem sap, causing leaf stippling, and indirect damage by transmitting viral diseases such as grassy stunt and rugged stunt (Reissig *et al.* 1986). The BPH is recognised for its ability to adapt in different ecological conditions and its regular impacts on rice yield (Lu *et al.* 2005).

Climate change is presently considered to be the most important factor influencing agricultural production. Greenhouse gases are contributing to the rise of temperature, among which carbon dioxide (CO₂) is the major gas. The atmospheric CO₂ concentration in the terrestrial troposphere was about 280 µl during pre industrial period and increased

to 398 µl/l in 2014 (NOAA/ESRL 2014). These variations might have a direct impact on plant chemical constituents and metabolic enzymes of herbivorous insects.

The impact of elevated CO₂ on ecosystem is expectable to be enormous, especially on the growth and biochemical constituents of plants (Penuelas *et al.* 2002). Foliar amino acids and protein content in cotton plants lowered significantly under elevated CO₂ compared to ambient CO₂ (Wu *et al.* 2007). Changes in atmospheric CO₂ not only affect the plant quality but also herbivore performance (Awmack *et al.* 2004). Most herbivorous insects responded to elevated CO₂ by increasing developmental time, and reducing growth, survival rates, population densities and fitness, presumably because of the increased levels of carbohydrates in host plants (Yin *et al.* 2009). The carbon: nitrogen (C: N) ratio of the plant foliage generally increases when plants are grown under elevated CO₂ than in the ambient CO₂. As a result, insect larvae increase leaf consumption and their sucking rates under elevated CO₂ to compensate for lower nitrogen in plant foliage (Prasannakumar *et al.* 2012). Besides, size of plant canopy may also influence pest populations especially rice planthoppers through changes in micro-environment. Increase in C: N ratio of plants under elevated CO₂ leads to higher feeding rates of insects that may impinge upon level of host plant resistance against the pest. *N. lugens* fed on resistant rice varieties significantly enhanced glutathione S-transferase (GST) enzyme activity (Zhou *et al.* 2003).

Plants respond to insect infestation and/ or climate

¹Scientist (e mail: veerudharavath67@gmail.com), DAATTC, Adilabad, Telangana. ²Principal Scientist and Professor (e mail: schanderthakur@gmail.com), ³Scientist (e mail: garuda344@gmail.com), Division of Entomology, IARI, New Delhi.

change by the development of the induced defence system by means of protective enzymes, viz. superoxide dismutase (SOD), catalase (CAT), glutathione S-transferase (GST), peroxidase (POX), and glutathione reductase (GR) (Foyer *et al.* 1994). Morphological, physiological and biochemical changes in the form of accumulation of defensive compounds occur in the plants following herbivory. Reactive oxygen species (ROS) and antioxidative enzymes are recognised in response to herbivory and wounding (Orozco-Cardenas and Ryan 1999). According to Kogan and Paxton (1983), phenolic compounds were accumulated in plants following feeding of insects. Because of the difference in species and genotype of insects, there is a variation in the response of antioxidative systems to elevated CO₂ and temperature (Geetika 2003).

Elevated CO₂ is expected to alter plant physiological processes especially C: N ratio, which would subsequently influence the crop-pest interactions. It thus becomes imperative to investigate the effect of elevated CO₂ on both plants as well as insects at biochemical level. With this background, the present study was planned to estimate the protein content, catalase and peroxidase activity of rice plants subjected to elevated CO₂ and BPH infestation and also to estimate the protein content, catalase and peroxidase activity in BPH that fed on rice plants under ambient and elevated CO₂.

MATERIALS AND METHODS

The study was undertaken at the ICAR-Indian Agricultural Research Institute (IARI), New Delhi (28°38'N latitude, 77°09'E longitude, 228.61 m altitude) during July-October 2016. Seeds of Pusa Basmati 1121, susceptible rice cultivar to *Nilaparvata lugens*, were sown in a nursery (measuring 3×3 sq m). One month old rice seedlings were transplanted in 20 experimental pots (15×9×9 cm³) under natural condition. Two rice seedlings were transplanted per pot and allowed to establish for 10 days. Ten potted plants each of 10-day old were transferred to open top chamber (OTC) with elevated CO₂ (570±25) µl/l and ambient CO₂ (400±25) µl/l level. No insecticide was applied to the plants throughout the study.

Brown planthopper collected from rice fields of IARI were reared and maintained under glasshouse in wire-mesh wooden cages (60×45×45 cm). Out of 10 pots each under elevated and ambient CO₂, five pairs (1:1 sex ratio) of BPH adults were released in five pots apiece, after 10 days of exposure of potted plants to CO₂ in OTCs. Remaining pots were maintained as control without BPH under both the OTCs. The biochemical assay was carried out on BPH infested and uninfested rice plants after one month of exposure of plants to CO₂ in OTCs. Likewise, BPH nymph were collected and analysed for biochemical changes, after one month of release of BPH on rice seedlings at both elevated and ambient CO₂.

BPH infested and uninfested plant samples (tillers) were collected from the experimental potted plants maintained at elevated and ambient CO₂ and were stored at -80°C

for further analysis. The samples were homogenised by hand held motorised homogeniser with addition of 150 µl homogenization buffer (potassium phosphate, 50 mM; pH 7). The homogenate was centrifuged at 12,000 rpm for 30 min at 4°C, and the supernatant was used for energy assay. Protein content was determined using Bradford reagent (Bradford 1976), with bovine serum albumin as the standard. The activity of catalase and peroxidase was analyzed according to the protocol described by Sinha *et al.* (1972) for catalase and Reddy *et al.* (1995) for peroxidase with suitable modifications. Enzyme activity of catalase and peroxidase was computed relative to total protein content. Each test was replicated at least three times. The biochemical assay was carried in infested plants to assess whether the host plant defensive system varied with the injury by the BPH nymphs.

After one month of release of BPH on rice plants in OTCs, five third instar nymphs were collected in 2 ml micro centrifuge tubes in three replicates each from rice plants both under elevated and ambient CO₂ and stored at -80°C for further analysis. The frozen samples were ground with homogenizer after adding homogenization buffer (Potassium phosphate 50 mM, pH 7). The homogenate was centrifuged at 10,000 rpm for 10 min at 4°C, and the supernatant was analyzed. Protein content and the activity of protective enzymes were estimated as described above. Catalase activity (µmoles of H₂O₂ decomposed/min/mg protein) and peroxidase activity (Δ in optical density (OD)/min/mg protein) was calculated relative to total protein content. Each test was replicated at least three times.

The mean protein content, and catalase and peroxidase activity in BPH infested and uninfested rice plant subjected to elevated and ambient CO₂ were analysed through two factorial one way analysis of variance (ANOVA) in Microsoft Excel. Mean differences were compared by least significant difference (LSD) test when significant F-values were obtained (P<0.05). The mean protein content and enzyme activities in *N. lugens* nymphs were compared by paired t-test (P<0.05).

RESULTS AND DISCUSSION

The results of the study on effect of elevated CO₂ and BPH infestation on rice plants revealed that protein content differed significantly across CO₂ levels (P=0.0002, Fig 1A₁) and infestation levels (P=0.001, Fig 1A₂). Rice plants had higher protein content (20.37 g/l) under ambient CO₂ as compared to elevated CO₂ (15.22 g/l) across infestation levels (Fig 1A₁). Similarly, uninfested rice plants had higher protein content (19.71 g/l) as compared to infested rice plant (15.89 g/l) across CO₂ levels (Fig 1A₂). The elevated CO₂ and BPH infestation thus reduced the protein content in rice plants. However, interaction effect between the BPH infestation and CO₂ levels in regard to protein content in plants was not observed.

Catalase (P=0.0001) and peroxidase activity (P=0.00007) in rice plants differed significantly between CO₂ levels across BPH infestation levels, while enzymes activity did not differ




Fig 1 Protein content and protective enzymes activity in BPH infested and uninfested rice plants at elevated and ambient CO₂ condition. (A) Protein content, i.e. g/l (mean±SE), (B) Catalase activity, i.e. μmoles of H₂O₂ decomposed/min/mg of protein (mean±SE) and (C) Peroxidase activity, i.e. ΔOD/min/g of protein (mean±SE). Different lowercase letters indicate significant difference between treatments by least significant difference (LSD) test. A₁: Comparison between CO₂ levels (LSD: 1.935), A₂: Comparison between BPH infestation levels (LSD: 1.935), B₁: Comparison between CO₂ levels (LSD: 0.32), B₂: Comparison between BPH infestation levels (Non-significant), C₁: Comparison between CO₂ levels (LSD: 0.111), C₂: Comparison between BPH infestation levels (Non-significant).

significantly between infestation levels across CO₂ levels. The interaction effect between the BPH infestation and CO₂ levels in regard to catalase and peroxidase activity in plants was not observed.

Rice plants had significantly higher catalase activity (1.39 μmoles of H₂O₂ decomposed/min/mg of protein, Fig 1B₁) under elevated CO₂ compared to ambient CO₂ (0.45 μmoles of H₂O₂ decomposed/min/mg of protein) across infestation levels. The BPH fed rice plants had reduced catalase activity (0.78 μmoles of H₂O₂ decomposed/min/

mg of protein, Fig. 1B₂) compared to uninfested plants (1.07 μmoles of H₂O₂ decomposed/min/mg of protein) across the CO₂ levels thereby impairing the plant defence against elevated CO₂.

Similarly, rice plants had higher peroxidase activity (0.41 ΔOD/min/g of protein, Fig. 1C₁) under elevated CO₂ compared to ambient CO₂ (0.06 ΔOD/min/g of protein), across BPH infestation levels. The peroxidase activity increased non-significantly in infested rice plants (0.264 ΔOD/min/g of protein, Fig 1C₂) compared to uninfested rice plant (0.215 ΔOD/min/g of protein) across CO₂ levels. BPH feeding thus increased peroxidase activity in rice plants. Catalase activity decreased, whereas peroxidase activity increased in response to BPH infestation indicating perhaps peroxidase could be involved in plant defence against BPH besides stress posed due to elevated CO₂.

The protein content in BPH nymphs ($t=3.875$, $df=2$, $P=0.03$; Fig 2A) under elevated CO₂ was significantly lower than ambient CO₂. The BPH that fed on rice plants under elevated CO₂ having lower protein content resulted in lower protein in insects. However, the activity of catalase ($t=9.066$, $df=2$, $P=0.012$; Fig 2B) and peroxidase in BPH nymphs ($t=5.937$, $df=2$, $P=0.027$; Fig 2C) was significantly higher under elevated CO₂ compared to ambient CO₂.

In most of the situation, rice plant responded to the elevated CO₂ by increasing the growth and development, and decreasing the foliar nitrogen and total amino acids, primarily because of build up of non-structural carbohydrates (Wu *et al.* 2007). In the present study, we observed reduced protein content under elevated CO₂ but the activity of catalase and peroxidase was enhanced in rice plant under elevated CO₂ that was consistent with earlier reports of lower protein content in rice at elevated CO₂ (Zeng *et al.* 2012). However, earlier studies reported reduced catalase activity in spruce and tobacco (Havir and McHale 1989), but there was no effect of elevated CO₂ on catalase activity in oak, pine and orange leaves (Schwanz *et al.* 1996). Under elevated condition, the amount of sucrose, glucose, free amino acids, soluble proteins, total non-structural carbohydrates, free amino acids, and nitrogen had increased (Chen *et al.* 2004). Although the effect of elevated CO₂ on plant and insect biochemical constituents often change with time, this has not been taken into account in most studies attempting to characterize the effect of CO₂ enrichment on antioxidant activity. Furthermore, there may be significant intra- and inter-specific changes in response of plant antioxidative enzymes to elevated CO₂ (Badiani *et al.* 1993).

It has been observed earlier that, there was a quantitative variation in biochemical constituents of the most infested plants in terms of increase in levels of proteins, phenols, carbohydrates and enzymes such as peroxidase and catalase activities (Willaims *et al.* 1998). Our study showed that protein content and catalase activity were reduced, while peroxidase activity increased in BPH infested rice plants compared to uninfested rice plants. This has also been reported due to Russian wheat aphid infestation (Mohase and Van der Westhuizen 2002). In




Fig 2 Protein content and protective enzymes activity in BPH nymphs at elevated and ambient CO₂ condition. (A) Protein content, i.e. g/l (mean±SE), (B) catalase activity, i.e. μmoles of H₂O₂ decomposed/min/mg of protein (mean±SE) and (C) peroxidase activity, i.e. ΔOD/min/g of protein (mean±SE). Different lowercase letters indicate significant difference between treatments by the paired T test ($P < 0.05$).

contrast, the increased catalase activity had been reported in yellow stem borer and leaf roller infested rice plants (Usha Rani and Jyothsna 2009) and also in *Helicoverpa zea* infested soybean plants (Bi and Felton 1995), probably because plant response depends on the type of insect feeding. The homopterans feed from the contents of vascular tissue by inserting a stylet between the adjacent cells, thus limiting cell damage and lowering the induced response. The increase in peroxidase activity in BPH damaged plants could be attributed to the fact that it is the key enzyme which participates in several plant cell wall building processes (Chittoor *et al.* 1999).

Insects responded to reduced nutrient content by lowering their behaviour performance and to increased amount of carbohydrates of plants under elevated condition by increasing development time and by reducing growth and survival rates, population densities and fitness (Yin *et al.* 2009). Our results showed that elevated CO₂ reduced the protein content but the activity of protective enzymes, viz. catalase and peroxidase was enhanced in BPH which is was consistent with the earlier reports of enhanced catalase and superoxide dismutase activity in BPH under adverse environmental condition or host plants as protective enzymes provided most important defence to the herbivory (Wu *et al.* 2011). Activities of catalase and cholinesterase of cotton bollworm was significantly affected under elevated condition (Wu *et al.* 2010). A strong correlation occurred between enhanced activities of protective enzymes catalase and peroxidase, and insect adaptation to harsh ecological condition rather than detoxification and digestive enzymes (Zeng *et al.* 2012).

The present study on impact of elevated CO₂ on rice plants and BPH nymphs concludes that elevated CO₂ decreased protein content and increased activity of protective enzymes in both rice plant and insects. This increase in activity of protective enzymes, adapts the insects and plants to accustom in adverse climatic conditions. BPH infestation on rice plants reduced the protein content and catalase activity in rice plants but increased peroxidase activity indicating that it is an important enzyme involved in plant defence against insects.

ACKNOWLEDGEMENTS

The authors are thankful to Head, Division of Entomology, ICAR-IARI, New Delhi for providing necessary facility and UGC for providing fellowship for pursuing Ph D and our thanks extends to Dr Madan Pal Singh, Division of Plant Physiology, Indian Agricultural Research Institute, New Delhi for providing OTC's to carry out the research work.

REFERENCES

- Asghar M, Suhail A, Afzal M and Khan M A. 2009. Determination of economic threshold levels for the stem borers (*Scirpophaga* sp.) and leaf folder (*Cnaphalocrosis medinalis*) of rice (*Oryza sativa*) in the Kallar tract of Punjab, Pakistan. *International Journal of Agriculture and Biology* **11**: 717–20.
- Awmack C S, Harrington R and Lindroth R L. 2004. Aphid individual performance may not predict population responses to elevated CO₂ or O₃. *Global Change Biology* **10**: 1414–23.
- Badiani M D Annibale A, Paolacci A R, Miglietta F and Raschi A. 1993. The antioxidant status of soybean (*Glycine max*) leaves grown under natural CO₂ enrichment in the field. *Australian Journal of Plant Physiology* **20**(3): 275–84.
- Bi J L and Felton G W. 1995. Foliar oxidative stress and insect herbivory: primary compounds, secondary metabolites and reactive oxygen species as components of induced resistance. *Journal of Chemical Ecology* **21**(10):1511–30.
- Bradford M M. 1976. A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye-binding. *Analytical Biochemistry* **72**: 248–54.

- Chen F J, Wu G and Ge F. 2004. Impacts of elevated CO₂ on the population abundance and reproductive activity of aphid *Sitobion avenae* Fabricius feeding on spring wheat. *Journal of Applied Entomology* **128**(9-10): 723–30.
- Chittoor J M, Leach J E and White F F. 1999. Induction of peroxidase during defense against pathogens. (In) *Pathogenesis-related Proteins in Plants*, pp 171–93. Datta S K and Muthukrisnan S (Eds). CRC Press, Boca Raton.
- Foyer C H, Descourvières P and Kunert K J. 1994. Protection against oxygen radicals: an important defence mechanism studied in transgenic plants. *Plant Cell and Environment* **17**(5): 507–23.
- Geetika S. 2003. Polyphenol oxidase production and activity under high temperature exotherm in vegetative organs of *Thuja orientalis*. *Environment Information Archives* **1**: 574–80.
- Havir E A and McHale N A. 1989. Regulation of catalase activity in leaves of *Nicotiana sylvestris* by high CO₂. *Plant Physiology* **89**: 952–7.
- Hughes L and Bazazz F A. 2001. Effects of elevated CO₂ on five plant aphid interactions. *Entomologia Experimentalis et Applicata* **99**(1): 87–96.
- Karnosky D F. 2003. Impacts of elevated atmospheric CO₂ on forest trees and forest ecosystems: knowledge gaps. *Environment International* **29**(2-3): 161–9.
- Khush G S. 2004. Harnessing science and technology for sustainable rice-based production systems. (In) *FAO Rice Conference 04/CRS.14*, 12–13 February 2004, Rome, Italy, p 13.
- Kogan M and Paxton J. 1983. Natural inducers of plant resistance to insects. (In) *Plant Resistance to Insects*, pp 153–71. Hedin P A (Ed). American Chemical Society Symposium, Series 208 American Chemical Society, Washington DC.
- Lu Z X, Heong K L and Yu X P. 2005. Effects of nitrogen on the tolerance of brown planthopper, *Nilaparvata lugens*, to adverse environmental factors. *Insect Science* **12**(2): 121–8.
- Magunmder S K G, Ali M P, Choudhury T R and Rahin S A. 2013. Effect of variety and transplanting date on the incidence of insect pests and their natural enemies. *World Journal of Agriculture Science* **1**(5): 158–67.
- Mohase L and Van der Westhuizen A. 2002. Salicylic acid is involved in resistance responses in the Russian wheat aphid wheat interaction. *Journal of Plant Physiology* **159**(6): 585–90.
- NOAA/ESRL. 2014. www.esrl.noaa.gov/gmd/ccgg/trends/
- Prasannakumar N, Chander S and Pal M. 2012. Assessment of impact of climate change with reference to elevated CO₂ on rice brown planthopper, *Nilaparvata lugens* (Stål.) and crop yield. *Current Science India* **103**(10): 1201–5.
- Reddy K P, Subhani S M, Khan P A and Kumar K B. 1995. Effect of light and benzyladenine and desk treated growing leaves, changes in the peroxidase activity. *Cell Physiology* **26**: 984.
- Reissig W H, Heinrichs E A, Litsinger J A, Moody K, Fiedler L, Mew T W and Barrion A T. 1986. Illustrated guide to integrated pest management in rice in tropical asia. International Rice Research Institute, Manila, Philippines, p 411.
- Schwanz P, Kimball B A, Idso S B, Hendrix D L and Polle A. 1996. Antioxidants in sun and shade leaves of sour orange trees (*Citrus aurantium*) after long-term acclimation to elevated CO₂. *Journal of Experimental Botany* **47**(305): 1941–50.
- Sinha A K. 1972. Colorimetric assay of catalase. *Analytical Biochemistry* **47**(2): 389–94.
- Usha Rani P and Jyothsna Y. 2009. Physiological changes in the groundnut, (*Arachis hypogaea* L.) plant due to the infection of a fungal pathogen, *Cercosporidium personatum* Deighton. *Allelopathy Journal*: 369–78.
- Wu G, Chen F J and Ge F. 2007. Effects of elevated CO₂ on the growth and foliar chemistry of transgenic Bt cotton. *Journal of Integrated Plant Biology* **49**(9): 1361–9.
- Wu G, Chen F J and Xiao N W. 2010. Plant allocation to defensive compounds of transgenic Bt cotton in response to infestation by cotton bollworm under elevated CO₂. *International Journal of Pest Management* **56**(2): 81–9.
- Wu G, Chen F J and Xiao N W. 2011. Influences of elevated CO₂ and pest damage on the allocation of plant defense compounds in Bt-transgenic cotton and enzymatic activity of cotton aphid. *Insect Science* **18**: 401–8.
- Yin J, Yun Y C and Wu G. 2009. No effects of elevated CO₂ on the population relationship between cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae), and its parasitoid, *Microplitis me-diator* Haliday (Hymenoptera: Braconidae). *Agriculture Ecosystem and Environment* **132**: 267–75.
- Zeng Y Y, Huang W, Su L, Wu G, Zhuang J, Zhao W, Hua H, Li J, Xiao N and Xiong Y. 2012. Effects of elevated CO₂ on the nutrient compositions and enzymes activities of *Nilaparvata lugens* nymphs fed on rice plants, *Science China-Life Sciences* **55**(10): 920–6.
- Zhou Y H, Han Z J and Zhang X M. 2003. Influence of rice varieties on the metabolizing enzymes of *Nilaparvata lugens* Stål (in Chinese). *Journal of Nanjing Agriculture University* **26**: 24–8.