Indian Journal of Animal Sciences **93** (6): 640–645, June 2023/Article DOI: https://doi.org/10.56093/ijans.v93i6.129295

Comparative production performance and egg quality of Kuzi Ducks (Anas platyrhynchos) and its crosses with Khaki Campbell

M K PADHI¹⊠, S C GIRI¹ and S K SAHOO¹

Regional Station, ICAR-Directorate of Poultry Research, Bhubaneswar, Odisha 751 003 India

Received: 17 October 2022; Accepted: 6 April 2023

Supplementary Table 1. Age of the flock at different level of production

Genetic group	First egg laid by the flock (day)	Age of the flock at 20% DDEP (day)	Age of the flock at 50% DDEP (day)	Age of the flock at 80% DDEP (day)
Kuzi	102.00±3.46	121.00±3.06a	138.00±3.79a	191.67±1.33ª
DK	105.00 ± 1.53	116.33 ± 0.88^{ab}	122.67±0.67b	$136.67 \pm 2.40^{\circ}$
KD	104.67 ± 1.20	111.00 ± 0.58^{b}	125.00 ± 0.58^{b}	155.33 ± 1.20^{b}

KD, Khaki Campbell \times Kuzi and DK, Kuzi \times Khaki Campbell; DDEP, Duck day egg production. Means having even one superscript in common in a column did not differ significantly ($P \le 0.05$).

Supplementary Table 2. Egg quality parameters at different weeks of age in Kuzi S2 generation and KD , DK (content % and shell thickness and Haugh unit)

Parameter	Genetic	40 wk	44 wk	48 wk	52 wk	60 wk	68 wk	72 wk
	group							
Albumen	KD	58.44±0.43 _A	59.21 ± 0.46	58.25 ± 0.52	57.90 ± 0.38	$55.82\pm0.37_{\rm B}$	$57.22 \pm 0.34_{\rm B}$	57.50 ± 0.56
%	DK	58.35ab±0.34 _A	$58.25^{ab} \pm 0.34$	$58.62^{ab} \pm 0.38$	56.21°±0.61	58.03ab±0.40 _A	59.50°±0.60°	$56.89^{bc} \pm 0.76$
	Kuzi	56.75 ^{bc} ±0.36 _B	$59.16^{a} \pm 0.31$	$57.46^{b} \pm 0.56$	$57.68^{b}\pm0.48$	55.52°±0.45 _B	56.89b±0.43 _B	$55.96^{bc} \pm 0.32$
Yolk %	KD	32.20±0.38	31.88 ± 0.40	32.81 ± 0.48	33.59 ± 0.38	34.16±0.34 _A	34.03±0.29 _A	33.37 ± 0.51
	DK	$32.10^{b} \pm 0.30$	$32.64^{b} \pm 0.35$	$32.28^{b} \pm 0.39$	$35.10^{a} \pm 0.53$	32.26 ^b ±0.45 _B	31.88 ^b ±0.55 _B	$34.14^{a}\pm0.74$
	Kuzi	$33.17^{b} \pm 0.37$	$32.01^{\circ} \pm 0.32$	$33.16^{b} \pm 0.48$	$33.62^{b} \pm 0.36$	34.91°±0.46°	34.02ab±0.48 _A	$34.92^a \pm 0.29$
Shell %	KD	$9.37^{a}\pm0.25$	$8.92^{abc} \pm 0.17$	$8.94^{abc} \pm 0.21$	$8.51^{\circ}\pm0.18$	9.22ab±0.16	8.75 ^{bc} ±0.16	$9.12^{abc} \pm 0.17$
	DK	$9.55^{a}\pm0.17$	$9.11^{ab} \pm 0.13$	$9.10^{ab} \pm 0.15$	$8.70^{b}\pm0.21$	$9.71^{a}\pm0.17$	$8.63^{b} \pm 0.24$	$8.97^{b}\pm0.29$
	Kuzi	$10.08^{a}\pm0.26$	$8.83^{cd} \pm 0.11$	$9.38^{bc} \pm 0.22$	$8.70^{d}\pm0.20$	$9.58^{ab} \pm 0.21$	$9.09^{bcd} \pm 0.17$	$9.12^{bcd} \pm 0.12$
STM	KD	$0.46^{ab} \pm 0.008_{A}$	$0.42^{c}\pm0.004_{B}$	$0.44^{b}\pm0.008$	$0.39^{d}\pm0.007_{B}$	$0.45^{ab} \pm 0.007_{A}$	$0.46^{a}\pm0.009$	$0.35^{\rm e} \pm 0.005_{\rm B}$
(mm)	DK	$0.45^{a}\pm0.004_{A}$	$0.45^{a}\pm0.007_{A}$	$0.43^{ab} \pm 0.008$	$0.40^{\circ} \pm 0.007_{AB}$	$0.43^{\rm b} \pm 0.005_{\rm B}$	$0.44^{ab} \pm 0.010$	$0.33^{d} \pm 0.007_{c}$
	Kuzi	$0.41^{\text{cd}} \pm 0.0069_{\text{B}}$	$0.43^{\rm b} \pm 0.005_{\rm B}$	$0.44^{b}\pm0.007$	0.42bc±0.007	0.43b±0.007 _{AB}	$0.46^{a}\pm0.008$	$0.39^{d} \pm 0.007_{A}$
ST (mm)	KD	0.359ab±0.005 _A	$0.37^{a}\pm0.007_{A}^{b}$	$0.339^{cd} \pm 0.006_{AB}$	$0.328^{d} \pm 0.005$	$0.344^{bcd} \pm 0.005$	$0.354^{abc} \pm 0.008$	$0.301^{\circ}\pm0.004_{_{\rm B}}$
	DK	$0.360^{a}\pm0.005_{A}$	$0.352^{a}\pm0.005_{B}$	$0.325^{b}\pm0.006_{B}$	$0.324^{b}\pm0.005$	$0.342^{ab}\!\!\pm\!0.005$	$0.356^a \pm 0.008$	0.271°±0.006°
	Kuzi	$0.298^{d} \pm 0.005_{B}$	0.356ab±0.005 _{AB}	0.351ab±0.005	$0.334^{\circ}\pm0.006$	$0.340^{bc} \pm 0.006$	$0.359^a \pm 0.008$	0.332°±0.005
Haugh	KD	91.95°±1.21 _A	91.63°±1.49	90.16ab±1.40	$91.98^{a}\pm1.56$	$86.85^{b} \pm 0.62$	$90.26^{ab} \pm 1.15$	91.92a±1.11
unit	DK	93.52°±0.73°	$89.24^{bc}\pm1.57$	$90.80^{abc} \pm 1.95$	$88.80^{\circ} \pm 1.16$	$87.10^{c} \pm 1.16$	$93.12^{ab} \pm 0.99$	$93.19^{ab} \pm 1.16$
	Kuzi	87.13 ^b ±1.11 _B	$90.52^{a}\pm0.97$	$88.92^{ab}{\pm}1.49$	$91.18^{a}\pm1.24$	$89.27^{ab}\!\!\pm\!0.71$	91.13°±1.12	92.25°±0.86

STM, shell thickness with membrane; ST, shell thickness; KD, Khaki Campbell ×Kuzi; DK, Kuzi ×Khaki Campbell. Means having even one common superscript in a row for a particular traits in a genetic groups at different ages did not differ significantly ($P \le 0.05$). Average showing even one subscript in capital letters for a particular trait at same age between different genetic groups did not differ significantly ($P \le 0.05$).

Supplementary Table 3. Mortality % during different period in the three genetic groups

Period	Kuzi (N= 88 at 20 week)	DK (n= 68 at 20 week)	KD (n= 69 at 20 week)
20-40 week	1.14	1.45	4.35
40-60 week	6.82	0.00	1.45
60-72 week	9.09	0.00	8.70
20-72 week	17.05	1.45	14.49
72 to 80 week	1.14	0.00	0.00

KD, Khaki Campbell \times Kuzi and DK, Kuzi \times Khaki Campbell.