



Effect of rice distiller dried grains with solubles (RDDGS) in commercial broiler chicken ration on carcass traits, chemical composition, and fatty acid profile of meat

SRIKANTH KANINDE¹, A ASHOK¹✉, K PREMAVALLI², C BANDESWARAN¹ and R RICHARD CHURCHIL¹

Madras Veterinary College, Chennai, Tamil Nadu 600 007 India

Received: 22 November 2022; Accepted: 18 April 2023

Supplementary Table 1. Effect of feeding different levels of RDDGS in commercial broiler chickens

Sex	Live weight	Giblet weight	Intestinal length	Intestinal weight
Male	2424.34±51.76	3.464±0.23	176.01±3.34	62.75±1.63
Female	2234±47.57	3.607±0.22	161.26±3.07	64.35±1.45
F value	7.21	0.12	10.50*	0.52

Means bearing different superscripts within a row differ significantly (P<0.05). **Significant (P<0.01); *Significant (P<0.05); NS, Not significant; No. of observations (N), 6.

Supplementary Table 2. Effect of feeding different levels of RDDGS in commercial broiler chickens on chemical composition of meat (Mean±SE)

Chemical composition	T1 (control)	T2 (5%)	T3 (10%)	T4 (15%)	F value
Moisture (%)	73.32±0.28	74.23±0.18	72.73±2.46	74.89±1.40	0.55 ^{NS}
Crude protein (%)	21.74±0.61	21.12±0.22	21.15±0.55	20.58±0.36	1.16 ^{NS}
Ether extract (%)	1.07±0.18	1.20±0.38	1.87±0.78	1.20±0.10	0.5 ^{NS}
Total ash (%)	0.99±0.12	1.19±0.95	1.12±0.17	1.49±0.18	2.59 ^{NS}

Means bearing different superscripts within a row differ significantly (P<0.05). **Significant (P<0.01); *Significant (P<0.05); NS, Non significant; No. of observations (N), 6.