

Effect of mycotoxins contaminated corn on growth nutrient digestibility and *in vitro* rumen fermentation in goats

BAI XUE¹, GUO CHUNHUA², ZHANG ZHENGFAN³ and HUANG YANLIN⁴

South West University for Nationalities, Chengdu, Sichuan 610 041 China

Received: 20 May 2013; Accepted: 9 July 2015

Key words: Aflatoxin B₁, Growing goats, Metabolism, Rumen fermentation

Aflatoxins (AFs) are the toxic secondary metabolites produced mainly by the fungi *Aspergillus flavus* and *Aspergillus parasiticus*, which grow on numerous feedstuffs (mainly cereals) when environmental conditions are favorable. The major forms of AFs in feedstuffs include aflatoxin B₁ (AFB₁), aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁) and aflatoxin G₂ (AFG₂). The order of toxicity is AFB₁ > AFG₁ > AFB₂ > AFG₂. AFB₁ is more common than the other AFs (Fandohan *et al.* 2005).

Ruminants seem to be more resistant to aflatoxicosis than monogastrics, but symptoms of aflatoxicosis were reported in bovine (Bodine and Mertens 1983, Cook *et al.* 1986, Diekmann and Green 1992, Auerbach *et al.* 1998, Upadhyaya *et al.* 2010). However, little information about the effects of AFB₁ on nutrient digestion and rumen fermentation is available (Jouany 2001). In addition, these studies were conducted mainly with purified toxins, but rarely with the naturally contaminated feed. In actual production, attention should also be paid to the modification of chemical and physical properties of feedstuffs caused by fungal invasion (Jouany *et al.* 2006, Seeling *et al.* 2006). Therefore, study was conducted to investigate the effects of corn naturally contaminated with aflatoxin B₁ on nutrient digestion and rumen fermentation of goats.

AFB₁-contaminated corns were collected from a feed factory in Sichuan by using a fluorescence detector. The corns naturally mildewed for 40 days at 27° to 30°C and 0.99 water activity. The types and concentrations of mycotoxins in the corns were detected by high performance liquid chromatography (HPLC).

The *in vivo* experiment was conducted at China Sichuan Lezhi Black Goats Sci-Tech Park from August to October 2012. Growing male China Lezhi black goats (12) with an average weight of 16.39 to 16.45 kg were equally divided to a trial group and a control group. Each group was kept in 6 folds (1 goat in each fold). All goats were fed with 40%

concentrated feed (60% corn) in the morning and 60% roughage (corn stover silage + rapeseed shell) in the afternoon and evening. The formula and chemical composition of the concentrate feed are shown in Table 1. The control group was fed with control corns free of mycotoxins; the trial group was fed with AFB₁-contaminated corns for 28 days.

The control corns and mycotoxin contaminated corns were fermented *in vitro* with goat rumen fluid for 3, 6, 12, 24 and 48 h, separately. Each sampling period was conducted in sextuplicate. The rumen fluids were collected from 4 China Lezhi black goats (32±5.3 kg), which were fed with 42.5% concentrate feed, 23.5% silage, 21% grass and 13% alfalfa hay. The rumen fluids were collected by using a rubber stomach tube 0 hr post feeding and squeezed through eight layers of cheesecloth into an Erlenmeyer flask. The rumen fluids were gassed with oxygen-free CO₂ for 15 min and diluted with the McDougall buffer (McDougall 1948) at the ratio of 1:1. The corn samples were fermented *in vitro* with the rumen fluids in calibrated glass syringes following the procedures of Menke *et al.* (1979). The corn samples were milled to pass a 1 mm mesh and weighed 400 mg of DM into the 100 ml calibrated glass syringes. The syringes were pre-warmed at 39°C before injection.

Table 1. Formula and chemical composition of concentrate feed

Item	Diet	
	Control diet	AFB ₁ -contaminated diet
Ingredient	% as fed	
Corn	60.00	0
AFB ₁ -contaminated corn	0	60.00
Commercial weaning feed	3.00	3.00
Silkworm faeces	23.53	23.53
Fermentation feed	7.90	7.90
Distiller's grains	5.57	5.57
Nutrient composition	%DM	
Crude protein (CP)	14.89	15.25
Crude fiber (CF)	10.00	10.16
Acid detergent fiber (ADF)	22.14	22.77
Neutral detergent fiber (NDF)	32.72	33.49

*The work was supported by Animal Science Discipline program of Southwest University for Nationalities (2011XWD-S0905)

Present address: ¹(yukivy@163.com), College of Life Science and Technology.

Rumen fluid-buffer mixture (30 ml) was injected into each syringe. The corn samples were fermented in a thermostatic incubator at 39 ± 0.5 °C for 3, 6, 12, 24 and 48 h, separately.

The 100 g corn and diet were collected twice each week and stored at -20 °C. All of the corn samples were mixed at the end of experiment to quantify the mycotoxin contents by high performance liquid chromatography.

Growing goats were weighed at the beginning of the experiment and on day 28, separately. Feed (concentrated feed, silage and rapeseed shell) intake was recorded each day. Thus average daily gain (ADG), feed intake (FI) and feed conversion ratio (FCR) were determined. Abnormal behavior and daily mortality were recorded, and dead goats were dissected. Three goats in each group were killed on day 28 and the ruminal fluids were collected for measurements of pH value, ammonia nitrogen ($\text{NH}_3\text{-N}$) and volatile fatty acid (VFA).

The concentrated feed, silage, rapeseed shell and fecal samples were collected at day 26, 27, 28. The samples of the 3 days were mixed and stored at -20 °C for determination of dry matter (DM), crude protein (CP), crude fiber (CF), neutral detergent fiber (NDF), acid detergent fiber (ADF) and 4N Acid-insoluble ash (AIA). Apparent digestion coefficients were calculated by using the formula given by Church and Pond (1982) with AIA as the internal indicator. In the *in vitro* experiment, the volume of gas was measured at 3, 6, 12, 24 and 48 h after initiation of incubation. Total gas values were corrected by blank incubation. The pH, $\text{NH}_3\text{-N}$ and VFA of fermentation fluid were measured at 3, 6, 12, 24 and 48 h.

The mycotoxins in corns and diets were quantified via HPLC. Separation was achieved on an SB-C18 column (4.6 × 250 mm; 5 μm). The mobile phases were methanol and water (45:55, v/v) for aflatoxins (AFB_1 , AFB_2 and AFG_2); acetonitrile and water (15:85, v/v) for deoxynivalenol (DON); acetonitrile and water (80:20, v/v) for T-2 toxin; acetonitrile, water and methanol (46:46:8, v/v/v) for zearalenone (ZEN). The flow rate was 1 ml/min, and the injection volume was 50 μl .

For the feed and fecal samples, DM and CP were determined with AOAC (1990) procedures; CF, NDF, and ADF were determined with the procedure of Goering and Van Soest (1970); 4N AIA was determined with the procedure of Van Keulen and Young (1977).

For both ruminal fluid (*in vivo*) and fermentation fluid (*in vitro*), the pH values were measured immediately with a pH meter, the $\text{NH}_3\text{-N}$ levels were detected with indophenol colorimetry (Zongci and Ming 2010), and VFA (acetic acid, propionic acid, butyric acid) concentrations were determined by using HPLC with a C_{18} column.

All data were analyzed by SPSS 18.0. *In vivo* statistical analyses were performed by using T-test Duncan, and *in vitro* statistical analyses were performed by using GLM with repeated statement. The fixed effects in the model were AFB_1 level, incubation time, and the interaction between AFB_1 level and incubation time. Individual incubation bottle nested within treatment was a random effect.

The gas production, gas production rate, pH and $\text{NH}_3\text{-N}$ at each incubation time point were analyzed by using T-test. All means were considered significantly different at $P<0.05$.

The concentrations of mycotoxins in control corn, mycotoxin naturally contaminated corn, control diet, and mycotoxin naturally contaminated diet are given in Table 2. The mycotoxins in the control diet were below the detection limit. The contaminated diet had 74.49 $\mu\text{g/kg}$ AFB_1 , 2.08 $\mu\text{g/kg}$ AFB_2 , 59.71 $\mu\text{g/kg}$ DON and 36.51 $\mu\text{g/kg}$ ZEN. Except AFB_1 , the other mycotoxins were below the legal limits established by the European Union (European Commission 2010).

During the experiment, no typical clinical illness attributable to aflatoxicosis was observed. One goat in the control group died due to the parasitism of brain spore worm. The AFB_1 -caused alterations in the growing goats' performance are shown in Table 3. The AFB_1 -contaminated corn showed no significant effects on the intake of concentrated feed, silage, or rapeseed shell of goats ($P>0.05$) during the whole experimental period. The average daily gain (ADG) of the trial group was 81.4g, which was lower than the control group (20.2g/d, $P=0.086$). The FCR of the control group and the trial group were calculated to

Table 2. The concentrations of mycotoxins in corns and diets ($\mu\text{g/kg}$)

	Control corn	Mycotoxin-contaminated corn	Control diet	Mycotoxin-contaminated diet
AFB_1	N. D	289.54	N. D	74.49
AFB_2	N. D	19.54	N. D	2.08
AFG_1	N. D	10.34	N. D	ND
DON	76.49	165.01	N. D	59.71
ZEN	42.9	102.14	N. D	36.51
T-2	N. D	23.47	N. D	ND

ND, No detect.

Table 3. The effects of AFB_1 on the performance of growing goats

	Control group	AFB_1 -corn group	P value
Daily feed intake (g/day)	923.86 $\pm$ 48.42	905.78 $\pm$ 60.49	0.142
Intake of concentrated feed (g/day)	372.18 $\pm$ 32.62	375.35 $\pm$ 13.24	0.831
Intake of silage (g/day)	460.75 $\pm$ 14.56	454.45 $\pm$ 39.92	0.747
Intake of rapeseed shell (g/day)	81.28 $\pm$ 2.59	75.97 $\pm$ 8.37	0.208
Initial body weight (kg)	16.39 $\pm$ 0.74	16.45 $\pm$ 0.5	0.638
Final body weight (kg)	19.09 $\pm$ 1.09	18.83 $\pm$ 0.20	0.83
Average daily gain (ADG), (g/day)	101.69 $\pm$ 17.26	81.4 $\pm$ 13.42	0.086
Feed conversion ratio (FCR), (g/g)	9.08 $\pm$ 1.56	10.54 $\pm$ 1.72	0.106

be 9.08 ± 1.56 and 10.54 ± 1.72 , respectively. AFB₁ improved the feed conversion ratio and decreased the feed conversion efficiency in growing goats ($P > 0.05$).

The effects of AFB₁ on nutrient digestibility of growing goats are shown in Table 4. The CP digestibility of the growing goats fed with AFB₁-contaminated corn was significantly reduced by 7.05 % ($P < 0.05$) compared to the control group. The digestibility of DM, CF, NDF or ADF had no significant change ($P > 0.05$). Ruminants, however, are considered generally more resistant to the adverse effects of mycotoxins, because the rumen microbiota could bio-transform the mycotoxins. But some of the metabolites of mycotoxins are still toxic (Fink-Gremmels 2008). In this study, the aflatoxin had no significant effect on feed intake of growing goats, but the body weight gain and feed consumption were lower ($P > 0.05$). Similar results were

found in ruminants like lamb (Fernandez 2000), deer (Quist *et al.* 1997), steer (Richard *et al.* 1983) and dairy cow (Korosteleva *et al.* 2009).

Mycotoxin contamination may change the digestibility of individual feed components (Gremmels *et al.* 2008). Many experiments about mycotoxin were for the absorbents, but some experimental design had no normal feed treatment. For example, Kiyothong *et al.* (2012) reported that the pH, NH₃-N, VFA, amounts of rumen bacteria, and the digestibility of CP, NDF were significantly ($P < 0.05$) higher in mycotoxin deactivator supplemented than in cows fed with contaminated mycotoxin, and some of those indexes were not within the normal physiological range. Therefore, if we could suppose that the mycotoxin had some negative influence on the metabolism of cows. In this study, the AFB₁ significantly reduced the CP

Table 4. The effects of AFB₁ on the nutrient digestion of growing goats

	DM (%)	CP (%)	CF (%)	ADF (%)	NDF (%)
Control group	61.52±2.30	60.74±2.41*	52.83±3.77	62.46±4.32	61.54±3.63
AFB ₁ -corn group	59.88±2.61	56.46±2.72	54.65±4.58	59.44±7.15	57.02±10.97
P value	0.419	0.035	0.531	0.515	0.456

*, $P < 0.05$.

Table 5. The effects of AFB₁ on the rumen fermentation of growing goats

	pH	NH ₃ -N (mg/dl)	acetic acid (mmol/l)	propionic acid (mmol/l)	butanoic acid (mmol/l)
Control group	7.32±0.04	8.34±0.74*	44.52±1.70	10.93±0.42	5.76±0.69
AFB ₁ -corn group	7.22±0.04	6.21±0.03	43.04±3.32	10.05±0.48	5.51±0.99
p value	0.441	0.046	0.631	0.194	0.796

*, $P < 0.05$.

Table 6. *In vitro* rumen fermentation parameters

	3h	6h	12h	24h	48h	SEM	P value ^a		
							Time	AFB ₁	Time×AFB ₁
Gas (ml)									
Control group	11.45**	47.14**	83.33**	90.67	97.67	3.738	<0.001	<0.001	0.015
AFB ₁ -corn group	1.82	16.67	64.67	75.00	90.67				
T-test P value	<0.001	<0.001	0.005	0.06	0.588				
Gas rate (ml/h)									
Control group	3.82**	7.86**	6.94**	3.78	2.11	0.342	<0.001	<0.001	<0.001
AFB ₁ -corn group	0.61	2.78	5.39	3.13	1.89				
T-test P value	<0.001	<0.001	0.005	0.06	0.588				
pH									
Control group	7.28	6.88	6.47	6.23	5.88*	0.178	<0.001	0.105	0.332
AFB ₁ -corn group	7.15	6.87	6.44	6.16	5.60				
T-test P value	0.178	0.67	0.833	0.136	0.045				
NH ₃ -N (mg/dl)									
Control group	17.52	10.88**	5.43**	4.30**	3.26	0.931	<0.001	<0.001	0.003
AFB ₁ -corn group	15.56	15.06	13.37	8.73	5.83				
T-test P value	0.486	0.001	0.001	0.008	0.252				

*, $P < 0.05$; **, $P < 0.01$.^a Time, fermentation time; AFB₁, AFB₁ contamination treatment; time×AFB₁, time×AFB₁ contamination treatment.

digestibility ($P < 0.05$). Rao (2003) reported that AF administration (300 ppb) resulted in lower CP digestibility ($P < 0.01$), and it was not improved even by supplementation with Sodium Bentonite or Activated Charcoal. The same result was reported in duck (Han *et al.* 2008) and laying hens (Kermanshahi *et al.* 2007). AFB₁ could inhibit the rumen microbial activity, such as *Streptococcus bovis* and *Bacillus megaterium*, and affect rumen microbial flora diversity, such as four different strains of *B. fibrisolvens* (Westlake *et al.* 1987, Kiessling *et al.* 1984, Mertens *et al.* 1979). Those studies showed that the AFB₁ inhibited rumen microbial fermentation. Cook *et al.* (1986) reported that aflatoxin could change the rumen motility in steers, like the amplitude (strength) and frequency of rumen contractions decreased. Therefore, AFB₁ may affect nutrient digestibility in ruminants by those ways.

The activities of carboxymethyl cellulase and microcrystalline cellulase in dairy cow rumen fluid were inhibited by AFB₁ *in vitro* (Jiang *et al.* 2010). Dvorak (1977) reported that aflatoxin decreased cellulose digestion in dairy cows. In this study, the NDF and ADF digestibility decreased too, but not significantly ($P > 0.05$).

The rumen fermentation of growing goats is showed in Table 5. The NH₃-N content of the trial group (6.21 ± 0.03 mg/dl) was significantly lower than the control group (8.34 ± 0.74 mg/dl) ($P < 0.05$). The pH and fractions of VFA were similar in control and treatment group in the trial group were lower than the control group, but not significant ($P > 0.05$).

The *in vitro* effects of AFB₁ on gas production, gas production rate, rumen fluid pH value and NH₃-N are shown in Table 6. The gas production and gas production rate were significantly influenced by the incubation time, AFB₁ and the interaction between incubation time and AFB₁ ($P < 0.05$). The gas production increased with the prolongation of incubation time ($P < 0.01$). During fermentation period from 0 to 3 h, the cumulative gas production of the AFB₁-corn group (1.82 ml) was significantly lower than that of the control group (11.45 ml) ($P < 0.01$). However, the total gas production showed no significant difference between the control group (97.67 ml) and AFB₁-corn group (90.67 ml) ($P = 0.588$). The gas production rate of the control group was always higher than the AFB₁-corn group in the incubation period from 0 to 48 h. The highest gas production rate appeared at 6 h (7.86 ml/h) in the control group and later at 12 h (5.39 ml/h) in the AFB₁-corn group.

AFB₁ had no significant effect on pH ($P = 0.105$). The pH value became increasingly lower with extension of incubation time ($P < 0.001$). At 48 h, the pH values of the control group and AFB₁-corn group were 5.88 and 5.6, respectively. Because the incubation metabolic products were accumulated, the pH at 48 h was not within the physiological rumen pH range (6 to 7).

The NH₃-N levels in the two groups continued to decrease in total fermentation ($P < 0.001$), and the decline in the AFB₁-corn group was slower than in the control group. The two groups showed no significant differences

at 3 h or 48 h. But at 6 h, 12 h and 24 h, the NH₃-N levels in the AFB₁-corn group were significantly higher than in the control group ($P < 0.001$). And there was a significant interaction between incubation time and AFB₁ content ($P = 0.003$).

NH₃-N is the decomposition product of rumen nitrogenous substances. Under adequate supply of soluble carbohydrate and energy, NH₃-N could be synthesized for bacterial protein by rumen microorganisms. In this *in vitro* experiment, the NH₃-N content of AFB₁-corn group was lower than the control group ($P > 0.05$) at 0 h to 3 h. The result was similar to the *in vivo* experiment and consistent with the lower degradation rate of crude protein of the AFB₁-corn group in the digestion test too. This was because the rumen microorganism was inhibited by AFB₁, so the nitrogenous substance in corn could not be effectively degraded to NH₃-N. But the *in vitro* trial was a closed system, the large amount of NH₃-N produced by the rumen microorganism fermentation at early stage cannot be absorbed by the rumen wall. And the ability of microbial protein synthesis was weakened by AFB₁. So the NH₃-N content of the AFB₁-corn group was higher than the control group ($P < 0.05$) in the late stage of *in vitro* fermentation.

The pH value and VFA had no significant changes ($P > 0.05$). Edrington *et al.* (1994) reported that after goats were fed with 2.5 mg of AF kg diet for 35 days, the rumen pH and VFA had no significant changes. Helferich *et al.* (1986) got similar results on steer. But Bodine and Mertens (1983), Jiang (2010) presumed that rumen microorganisms were very sensitive to aflatoxin, the VFA production and proportion were changed by aflatoxin *in vitro* test. The different results may be related to the dose, animal species and differences between naturally contaminated mycotoxin and pure mycotoxin.

Gas production and gas production rate are intuitive indicators of rumen fermentation. In this *in vitro* experiment, the gas production and gas production rate in the AFB₁-corn group were both significantly lower than in the control group ($P < 0.01$). It showed that the AFB₁ inhibited rumen microbial fermentation. Westlake *et al.* (1989) reported that with the addition of 1~10 µg/ml aflatoxin M, the rumen microbial digestion decreased by 50 to 67%. The reason may be that the rumen microbes in fermentation fluid changed. In addition, we found that the incubation time and AFB₁ content had significant effects on the rumen fermentation of goats. As early as 0 h to 3 h, the gas production and gas production rate of the AFB₁-corn group extremely significantly decreased, which indicated that AFB₁ may affect the rumen fermentation in the initial administration.

Corns naturally contaminated with harmful mycotoxins had no significant effect on the feed intake of growing goats, but decreased the nutrient digestibility and adversely affected the *in vitro* rumen fermentation. These lead to the reduction of ADG and feed consumption. Although ruminants have certain tolerance to mycotoxin, the rumen is negatively affected by mycotoxins when the rumen

degrades the mycotoxin. Accordingly, mycotoxin contamination on ruminant feed cannot be ignored.

SUMMARY

Two trials (*in vivo* and *in vitro*) were conducted to evaluate corn naturally contaminated with mycotoxins, majority being aflatoxin B₁ (AFB₁) on the performance, nutrient digestion and rumen fermentation in growing goats. China Lezhi black goats (12), weighing 16.39 to 16.45 kg, were fed with the diet of 40% concentrate (the mycotoxin naturally contaminated diet containing 74.49 µg/kg AFB₁, 2.08 µg/kg AFB₂, 59.71 µg/kg DON and 36.51 µg/kg ZEN) for 28 days. The results showed that the contaminated corn had no significant effect on feed intake but decreased the average daily gain (ADG) and feed conversion ratio (FCR) in growing goats. Digestibility of crude protein (CP) in the trial group was significantly lower than the control group and the digestibilities of acid detergent fibre (ADF) and neutral detergent fibre (NDF) decreased too, but not significantly. Neither volatile fatty acid (VFA) nor pH was significantly different between the 2 groups. The ammonia nitrogen (NH₃-N) in the trial group was lower in both *in vivo* trial and *in vitro* trial (0 h to 3 h). In *in vitro* experiment, ruminal fluids were collected from 4 China Lezhi goats and incubated at 39°C for 48 h with control corn or AFB₁ contaminated corn. The total gas production and gas production rate in the trial group were significantly lower than the control group. These reductions showed the negative effects of the naturally contaminated AFB₁ corn on nutrient digestibility and rumen function in growing goats.

REFERENCES

- AOAC. 1990. Official Methods of Analysis. *Association of Analytical Chemists*. 15th Ed. Arlington, Virginia, USA.
- Auerbach H, Maas R F M, Op Den Camp H J M, Pol A and Fink Gremmels J. 1998 b. Biodegradation of aflatoxin B₁ by bovine rumen microorganisms *in vitro* and its effects on rumen fermentation. *Revue de Médecine Vétérinaire* **149**: 573.
- Bodine A B and Mertens D R. 1983. Toxicology, metabolism and physiological effects of aflatoxin in the bovine. *Aflatoxin and Aspergillus flavus in Corn*. (Eds) Diener U L, Asquith R L and Dickens J W. Alabama Agnc. Exp. Sta. Coop. Ser. Bull 279:46
- Church D C and Pond W G. 1982. *Basic Animal Nutrition and Feeding*. 3rd edn. John Wiley & Sons, New York.
- Cook W O, Richard J L, Osweiler G D and Trampel D W. 1986. Clinical and pathologic changes in acute bovine aflatoxicosis: rumen motility and tissue and fluid concentrations of aflatoxins B₁ and M₁. *American Journal of Veterinary Research* **47**: 1817–25.
- Diekman M A and Green M L. 1992. Mycotoxins and reproduction in domestic livestock. *Journal of Animal Science* **70**: 1615–27.
- Dvorak R, Jagos J, Bouda J, Piskac A and Zapetal O. 1977. Changes in the clinico-biochemical indices in the rumen liquor and urine in cases of experimental aflatoxicoses in dairy cows. *Veterinary Medicine. (Praha)* **22** (3): 161.
- Erdington T S, Harvey R B and Kubena L. 1994. Effects of aflatoxin in growing lambs fed ruminally degradable or escape protein sources. *Journal of Animal Science* **72**: 1274–81.
- European Commission. 2010. Commission Regulation (EU) No. 165/2010 of 26 February 2010 amending Regulation (EC) No. 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards aflatoxins. *Official Journal of the European Union* **L50**: 8–12.
- Fandohan P, Zoumenou D, Hounhouigan D J, Marasa W F, Wingfield M J and Hell K. 2005. Fate of aflatoxin and fumonisins during the processing of maize into food products in Benin. *International Journal of Food Microbiology* **98**: 249–59.
- Feng Zongci and Gao Ming. 2010. Improved method for measurement of rumen fluid ammonia content by colorimetric. *Animal Husbandry and Feed Science* **31**: 37.
- Fernández A, Hernández M, Verde M T and Sanz M. 2000. Effect of aflatoxin on performance, hematology, and clinical immunology in lambs. *Canadian Journal of Veterinary Research* **64**: 53–58.
- Fink-Gremmels J. 2008. Mycotoxins in cattle feeds and carryover to dairy milk: A review. *Food Additives and Contaminants* **25** (2): 172–80.
- Goering H J and Van Soest P J. 1970. Forage fiber analysis (Apparatus, reagents, procedure and some applications). *USDA Agricultural Handbook No. 379*. US Print Office Washington, DC, USA.
- Han Xinyan, Huang Qichun, Li Weifen, Jiang Junfang and Xu Zirong. 2008. Changes in growth performance, digestive enzyme activities and nutrient digestibility of cherry valley ducks in response to aflatoxin B₁ levels. *Livestock Science* **119**: 216–20.
- Helferich W G, Garrett W N, Hsieh D P H and Baldwin R L. 1986. Feedlot performance and tissue residues of cattle consuming diets containing aflatoxins. *Journal of Animal Science* **62**: 691–96.
- Jiang Yahui and Yang Hongjian. 2010. Inhibitory effect of different levels of aflatoxin B₁ addition on fermentation characteristic of mixed rumen microorganisms *in vitro*. *China Agricultural Science. Bulletin* **26** (16): 8–13.
- Jouany J P. 2001. The impact of mycotoxins on performance and health of dairy cattle. Science and Technology in the Feed Industry. *Proceedings of Alltech's 17th Annual Symposium*. pp. 191–224. (Eds) Lyons T P and Jacques K A. Nottingham University Press, UK.
- Kermanshahi H, Akbari M R, Maleki M and Behgar M. 2007. Effect of prolonged low level inclusion of aflatoxin B into diet on performance, nutrient digestibility, histopathology and blood enzymes of broiler chickens. *Journal of Animal and Veterinary Advances* **6** (5): 686–92.
- Kiessling K H, Pettersson H, Sandholm K and Olsen M. 1984. Metabolism of aflatoxin, ochratoxin, zearalenone and three tricothecenes by intact rumen fluid, rumen protozoa and rumen bacteria. *Applied and Environmental Microbiology* **47**: 1070–73
- Kiyothong K, Rowlinson P, Wanapat M and Khampa S. 2012. Effect of mycotoxin deactivator product supplementation on dairy cows. *Animal Production Science* **52** (9): 832–41.
- Korosteleva S N, Smith T K and Boermans H J. 2009. Effects of feed naturally contaminated with fusarium mycotoxins on metabolism and immunity of dairy cows. *Journal of Dairy Science* **92**: 1585–93.
- McDougall E T. 1948. Studies on ruminant saliva 1. The composition and output of sheep's saliva. *Biochemical Journal* **43**: 99–108.
- Menke K L, Raab A, Salewski H, Steingass D F and Schneider

- W. 1979. The estimation of the digestibility and metabolizable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen liquor in vitro. *Journal of Agricultural Science (Camb)* **93**: 217–22.
- Mertens D R. 1979. Biological effects of mycotoxins upon rumen fermentation and lactating dairy cows. *Interactions of Mycotoxins in Animal Production*. pp 118–36. National Academy Press, Washington, DC.
- Quist C F, Howerth E W, Fisher J R, Wyatt R D, Miller D M and Nettles V F. 1997. Evaluation of low-level aflatoxin in the diet of white-tailed deer. *Journal of Wildlife Diseases* **33**: 112–21.
- Rao S B. and Chopra R C. 2003. Effect of sodium bentonite and activated charcoal on nutrient utilization in goats receiving diets added with aflatoxin B1. *Indian Journal of Animal Nutrition* **20**: 23–29.
- Richard J L, Pier A C, Stubblefield R D, Shotwell O L, Lyon R L and Cutlip R C. 1983. Effect of feeding corn naturally contaminated with aflatoxin on feed efficiency, on physiologic, immunologic, and pathologic changes, and on tissue residues in steers. *American Journal of Veterinary Research* **44** (7): 1294–98.
- Seeling K, Danicke S, Valenta H, Egmond H P van, Schothorst R C, Jekel A A, Lebzein P, Schollenberger M, Razzazi-fazeli E and Flachowsky. G 2006. Effects of Fusarium toxin contaminated wheat and feed intake level on biotransformation and carry-over of deoxynivalenol in dairy cows. *Food Additives and Contaminants* **23** (10): 1008–20.
- Upadhaya S D, Park M A and Jong K Ha. 2010. Mycotoxins and their biotransformation in the rumen: a review. *Asian-Australasian Journal of Animal Sciences* **23** (9): 1250–60.
- Van Keulen J and Young B A. 1977. Evaluation of acid-insoluble ash as a natural marker in ruminant digestibility studies. *Journal of Animal Science* **44**: 282.
- Westlake K R, Mackie I and Dutton M F. 1989. In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations. *Animal Feed Science and Technology* **25**:169–78.
- Westlake K R, Mackie I and Dutton M F. 1987. Effects of several mycotoxins on specific growth rate of *butyrivibrio fibrisolvens* and toxin degradation in vitro. *Applied and Environmental Microbiology* **53** (3): 613–14.

BOOK REVIEW

Parasitic Zoonoses in the New Millennium. 2015. M.B. Chhabra and K.M.L. Pathak. Today & Tomorrow's Printers and Publishers 4436/7, Ansari Road, Daryaganj, New Delhi 110012. 15.5 cm×23.5 cm. pp i – xii + 1-269. ISBN: 81-7019-519-1 (India), 1-55528-575-6 (USA).

Parasitic zoonoses or diseases caused by parasites transmissible between men and animals, is one of the major problems faced by humanity worldwide. In recent times, it has been exacerbated by several environmental and human factors, thereby enhancing the risk perceptions. Global agencies like WHO, and FAO, through a series of Expert Committee Reports, have brought increased attention and heightened concern on the impact of these diseases on the health and economy of vast populations. Parasitic Zoonoses is a highly relevant and rapidly-evolving sub-discipline of parasitology involving both medical and veterinary interesting.

The contents are arranged in six sections: first is Introduction which includes different classification infectious diseases, epidemiology and socio-economic impact of zoonoses. The other five sections deal with 52 diseases under major groups of etiological agents, namely Protozoa (14), Trematoda (10), Cestoda (8), Nematoda (14) and Arthropod zoonoses (6). Each disease has been described under appropriate sub-headings, namely synonyms, etiology, history (where available), life-cycle, epidemiology, clinical syndrome in man and animals, diagnosis and control who may wish to do additional reading. The Illustrations and Data

are given to explain the facts in 10 Tables. In view of the emerging significance of wildlife as additional reservoirs of zoonoses and Appendix listing the various zoonotic parasites recorded from different wildlife species in India, has been added.

Coming at a time palpable paucity of comprehensive texts on this subject, particularly in India, the present authors deserve kudos for their effort to fill the void. This book scores over its contemporaries in certain special features. In keeping with its title, most of the literature cited pertains to the post-2000 period, material has been drawn from both medical and veterinary sources and despite its worldwide outreach, the Indian perspective has given special emphasis. It is heartening to note that the often-neglected area of arthropod zoonoses has been given adequate attention. Emerging (and re-emerging) zoonoses like cryptosporidiosis, toxoplasmosis, cysticercosis, hydatidosis and myiasis like have been allotted greater space in proportion to their growing importance. The certain disease entities like fascioliasis, larva migrans and zoonotic giardiasis, have made progressive inroads in parts of India. The language is simple, lucid and concise. The overall get-up and production of the book is excellent. High quality printing and aesthetically-pleasing cover design, add to the value of the book. The book will be eminently suitable reference manual for post-graduate, as well as advanced undergraduate students of both veterinary and human medicine, parasitological and public health in particular, and those of biomedical sciences, in general.

Dr Sudhir Pradhan
ICAR