



## Effect of solid state fermentation on nutrient content and ileal amino acids digestibility of palm kernel cake in broiler chickens

MOHAMED IDRIS ALSHELMANI<sup>1</sup>, TECK CHWEN LOH<sup>2</sup>, HOOI LING FOO<sup>3</sup>,  
AWIS QURNI SAZILI<sup>4</sup> and WEI HONG LAU<sup>5</sup>

*Universiti Putra Malaysia, Serdang, Selangor 43400 Malaysia*

Received: 27 February 2017; Accepted: 27 March 2017

### ABSTRACT

Digestibility trial was conducted to determine the apparent ileal digestibility (AID) of crude protein (CP) and amino acids (AA) in untreated palm kernel cake (PKC) and fermented palm kernel cake (FPKC) on finisher broiler. *Paenibacillus polymyxa* ATCC 842 and *P. curdlanolyticus* DSMZ 10248 were used to produce FPKCa and FPKCb, respectively through solid state fermentation (SSF). Broiler male chickens were fed with diets containing 15% PKC from day one until 41 days of age. Birds (36) were selected with uniform body weight, and randomly distributed into 3 groups with 6 replicates in each treatment and 2 birds per replicate. The chickens were deprived from food overnight with free access to drinking water. The birds were fed PKC, FPKCa and FPKCb with indigestible marker. All the chickens were allowed free access to the test ingredients and drinking water for 4 days. The birds were slaughtered; ileal digesta were individually collected, pooled within each replicate in plastic cups; and immediately kept at  $-20^{\circ}\text{C}$  for chemical analysis. The findings showed that the process of SSF by cellulolytic bacteria increased the levels of CP from 16.43% in the PKC to 16.68% and 16.80% in FPKCb and FPKCa, respectively. The AID of CP was increased in FPKC compared to the PKC. Additionally, there was an increase in the digestibilities of AA in FPKC compared to untreated PKC. The process of SSF decreased the fibres in FPKC, and there was improvement in the nutrient value of FPKC by cellulolytic bacterial cultures in terms of nutrient content and digestibility.

**Key words:** Amino acids, Broiler, Cellulolytic bacteria, Digestibility, Fermented palm kernel cake

The most common feedstuffs used in poultry diets are soybean meal and yellow corn as important sources for protein and energy, respectively. The soybean meal provides protein and most essential amino acids (AA), whereas the yellow corn is considered as a main source of energy for the chickens. It is also known that availability of AA in these feedstuffs is high, and therefore enable the chickens to exhibit high growth performance. There are attempts to substitute yellow corn and soybean meal with local feedstuffs in order to decrease the cost of feed, and to achieve the food security in the developing countries. However, the local feedstuffs are poor in nutrient digestibility, crude protein (CP) and energy content. Moreover, the local ingredients are high in crude fibres and non-starch polysaccharides (NSPs) as well as anti-nutritional factors.

Malaysia produces abundant amount of palm kernel cake (PKC) as an agro-industrial by-product from palm fruits during oil extraction. The nutritive value of PKC is 16%

CP and contains high content of crude fibres as well as NSPs, especially mannans (Sundu and Dingle 2002, Alimon 2004, Fadil *et al.* 2014). The challenge point of using the PKC in broiler diets is the content of NSPs and coarse texture (McDonald *et al.* 1995, O'Mara *et al.* 1999, Sundu and Dingle 2002). Therefore, there were attempts to improve the nutritive quality of the PKC by using exogenous enzymes (Ng and Chong 2002, Iyayi and Davies 2005, Lawal *et al.* 2010, Alvarez-Cervantes *et al.* 2013, Chen *et al.* 2013, Kocher *et al.* 2002, Adebisi and Olukosi 2015), or by applying solid state fermentation (SSF) using cellulolytic microorganisms (Marini *et al.* 2005, Iluyemi *et al.* 2006, Graminha *et al.* 2008, Lateef *et al.* 2008, Muangkeow and Chinajariyawong 2009, Sukaryana *et al.* 2010, Mirnawati *et al.* 2011, Alshelmani *et al.* 2013, Gao *et al.* 2013, Alshelmani *et al.* 2014, Alshelmani *et al.* 2016a, Alshelmani *et al.* 2016b). Although many researches were conducted by using fungi in SSF, the secondary metabolites resulting by fungi can suppress the growth of animals (Khin 2004, Marini *et al.* 2005, Moftah *et al.* 2012). Therefore, fermented PKC cellulolytic microorganisms in SSF may alleviate this problem.

The presence of NSPs in poultry diets may lead to increased viscosity, raise the speed of passage time of the ingesta in gastro-intestinal tract (GIT), and therefore

Present address: <sup>1</sup>(alshelmani@gmail.com), Department of Animal Production, University of Benghazi, Benghazi, Libya.  
<sup>2</sup>(tcloh@upm.edu.my), Department of Animal Science;  
<sup>3</sup>(hlfoo@upm.edu.my), Department of Bioprocess Technology;  
<sup>4</sup>(awis@upm.edu.my), <sup>5</sup>(lauweih@upm.edu.my), Department of Plant Protection.

decrease the absorption and utilization of nutrients in the small intestines. Consequently, it appears to be important to examine this by-product in poultry nutrition, and study the AA digestibility to the PKC after fermentation process. Since *P. polymyxa* and *P. curdlanolyticus* exhibited high capabilities in degrading crude fibre and NSPs in our previous studies (Alshelmani *et al.* 2013, 2014), the objective of this experiment was to investigate the digestibility of CP and AA in the fermented PKC (FPKC) by cellulolytic bacteria.

#### MATERIALS AND METHODS

**Preparation of fermented palm kernel cake:** PKC was fermented by *Paenibacillus polymyxa* ATCC 842 and *P. curdlanolyticus* DSMZ 10248. The fermentation process was conducted as described by Alshelmani *et al.* (2013, 2014). The bacterial cultures were revived (sub-cultured), and the optical density (OD) was adjusted to 1.00 at 600 nm using spectrophotometer. Inoculum [10% (v/w)] was then transferred into each conical flask. The conical flasks inoculated with cellulolytic bacteria were incubated for 9 days under humidified conditions by placing sterile distilled water inside the incubator. The FPKC was autoclaved and dried after SSF.

**Birds and experimental design:** The study was conducted according to the guidelines of the Research Policy on Animal Ethics of the Universiti Putra Malaysia. Day-old male broiler chicks (Cobb 500) were fed with diet containing 15% PKC in starter and finisher phase. The birds received a commercial broiler starter diet (21.84% crude protein) from days 1 to 21 and finisher diet (20.02% crude protein) from 22 to 41 days (Table 1). Thirty six birds were selected with uniform body weight (1837.80 ± 100 g) for this trial. The birds were randomly distributed into three treatments with six replicates in each treatment, and two birds per replicate. The chickens were deprived from food overnight with free access to drinking water in order to evacuate their GIT from any previous feed. After a fasting period, the birds were fed untreated PKC, FPKCa and FPKCb (Table 2). All the chickens were allowed free access to the test ingredients and drinking water for 4 days. The birds were slaughtered, and ileal digesta were gently squeezed and few drops of distilled water were added to the ileum to collect the digesta (Ravindran *et al.* 2005). The ileal digesta was collected from the Meckel's diverticulum to 1 cm before ileo-cecal junction (Loh *et al.* 2008).

Ileal digesta of birds were individually collected, and pooled within each replicate (pen) in plastic containers resulting in 6 composite samples per test ingredients. The digesta were immediately kept at -20°C and subsequently dried at 60°C. The dried ileal digesta samples were ground to pass through a 0.5 mm sieve and stored at -20°C for chemical analyses.

**Chemical analyses:** Amino acids content in test ingredients and ileal digesta were determined by high performance liquid chromatography (HPLC) with post-

Table 1. Ingredient and nutrient compositions of starter and finisher diet

Ingredient (%)	Starter	Finisher
Yellow corn	39.15	43.84
Palm oil	8.24	7.32
PKC <sup>1</sup>	15.00	15.00
Soybean meal 44% CP	28.54	24.60
Fish meal 55% CP	5.00	5.00
DCP <sup>2</sup> 18%	1.70	1.80
Calcium carbonate	0.37	0.37
DL-Methionine	0.25	0.28
L-Lysine	0.22	0.23
L-Threonine	0.00	0.03
Salt	0.30	0.30
Vitamin premix <sup>a</sup>	0.50	0.50
Mineral premix <sup>b</sup>	0.50	0.50
Toxin binder	0.14	0.14
Antioxidant <sup>c</sup>	0.01	0.01
Choline chloride	0.08	0.08
Total	100.00	100.00
<i>Calculated analyses<sup>d</sup></i>		
ME* (kcal/kg)	3000.71	3000.00
Methionine + Cystine (%)	0.95	0.90
Lysine (%)	1.30	1.21
Arginine (%)	1.54	1.43
Lys:Arg ratio	1;1.18	1: 1.18
Calcium (%)	1.08	1.14
Available phosphorus (%)	0.57	0.58
<i>Chemical analyses (%)</i>		
DM	94.11	91.00
CP	21.84	20.02
NDF <sup>3</sup>	19.30	19.12
ADF <sup>4</sup>	11.42	11.20
EE	10.30	11.73
Ash	4.77	5.10
NFE	58.16	58.47

<sup>a</sup>Provided per kg diet: vitamin A, 6670 IU; vitamin D<sub>3</sub>, 1000 IU; vitamin E, 23 IU; vitamin K<sub>3</sub>, 1.33 mg; cobalamin, 0.03 mg; Thiamin, 0.83 mg; riboflavin, 2.0 mg; folic acid, 0.33 mg; biotin, 0.03 mg; pantothenic acid, 3.75 mg; niacin, 23.30 mg; pyridoxine, 1.33 mg. <sup>b</sup>Provided per kg diet: Fe, 100.0 mg; Mn, 110.0 mg; Cu, 20.0 mg; Zn, 100.0 mg; I, 2.0 mg; Se, 0.20 mg; Co, 0.60 mg. <sup>c</sup>Butyrate hydroxytoluene (BHT). DM, dry matter; CP, crude protein; EE, ether extract; NFE, nitrogen free extract. <sup>1</sup>PKC, Palm kernel cake; <sup>2</sup>DCP, Dicalcium phosphate; <sup>3</sup>NDF, Neutral detergent fibre; <sup>4</sup>ADF, Acid detergent fibre; \*ME, Metabolizable energy. <sup>d</sup>Diets were formulated using FeedLIVE software (FeedLIVE 1.52, Thailand).

column derivatization (*O*-phthaldehyde) and fluorescence detection (Model 1100, Agilent Technologies, Inc., Santa Clara, CA). The samples were hydrolysed with 6 M hydrochloric acid (HCl) at 110°C for 24 h, whereas methionine and cysteine were hydrolysed as methionine sulfone and cysteic acid by oxidation with performic acid at 4°C for 16 h (Moore 1963). The concentration of N, crude fibre (CF), dry matter (DM) and ash were determined based

Table 2. Composition of test ingredients in the digestibility trial

Item	Test ingredients (%)		
	PKC	FPKCa	FPKCb
PKC <sup>1</sup>	90.30	0.00	0.00
FPKCa <sup>2</sup>	0.00	90.30	0.00
FPKCb <sup>3</sup>	0.00	0.00	90.30
Palm oil	6.00	6.00	6.00
CaCO <sub>3</sub>	1.70	1.70	1.70
Salt	0.40	0.40	0.40
Vitamin premix <sup>4</sup>	0.50	0.50	0.50
Mineral premix <sup>5</sup>	0.50	0.50	0.50
Choline-Cl	0.30	0.30	0.30
TiO <sub>2</sub>	0.30	0.30	0.30
Total	100.0	100.0	100.0

<sup>1</sup>PKC, Untreated palm kernel cake. <sup>2</sup>FPKCa, fermented by *P. polymyxa* ATCC 842. <sup>3</sup>FPKCb, fermented by *P. curdlanolyticus* DSMZ 10248. <sup>4</sup>Provided per kg diet: vitamin A, 6670 IU; vitamin D<sub>3</sub>, 1000 IU; vitamin E, 23 IU; vitamin K<sub>3</sub>, 1.33 mg; cobalamin, 0.03 mg; Thiamine, 0.83 mg; riboflavin, 2.0 mg; folic acid, 0.33 mg; biotin, 0.03 mg; pantothenic acid, 3.75 mg; niacin, 23.30 mg; pyridoxine, 1.33 mg. <sup>5</sup>Provided per kg diet: Fe, 100.0 mg; Mn, 110.0 mg; Cu, 20.0 mg; Zn, 100.0 mg; I, 2.0 mg; Se, 0.20 mg; Co, 0.60 mg.

on the method described by AOAC (1995). The CP was calculated as  $N \times 6.25$  AA, CP and other nutrients were presented in percentages as dry matter basis. Neutral detergent fibres (NDF) and acid detergent fibres (ADF) were determined based on the method described by Goering and Van Soest (1970). Titanium dioxide was determined in the feed and digesta based on the method described by Short *et al.* (1996).

**Calculation:** Apparent ileal digestibility (AID) of CP and AA was calculated based on CP, AA and TiO<sub>2</sub> concentrations of test ingredients and ileal digesta using the following formula (Son *et al.* 2014):

$$\text{AID} = 100 - [100 \times (\% \text{ TiO}_2 \text{ in feed} / \% \text{ TiO}_2 \text{ in digesta}) \times (\% \text{ nutrient digesta} / \% \text{ nutrient feed})]$$

**Statistical analysis:** Data were analysed using One-way ANOVA to compare the differences among the nutrients in the test ingredients. The experimental design was based on complete randomized design (CRD) in the digestibility trial. Data were analyzed by using General Linear Model procedure of the statistical analysis system (SAS 2003). Tukey's test was used to compare the means of treatment at 5% probability ( $P < 0.05$ ). The statistical model used for the trial was  $Y_{ijk} = \mu + T_{ij} + E_{ijk}$ , where,  $Y_{ijk}$ , observation;  $\mu$ , population mean;  $T_{ij}$ , effect of test ingredients;  $E_{ijk}$ , experimental error

## RESULTS AND DISCUSSION

The content of nutrients in the PKC and FPKC by cellulolytic bacteria are shown in Table 3. The process of SSF by cellulolytic bacteria led to increased levels of CP and AA in the FPKC compared to the untreated PKC. The CP increased ( $P = 0.0722$ ) from 16.43% in the untreated PKC

to 16.68% and 16.80% in FPKC by *P. curdlanolyticus* DSMZ 10248 and *P. polymyxa* ATCC 842, respectively. The AA were significantly increased ( $P < 0.05$ ) in the FPKC as compared to the untreated PKC. For instance, the concentration of isoleucine ( $P = 0.0239$ ), phenylalanine ( $P = 0.0192$ ), threonine ( $P = 0.0118$ ), histidine ( $P = 0.0150$ ), methionine ( $P = 0.0003$ ), arginine ( $P = 0.0312$ ), glycine ( $P = 0.0489$ ), aspartic acid ( $P = 0.0155$ ), glutamic acid ( $P = 0.0033$ ) and serine ( $P = 0.0150$ ) were increased in FPKC compared to untreated PKC. On the other hand, the SSF decreased the NDF ( $P < 0.0001$ ), ADF ( $P = 0.0003$ ), CF ( $P < 0.0001$ ), hemicellulose ( $P = 0.0010$ ) and cellulose ( $P = 0.0010$ ) in FPKC compared with untreated PKC. The AA content in the untreated PKC were consistent with Sulabo *et al.* (2013) and Son *et al.* (2014). It was also consistent with Muangkeow and Chinajariyawong (2009) except histidine, arginine, glutamate and tyrosine in which they were higher than the current PKC. The increase of CP and AA content in the FPKC by cellulolytic bacteria could be due to the reduction of NDF (12.87% in FPKCa and

Table 3. Nutrient content of palm kernel cake and fermented palm kernel cake by cellulolytic bacteria (dry matter basis)\*

Nutrient (%)	PKC	FPKCa <sup>1</sup>	FPKCb <sup>2</sup>	SEM <sup>3</sup>	P-values
Crude protein	16.43	16.80	16.68	0.04	0.0822
Dry matter	91.42	92.62	92.44	0.38	0.5228
Ash	4.74	4.67	4.80	0.13	0.2201
Neutral detergent fibre (NDF)	82.29 <sup>b</sup>	71.70 <sup>a</sup>	73.54 <sup>a</sup>	0.52	<0.0001
Acid detergent fibre (ADF)	51.48 <sup>b</sup>	47.27 <sup>a</sup>	47.45 <sup>a</sup>	0.58	0.0003
Hemicellulose	30.81 <sup>b</sup>	24.43 <sup>a</sup>	26.42 <sup>a</sup>	0.75	0.0010
Cellulose	35.55 <sup>b</sup>	31.85 <sup>a</sup>	31.41 <sup>as</sup>	0.62	0.0010
<i>Indispensable amino acids</i>					
Lysine	0.37	0.41	0.38	0.02	0.1325
Leucine	0.89	0.94	0.95	0.02	0.0551
Isoleucine	0.50 <sup>b</sup>	0.59 <sup>a</sup>	0.53 <sup>a</sup>	0.02	0.0239
Valine	0.69	0.78	0.72	0.03	0.1433
Phenylalanine	0.57 <sup>b</sup>	0.66 <sup>a</sup>	0.63 <sup>ab</sup>	0.02	0.0192
Threonine	0.41 <sup>b</sup>	0.51 <sup>a</sup>	0.46 <sup>ab</sup>	0.02	0.0118
Histidine	0.23 <sup>b</sup>	0.29 <sup>a</sup>	0.24 <sup>ab</sup>	0.02	0.0150
Methionine	0.22 <sup>b</sup>	0.27 <sup>a</sup>	0.26 <sup>a</sup>	0.01	0.0003
Arginine	1.60 <sup>b</sup>	1.76 <sup>a</sup>	1.69 <sup>ab</sup>	0.04	0.0312
Glycine	0.60 <sup>b</sup>	0.78 <sup>a</sup>	0.71 <sup>ab</sup>	0.04	0.0489
<i>Dispensable amino acids</i>					
Aspartic acid	1.12 <sup>b</sup>	1.27 <sup>a</sup>	1.23 <sup>ab</sup>	0.03	0.0155
Glutamic acid	2.48 <sup>b</sup>	2.80 <sup>a</sup>	2.76 <sup>a</sup>	0.08	0.0033
Proline	0.44 <sup>b</sup>	0.59 <sup>a</sup>	0.52 <sup>ab</sup>	0.02	0.0018
Serine	0.56 <sup>b</sup>	0.69 <sup>a</sup>	0.66 <sup>ab</sup>	0.04	0.0150
Tyrosine	0.25	0.24	0.24	0.01	0.4435
Cysteine	0.20	0.22	0.21	0.01	0.3632
Alanine	0.62	0.70	0.71	0.06	0.3892

<sup>1</sup>FPKCa, fermented palm kernel cake by *P. polymyxa* ATCC 842; <sup>2</sup>FPKCb, fermented palm kernel cake by *P. curdlanolyticus* DSMZ 10248; <sup>3</sup>Pooled standard error of means. \*Means  $\pm$  SEM. <sup>a,b</sup>Means with different superscripts in the same row differ significantly ( $P < 0.05$ ). n = 6 (6 replicates per treatment with 2 birds per replicate).

Table 4. Amino acids and crude protein digestibility of palm kernel cake and fermented palm kernel cake by cellulolytic bacteria (dry matter basis)\*

Nutrient (%)	PKC	FPKCa <sup>1</sup>	FPKCb <sup>2</sup>	SEM <sup>3</sup>	P-values
Crude protein	57.92 <sup>b</sup>	61.83 <sup>a</sup>	60.88 <sup>a</sup>	0.63	0.0014
<i>Indispensable amino acids</i>					
Lysine	65.94	69.57	70.63	2.12	0.1479
Leucine	65.47	68.04	63.89	1.39	0.1375
Isoleucine	69.59	70.47	66.39	2.58	0.5152
Valine	62.89 <sup>b</sup>	70.42 <sup>a</sup>	65.08 <sup>b</sup>	1.26	0.0022
Phenylalanine	68.77	70.76	68.51	1.96	0.6802
Threonine	61.38	64.98	61.69	1.73	0.2935
Histidine	56.99 <sup>b</sup>	71.50 <sup>a</sup>	64.83 <sup>ab</sup>	2.77	0.0076
Methionine	61.67 <sup>b</sup>	71.92 <sup>a</sup>	69.20 <sup>a</sup>	0.90	<0.0001
Arginine	75.75 <sup>b</sup>	81.15 <sup>a</sup>	76.30 <sup>b</sup>	0.95	0.0019
Glycine	47.44	45.52	52.96	4.16	0.4424
<i>Dispensable amino acids</i>					
Aspartic acid	56.87 <sup>b</sup>	64.30 <sup>a</sup>	61.74 <sup>a</sup>	1.20	0.0018
Glutamic acid	62.64 <sup>b</sup>	72.37 <sup>a</sup>	65.45 <sup>b</sup>	0.91	<0.0001
Proline	53.76	58.73	51.20	3.06	0.2401
Serine	65.76	69.78	67.58	2.10	0.4186
Tyrosine	59.04 <sup>b</sup>	67.58 <sup>a</sup>	61.93 <sup>ab</sup>	1.84	0.0155
Cysteine	33.34 <sup>b</sup>	41.45 <sup>a</sup>	37.46 <sup>ab</sup>	2.01	0.0393
Alanine	52.07 <sup>b</sup>	66.87 <sup>a</sup>	59.84 <sup>ab</sup>	2.49	0.0029

<sup>1</sup>FPKCa, Fermented palm kernel cake by *P. polymyxa* ATCC 842.<sup>2</sup>FPKCb, fermented palm kernel cake by *P. curdlanolyticus* DSMZ 10248.<sup>3</sup>Pooled standard error of means. \* <sup>a,b</sup>Means±SEM. Means with different superscripts in the same row differ significantly (P<0.05). n= 6 (6 replicates per treatment with 2 birds per replicate).

10.63% in FPKCb), ADF (8.18% in FPKCa and 7.83% in FPKCb) and NSPs.

The findings were in agreement with Muangkeow and Chinajariyawong (2009), who found that the rest of AA were increased when the PKC was fermented by *Aspergillus wentii* TISTR 3075. These findings were also consistent with Gao *et al.* (2013), who reported that SSF by *Lacobacillus brevis* led to increased CP and AA in soybean meal, or by *L. salivarius* in canola meal (Ahmed *et al.* 2014). The increase in CP in FPKC by *P. polymyxa* ATCC 842 may be due to its capability of producing proteolytic enzymes (Alvarez *et al.* 2006). Therefore, proteolytic enzymes may produce more peptides and AA, and consequently the digestible and soluble protein could be increased (Pranoto *et al.* 2013).

Digestibility of CP and AA in the PKC and FPKC by cellulolytic bacteria as is presented in Table 4. The digestibility of CP was significantly (P<0.05) increased in FPKC by cellulolytic bacteria as compared to the untreated PKC.

The CP digestibilities were 57.92%, 60.88% and 61.83% in the untreated PKC, FPKC by *P. curdlanolyticus* DSMZ 10248 and FPKC by *P. polymyxa* ATCC 842, respectively. There was also significant increase in the digestibilities of valine (P=0.0022), histidine (P=0.0076), methionine (P<0.0001), arginine (P=0.0019), aspartic acid (P=0.0018),

glutamic acid (P<0.0001), tyrosine (P=0.0150), cysteine (P=0.0393) and alanine (P=0.0029) in FPKC compared to the untreated PKC. Overall, there was improvement in the FPKC by cellulolytic bacterial cultures as mentioned above in terms of CP and AA digestibility.

The digestibility of CP and AA in the PKC is poorer compared to other common feedstuffs used in poultry feed such as soybean meal and yellow corn. The digestibility of CP in the PKC was consistent with Mustafa *et al.* (2004), who mentioned that digestibility of CP in the PKC was 53.37% in male Muscovy ducks. On the other hand, the digestibility of CP and AA were improved in the FPKC by cellulolytic bacteria, especially *P. polymyxa* ATCC 842.

The findings were in agreement with Sulabo *et al.* (2013), who reported that reduction in CP and AA digestibility in the growing pigs may be due to presence of high content of fibres. They also attributed the reduction of AID in the PKC to the high content of NDF, ADF and lignin. Sulabo *et al.* (2013) also reported that digestibility of CP was decreased by 0.3–0.8% as NDF increased by 1% in the diet. In addition, the negative effect of the soluble fibres in monogastrics is greater than the insoluble (Dégen *et al.* 2007, Sulabo *et al.* 2013). Sulabo *et al.* (2013) reported that PKC, with exception of guar meal, has greatest levels of  $\alpha$ -mannans among all feedstuffs. It was also claimed by Annison and Choct (1991) and Aya *et al.* (2013) that presence of NSPs in poultry diet depressed the nutrient utilization.

The improvement in CP and AA digestibility in FPKC by cellulolytic bacteria in the current study may be attributed to the reduction of NDF, ADF, hemicellulose and cellulose as a result of their capabilities of producing cellulolytic enzymes such as cellulase, xylanase and mannanase. These findings were consistent with Liu *et al.* (2013), who reported that *Bacillus licheniformis* derived protease led to better CP and AA digestibility of broiler chickens fed with sorghum-based diets. Additionally, the improvement in AID could be attributed to the cellulolytic enzymes (Ravindran *et al.* 2007, Selle *et al.* 2009, Masey-O'neill *et al.* 2014).

The presence of NSPs in poultry diets may lead to increase in the speed of food passage in the gastrointestinal tract (GIT). Therefore, the utilization of AA and other nutrients will be decreased. The findings were also in agreement with Lawal *et al.* (2010), who reported that reduction in NSPs of FPKC by fungi led to better nutrient digestibility in broiler chickens.

Based on the current results, it can be concluded that nutritive value of FPKC by cellulolytic bacteria improved. The NDF and ADF were decreased by 16.92%, 12.87% and 8.18%, respectively in FPKCa. The bacterial culture of *P. polymyxa* ATCC 842 showed higher ability in degradation and improvement of the PKC during SSF.

#### ACKNOWLEDGEMENTS

This research project was supported by Long-Term Research Grant Scheme (LRGS) given by the Ministry of Education, Malaysia.

## REFERENCES

- Alvarez-Cervantes J, Hern´andez-Dom´ınguez E M, Arana-Cuenca A, D´ıaz-God´ınez G and Mercado-Flores Y. 2013. Purification and characterization of xylanase SRXL1 from sporisorium reilianum grown in submerged and solid-state fermentation. *BioResources* **8**: 5309–18.
- Adebiyi A O and Olukosi O A. 2015. Apparent and standardised ileal amino acid digestibility of wheat distillers dried grains with solubles with or without exogenous protease in broilers and turkeys. *British Poultry Science* **56**: 239–46.
- Ahmed A, Zulkifli I, Farjam A S, Abdullah N, Liang J B and Awad E A. 2014. Effect of solid state fermentation on nutrient content and ileal amino acids digestibility of canola meal in broiler chickens. *Italian Journal of Animal Science* **13**: 410–14.
- Alimon A R. 2004. The nutritive value of palm kernel cake for animal feed. *Palm Oil Development* **40**: 12–14.
- Alshelmani M I, Loh T C, Foo H L, Lau W H and Sazili A Q. 2013. Characterization of cellulolytic bacterial cultures grown in different substrates. *The Scientific World Journal* **2013**: 6.
- Alshelmani M I, Loh T C, Foo H L, Lau W H and Sazili A Q. 2014. Biodegradation of palm kernel cake by cellulolytic and hemicellulolytic bacterial cultures through solid state fermentation. *The Scientific World Journal* **2014**: 8.
- Alshelmani M I, Loh T C, Foo H L, Sazili A Q and Lau W H. 2016a. Effect of feeding different levels of palm kernel cake fermented by *Paenibacillus polymyxa* ATCC 842 on nutrient digestibility, intestinal morphology, and gut microflora in broiler chickens. *Animal Feed Science and Technology* **216**: 216–24.
- Alshelmani M I, Loh T C, Foo H L, Sazili A Q and Lau W H. 2016b. Effect of feeding different levels of palm kernel cake fermented by *Paenibacillus polymyxa* ATCC 842 on broiler growth performance, blood biochemistry, carcass characteristics, and meat quality. *Animal Production Science* **57**: 839–48.
- Alvarez V M, Von Der Weid I, Seldin L and Santos A L S. 2006. Influence of growth conditions on the production of extracellular proteolytic enzymes in *Paenibacillus peoriae* NRRL BD-62 and *Paenibacillus polymyxa* SCE2. *Letters in Applied Microbiology* **43**: 625–30.
- Annisson G and Choct M. 1991. Anti-nutritive activities of cereal non-starch polysaccharides in broiler diets and strategies minimizing their effects. *World's Poultry Science Journal* **47**: 232–42.
- AOAC. 1995. Official Methods of Analysis (16th ed.). Arlington, VA. Association of Official Analytical Chemists.
- Aya V, Ayanwale B, Ijaiya A and Aremu A. 2013. Performance and nutrient digestibility in broiler chicks as influenced by multienzyme addition to starter diets containing palm kernel meal. *Biotechnology in Animal Husbandry* **29**: 93–104.
- Chen W L, Liang J B, Jahromi M F, Ho Y W and Abdullah N. 2013. Optimization of multi-enzyme production by fungi isolated from palm kernel expeller using response surface methodology. *BioResources* **8**: 3844–57.
- Dégen L, Halas V and Babinszky L. 2007. Effect of dietary fibre on protein and fat digestibility and its consequences on diet formulation for growing and fattening pigs: A review. *Acta Agriculturae Scandinavica, Section A — Animal Science* **57**: 1–9.
- Fadil M, Alimon A R, Meng G Y, Ebrahimi M and Farjam A S. 2014. Palm kernel cake as a potential ingredient in Muscovy ducks diet. *Italian Journal of Animal Science* **13**: 112–15.
- Gao Y L, Wang C S, Zhu Q H and Qian G Y. 2013. Optimization of solid-state fermentation with *Lactobacillus brevis* and *Aspergillus oryzae* for trypsin inhibitor degradation in soybean meal. *Journal of Integrative Agriculture* **12**: 869–76.
- Goering H K and Van Soest P J. 1970. Forage fibre analysis (apparatus, reagents, procedures and some applications). USDA. *Agricultural Handbook. No 379*
- Graminha E B N, Gonçalves A Z L, Pirola R D P B, Balsalobre M A A, Da Silva R and Gomes E. 2008. Enzyme production by solid-state fermentation: Application to animal nutrition. *Animal Feed Science and Technology* **144**: 1–22.
- Iluyemi F B, Hanafi M M, Radziah O and Kamarudin M S. 2006. Fungal solid state culture of palm kernel cake. *Bioresource Technology* **97**: 477–82.
- Iyayi E A and Davies B I. 2005. Effect of enzyme supplementation of palm kernel meal and brewer's dried grain on the performance of broilers. *International Journal of Poultry Science* **4**: 76–80.
- Khin H S. 2004. 'Evaluation of solid state fermentation by *Aspergillus niger* to improve the nutritive value of palm kernel cake for broilers.' Ph. D. Thesis. Universiti Putra Malaysia.
- Kocher A, Choct M, Porter M D and Broz J. 2002. Effects of feed enzymes on nutritive value of soyabean meal fed to broilers. *British Poultry Science* **43**: 54–63.
- Lateef A, Oloke J K, Gueguim Kana E B, Oyeniyi S O, Onifade O R, Oyeleye A O, Oladosu O C and Oyelami A O. 2008. Improving the quality of agro-wastes by solid-state fermentation: enhanced antioxidant activities and nutritional qualities. *World Journal of Microbiology and Biotechnology* **24**: 2369–74.
- Lawal T E, Iyayi E A, Adeniyi B A and Adaramoye O A. 2010. Biodegradation of palm kernel cake with multienzyme complexes from fungi and its feeding value for broiler. *International Journal of Poultry Science* **9**: 695–701.
- Liu S Y, Selle P H, Court S G and Cowieson A J. 2013. Protease supplementation of sorghum-based broiler diets enhances amino acid digestibility coefficients in four small intestinal sites and accelerates their rates of digestion. *Animal Feed Science and Technology* **183**: 175–83.
- Loh T C, Ling H G, Thanh N T, Foo H L, Rajion M A and David S I. 2008. Effects of feeding phytogenic substances and phytase on growth performance and nutrient digestibility of young broilers. *Journal of Applied Animal Research* **33**: 187–92.
- Marini A M, Daud M J, Noraini S, Jame'ah H and Azahan E A E. 2005. Performance of locally isolated microorganism in degrading palm kernel cake (PKC) fibre and improving the nutritional value of fermented PKC. *Journal of Tropical Agriculture and Food Science* **33**: 311–19.
- Masey-O'neill H V, Singh M and Cowieson A J. 2014. Effects of exogenous xylanase on performance, nutrient digestibility, volatile fatty acid production and digestive tract thermal profiles of broilers fed on wheat- or maize-based diet. *British Poultry Science* **55**: 351–59.
- McDonald P, Edwards R A and Greenhalgh J F D. 1995. *Animal Nutrition*. Longman Singapore Publishers Pvt Ltd.
- Mirawati Y R, Marlida Y and Kompang P. 2011. Evaluation of palm kernel cake fermented by *Aspergillus niger* as substituent for soybean meal protein in the diet of broiler. *International Journal of Poultry Science* **10**: 537–41.
- Moftah B, Lah T, Nawi A, Kadir M and Aliyu-Paiko M. 2012. Maximizing glucose production from palm kernel cake (PKC) from which residual oil was removed supercritically via solid

- state fermentation (SSF) method using *Trichoderma reesi* isolate *Pro-AI*. *The Internet Journal of Microbiology* **10**(1): 1–7.
- Moore S. 1963. On the determination of cystine as cysteic acid. *Journal of Biological Chemistry* **238**: 235–37.
- Muangkeow N and Chinajariyawong C. 2009. Determination of true amino acid digestibility and metabolizable energy in fermented palm kernel meal with *Aspergillus wentii* TISTR 3075 for chickens. *Walailak Journal of Science and Technology* **6**: 231–41.
- Mustafa M F, Alimon A R, Zahari M W, Idris I and Hair Bejo M. 2004. Nutrient digestibility of palm kernel cake for Muscovy ducks. *Asian Australasian Journal of Animal Science* **17**: 514–17.
- Ng W K and Chong K K. 2002. The nutritive value of palm kernel meal and the effect of enzyme supplementation in practical diets for red hybrid tilapia (*Oreochromis* sp.). *Asian Fisheries Science* **15**: 167–76.
- O'Mara F P, Mulligan F J, Cronin E J, Rath M and Caffrey P J. 1999. The nutritive value of palm kernel meal measured *in vivo* and using rumen fluid and enzymatic techniques. *Livestock Production Science* **60**: 305–16.
- Pranoto Y, Anggrahini S and Efendi Z. 2013. Effect of natural and *Lactobacillus plantarum* fermentation on *in-vitro* protein and starch digestibilities of sorghum flour. *Food Bioscience* **2**: 46–52.
- Ravindran V, Hew L I, Ravindran G and Bryden W L. 2005. Apparent ileal digestibility of amino acids in dietary ingredients for broiler chickens. *Animal Science* **81**: 85–97.
- Ravindran V, Tilman Z V, Morel P C H, Ravindran G and Coles G D. 2007. Influence of  $\alpha$ -glucanase supplementation on the metabolisable energy and ileal nutrient digestibility of normal starch and waxy barleys for broiler chickens. *Animal Feed Science and Technology* **134**: 45–55.
- SAS. 2003. *Statistical Analytical System*. SAS Institute Inc., Cary, NC, USA.
- Selle P H, Ravindran V and Partridge G G. 2009. Beneficial effects of xylanase and/or phytase inclusions on ileal amino acid digestibility, energy utilisation, mineral retention and growth performance in wheat-based broiler diets. *Animal Feed Science and Technology* **153**: 303–13.
- Short F, Gorton P, Wiseman J and Boorman K. 1996. Determination of titanium dioxide added as an inert marker in chicken digestibility studies. *Animal Feed Science and Technology* **59**: 215–21.
- Son A R, Hyun Y, Htoo J K and Kim B G. 2014. Amino acid digestibility in copra expellers and palm kernel expellers by growing pigs. *Animal Feed Science and Technology* **187**: 91–97.
- Sukaryana Y, Atmomarsono U, Yunianto V D and Supriyatna E. 2010. Bioconversions of palm kernel cake and rice bran mixtures by *Trichoderma viride* toward nutritional contents. *International Journal of Science and Engineering* **1**: 27–32.
- Sulabo R C, Ju W S and Stein H H. 2013. Amino acid digestibility and concentration of digestible and metabolizable energy in copra meal, palm kernel expellers, and palm kernel meal fed to growing pigs. *Journal of Animal Science* **91**: 1391–99.
- Sundu B and Dingle J. 2002. Use of enzymes to improve the nutritional value of palm kernel meal and copra meal. *Proceeding of the Queensland Poultry Science Symposium*, Australia.