



Sero-epidemiology of Bluetongue and Caprine arthritis-encephalitis in goats of middle Indo-Gangetic plains

PANKAJ KUMAR¹, KAUSHAL KISHORE RAJAK², SANKARIAH JEGAVEERA-PANDIAN³, DINESH MAHTO⁴, RASHMI REKHA KUMARI⁵, RAMA KRISHNA ROY⁶ and ABHAY KUMAR⁷

ICAR Research Complex for Eastern Region, Patna, Bihar 800 014 India

Received: 16 June 2017; Accepted: 25 July 2017

ABSTRACT

The study was undertaken in middle of Indo-Gangetic plains of India. Bluetongue (BT) is endemic throughout the country. However, Caprine arthritis-encephalitis virus (CAE) is sporadically reported from India. The study was undertaken to evaluate the burden of these viral diseases of goat due to unavailability of systemic study in the region. Commercial ELISA kits were used for the study. BTV antibody sero-surveillance was undertaken in 504 random serum samples and Lentivirus antibody sero-surveillance for CAE in 280 random serum samples. None of the goats covered under the study had any form of clinical manifestations of these diseases. The study revealed moderate (15.64%) sero-positivity of BTV antibody and only two cases of sero-positivity of CAE. Two positive cases of CAE in Sirohi breed was brought from Rajasthan. Overall sero-positivity of BTV antibody in male was higher than female goats. The binary regression of circulating BTV antibody was significant for Sirohi and Black Bengal breeds under the study indicating breed differences.

Key words: Sero-epidemiology, Bluetongue, Caprine arthritis-encephalitis, Goat, Indo-Gangetic plains

The middle Indo-Gangetic plain is situated between 24°–27° N latitude and 82°–88° E longitude. It possess about 7.63% of India's total goat population and ranks 5th in country. The region is divided into Zone 1 (North West alluvial plains), Zone 2 (North East alluvial plains) and Zone 3 (South Bihar alluvial plains). Goat rearing in the region is mostly restricted to small and marginal farmers and is source of livelihood. Bluetongue (BT) is an important endemic viral disease of goat transmitted by *Culicoides* species and approximately 22 different serotypes has been reported based on isolation and serology. Rao *et al.* (2016) has isolated 13 serotypes from India. It is mostly asymptomatic in goats. Caprine arthritis and encephalitis (CAE) of Lentivirus (LV) in goats causes significant economic losses and hampers exports. CAE in goats is sporadically reported. Most CAE infections in goats are subclinical (Larruskain and Jugo 2013). Sero-

epidemiological studies for BT and CAE in ruminants has not been reported from middle Indo-Gangetic plains. The sero-epidemiology of diseases in particular geographical area is important for planning effective prevention and control measures. Thus the present study was undertaken to know the sero-prevalence of BT and CAE in goats of middle Indo-Gangetic plains.

MATERIALS AND METHODS

Study area: A total of eight districts were randomly selected for the study representing all the agro-climatic zone of the region viz. Sitamarhi and Vaishali in zone 1; Purnia and Katihar in zone 2; Buxar, Patna, Gaya and Jehanabad in zone 3. This study was conducted over a period of three years from February 2013 to January 2016.

Serum samples: Goats of all ages, breed and gender were selected randomly from the study area. Data were collected through pre-tested schedule developed for the study. Blood (5 ml) samples were collected from jugular vein using plain vacutainers or sterile disposable syringes and serum was harvested. With proper identification, serum samples were transported to laboratory under refrigeration in cool box and stored at –20°C till further analysis. A total of 504 random samples were used for BTV antibody sero-epidemiology and 280 random samples for LV antibody sero-epidemiology.

Assay for Bluetongue antibodies: Sero-epidemiology of BTV circulating antibody was done using monoclonal

Present address: ¹Senior Scientist (Pankaj.Kumar1@icar.gov.in), ³Scientist (jeganicar@gmail.com), Veterinary Medicine, Division of Livestock and Fisheries Management; ²Scientist (kaushalvirol@gmail.com), Veterinary Virology, Biological Products Division, ICAR-IVRI, Izatnagar, Bareilly, UP. ⁴SMS (drdineshgy@gmail.com), Animal Science, KVK, Gandhar, Banbhuganj, Jehanabad, Bihar. ⁵Assistant Professor (rrkvet@gmail.com), Bihar Veterinary College, Patna, Bihar. ⁶Programme Coordinator (dramakrishnaroy@gmail.com), KVK, Sipaya, Gopalganj, Bihar. ⁷Principal Scientist (akumar1904@rediffmail.com), Division of Socio-Economics and Extension.

antibody- Competitive Enzyme Linked Immunosorbent Assay (MAb c-ELISA) kits manufactured by Veterinary Medical Research and Development (VMRD), USA with a claimed specificity of 99.0% and sensitivity of 100%. Optical density (OD) value was measured by ELISA reader (ECIL Microscan MS5608A) at 620 nm wavelength. Sero-positivity was recorded based on test validation method provided by the manufacturer.

Assay for Lentivirus antibodies: Sero-epidemiology of CAE based on detection of circulating antibody of LV by measuring the titre using small ruminant Lentivirus antibody c-ELISA kit manufactured by Veterinary Medical Research and Development (VMRD), USA. The kit has 99.0% specificity and 100% sensitivity for LV antibodies. Optical density (OD) value was measured by ELISA reader (ECIL Microscan MS5608A) at 620 nm wavelength. Sero-positivity was recorded based on test validation method provided by the manufacturer.

Statistical analysis: Apparent prevalence was calculated by number of sero-positivity divided by total number of serum samples tested. Corrected prevalence was calculated using formula described (Kumar *et al.* 2017). Results obtained from c-ELISA test were statistically analysed for testing the significance by analysis of variance and odds ratio of age, gender, zones and breeds of goat using logistic regression model.

RESULTS AND DISCUSSION

The present study was carried out in three agro-climatic zones of middle Indo-Gangetic plains for investigating sero-positivity of BT and CAE in goats, its percent prevalence and physiological state, gender and/or breed as risk factors. None of the goats under the study had any form of reported clinical manifestations of either BT or CAE. The study revealed moderate sero-positivity of BTV antibody (apparent prevalence of 15.48% and corrected prevalence of 15.64%) and only two cases of sero-positivity for LV antibody corresponding to CAE infection. In neighbouring state of Jharkhand, comparatively higher sero-positivity (43.33%; n=91) of circulating BTV antibody had been

reported in goats (Tigga *et al.* 2015). Similar reports from Odisha and Asom also had higher sero-positivity (31.25% and 31.79%, respectively) of BTV antibody in goats (Joardar *et al.* 2013, 2014). The sero-prevalence of circulating BTV antibody was comparatively lower in goats of Bihar than those reported from Asom, Odisha and Jharkhand. It can be inferred that this lower prevalence could be due to possible low vector population and/or comparatively larger sample size in this study. The non-significantly higher prevalence % of circulating BTV antibody were observed in Purnia district of Zone 2 (Table 1). However, the binary regression between circulating BTV antibody and sero-positivity in Zone 2 was significant. Higher incidence of BTV sero-positivity was reported from Zone 2 could be attributed to seasonally water-logged topography with high organic content which favours the breeding and survival of *Culicoides* (Chand *et al.* 2015). The Zone 2 also has comparatively higher goat population (9.0% of the total goat population in Purnia and Katihar) compared to other districts of Bihar (DAHD 2012). Higher prevalence % of BTV antibody was also observed in Gaya district of zone 3 attributed to very high summer temperature which favours vector movement. Circulating BTV antibody was highest in Sirohi breed followed by Black Bengal type and other breeds under study (Table 2). The significant binary regression was observed between circulating BTV antibody and breeds of Sirohi and Black Bengal type (Table 3). The higher incidence of this circulating virus in non-native goat breed (Sirohi) indicates possible spread of diseases due to inter-state marketing of goat population. The variation in sero-positivity amongst breed may also be attributed to variable sample size of breeds and influx of Sirohi breed goat population from other states. Gender and age as risk factor were analysed for determining sero-positivity of BTV antibody. Overall sero-positivity of BTV antibody in male was higher than female goats though the probability of higher circulating BTV antibody in male was non-significantly different (Table 3). However, circulating BTV antibody was higher in adult compared to kids of any gender and the binary regression

Table 1. Agro-climatic zone-wise sero-prevalence of BTV antibody in goats of Bihar

Zone	District	Female (n=351)		Male (n=153)		Apparent prevalence %	Corrected prevalence %	Zone prevalence %
		Kid (n=42)	Adult (n=309)	Kid (n=29)	Adult (n=124)			
1	Vaishali (n= 56)	0.00 (n=5)	13.79 (n=29)	0.00 (n=4)	11.11 (n=18)	10.71	10.83	12.37 ^a
	Sitamarhi (n=41)	0.00 (n=1)	15.00 (n=20)	0.00 (n=3)	17.65 (n=17)	14.63	14.79	
2	Purnia (n= 43)	0.00 (n=4)	29.63 (n=27)	0.00 (n=2)	10.00 (n=10)	20.93	21.15	19.40 ^a
	Katihar (n=24)	0.00 (n=2)	17.65 (n=17)	0.00	20.00 (n=5)	16.67	16.85	
3	Patna (n=112)	0.00 (n=8)	12.50 (n=56)	0.00 (n=10)	15.79 (n=38)	11.61	11.74	14.71 ^a
	Jehanabad (n=97)	0.00 (n=9)	19.44 (n=72)	0.00 (n=3)	7.69 (n=13)	15.46	15.63	
	Gaya (n=90)	0.00 (n=9)	18.18 (n=66)	20.00 (n=5)	20.00 (n=10)	16.67	16.85	
	Buxar (n=41)	0.00 (n=5)	19.05 (n=21)	50.00 (n=2)	15.38 (n=13)	17.07	17.25	
	Total (n=504)	0.00	8.16±1.87	8.75±6.39	14.70±1.64	15.48±1.15	15.64±1.16	15.49±2.07
	Confidence interval at 95%	-	13.74 to 22.57	-6.36 to 23.86	10.82 to 8.58	12.76 to 18.18	12.90 to 18.37	6.60 to 24.39

Values with different superscript differ significantly ($P \leq 0.05$) and values in parenthesis indicates sample size.

Table 2. Breed-wise prevalence % of BTV antibody in goats of different districts of Bihar

Zone	District	Black Bengal type (n=231)	Sirohi (n=68)	Others (n=205)
1	Vaishali (n=56)	6.45 (n=31)	50.00 (n=2)	13.04 (n=23)
	Sitamarhi (n=41)	21.05 (n=19)	-	9.09 (n=22)
2	Purnia (n=43)	15.79 (n=19)	100.00 (n=3)	14.29 (n=21)
	Katihar (n=24)	23.08 (n=13)	-	9.09 (n=11)
	Patna (n=112)	9.84 (n=61)	16.67 (n=18)	12.12 (n=33)
3	Jehanabad (n=97)	17.07 (n=41)	40.00 (n=15)	4.88 (n=41)
	Gaya (n=90)	16.67 (n=36)	54.55 (n=11)	6.98 (n=43)
	Buxar (n=41)	9.09 (n=11)	26.32 (n=19)	9.09 (n=11)
	Total	14.88 ± 2.09 ^a	47.92 ± 11.93 ^b	9.82 ± 1.12 ^c
	Confidence interval at 95%	9.94 to 19.82	17.27 to 78.58	7.18 to 12.46

Values with different superscript differ significantly ($P \leq 0.05$) and values in parenthesis indicates samples size.

between BTV virus antibody and age was highly significant ($P=0.006$) with high odds ratio. In general, intact males were not sold for meat purpose and usually retained for longer period in the region. This may be the possible reason for higher sero-positivity of BTV antibody in addition to lesser representation of male goat population in the study. Thus, the prevalence of BT was not affected markedly by the sex. The gender as risk factor varied non-significantly for BTV antibody and corroborated with findings of Shringi and Shringi (2005) and Bitew *et al.* (2013). Higher sero-positivity of BTV in adult goats compared to kids indicates possible temporal factor for exposure to the BTV in adults. The present finding of higher sero-positivity in adults was in agreement to the findings of Christi (2010) who found that flock level sero-prevalance was higher among the adult population (25.93%) than young animals (3.7%). In contrast to our findings, Bitew *et al.* (2013) reported that risk factors like sex, age and body condition had no significant impact on the sero-prevalence. Sero-positivity of BTV antibody in kids also indicates possibility of trans-placental transmission of infection in agreement to the findings of Belbis *et al.* (2013).

Circulating LV antibody for CAE in goats was negative in most samples, except for the two cases of sero-positivity found in adult female Sirohi goats newly introduced in village Tetariya (24°47'27.8"N and 85°04'06.0"E) in Manpur block of Gaya district under zone 3. The goats were procured from middle man which were said to be brought from Rajasthan. Thus, apparent sero-prevalence of LV antibody circulating was very low (0.71%). Further analysis of data was not done for CAE. Similar low prevalence

Table 3. Binary logistic regression of BTV sero-positivity and zones, breeds, gender and age

Log regression of BT with	B	S.E.	Z	Sig.	Odds ratio Exp (B)
Gender	-0.16	0.30	-0.53	0.59	0.85
Age	2.04	0.74	2.76	0.006*	7.71
Black Bengal breed	0.53	0.31	1.70	0.09*	1.46
Sirohi breed	1.95	0.38	5.16	0.00*	5.22
ND breeds			Omitted		
Zone 1	0.20	0.36	0.57	0.57	1.23
Zone 2	0.54	0.36	1.49	0.14*	1.72
Zone 3			Omitted		
Constant	-4.37	0.78	-5.61	0.00	0.00

Log likelihood, -192.72; Pseudo R^2 , 0.098; B, Regression coefficient; SE, Standard error; Z, random variable that has a standard normal distribution and calculated as B/S.E; Sig., level of significance

(3.33%) of CAE in goats was reported by Waseem *et al.* (2015). Our findings of sero-positivity in female goats was in agreement with the findings of Waseem *et al.* (2015).

This indicates the importance of such surveillance study for emerging animal diseases which were not reported and occurring sporadically. Periodical surveillance will bring awareness about possible state migration of new diseases and consequently help the Government agencies to take suitable action for preventing the spread of these diseases and from becoming endemic.

ACKNOWLEDGEMENT

Authors acknowledge support and cooperation of Director, ICAR- Research Complex for Eastern Region, Patna for providing necessary funds and facilities for undertaking the institute project. Support rendered by Head, Division of Livestock and Fisheries Management is also gratefully acknowledged. Support provided by Programme Coordinators of the Krishi Vigyan Kendras during survey and sampling is duly acknowledged.

REFERENCES

- Belbis G, Bréard E, Cordonnier N, Moulin V, Desprat A, Saillieu C, Viarouge C, Doceul V, Zientara S and Millemann Y. 2013. Evidence of transplacental transmission of bluetongue virus serotype 8 in goats. *Veterinary Microbiology* **166**(3): 394–404.
- Bitew M, Nandi S, Ravishankar C and Somvanshi R. 2013. Serological and molecular evidence of bluetongue in sheep and goats in Uttar Pradesh, India. *African Journal of Biotechnology* **12**(19): 2699–704.
- Chand K, Biswas S K, Pandey A B, Muthuchelvan D and Mondal B. 2015. Bluetongue in India: A review. *Advances in Animal and Veterinary Sciences* **3**(11): 605–12.
- Christie M. 2010. Prevalence and risk factors associated with bluetongue virus among Colorado sheep flocks. M.Sc Thesis, Department of Clinical Sciences, Colorado State University, pp. 45–60.
- DAHD. 2012. 19th Livestock Census District Wise Report, Vol. I: 186–229.

- Joardar S N, Barkataki B, Halder A, Lodh C and Sarma D. 2013. Seroprevalence of bluetongue in north eastern Indian state of Assam. *Veterinary World* **6**(4): 196–99.
- Joardar S N, Halder A, Maity A, Mishra S K and Lodh C. 2014. Seroprevalence of bluetongue in ruminants of Orissa. *Indian Journal of Comparative Microbiology, Immunology and Infectious Diseases* **35**(2): 58–60.
- Kumar P, Sinha B S, Roy R K, Kumari R R and Kumar A. 2017. Peste des petits ruminants in goats: Sero-epidemiological study in middle Indo-Gangetic plains. *Indian Journal of Animal Sciences* **87**(4): 418–21.
- Larruskain A and Jugo B M. 2013. Retroviral infections in sheep and goats: small ruminant lentiviruses and host interaction. *Viruses* **5**(8): 2043–61.
- Rao P P, Hegde N R, Reddy Y N, Krishnajyothi Y, Reddy Y V, Susmitha B, Gollapalli S R, Putty K and Reddy G H. 2016. Epidemiology of Bluetongue in India. *Transboundary and Emerging Diseases* **63**(2): 151–64.
- Shringi S and Shringi B N. 2005. Comparative efficacy of standard AGID, CCIE and competitive ELISA for detecting bluetongue virus antibodies in indigenous breeds of sheep and goats in Rajasthan, India. *Journal of Veterinary Science* **6**(1): 77–79.
- Tigga P, Joardar S N, Halder A, Lodh C, Samanta I, Isore D V, Batabyal K and Dey S. 2015. Seroprevalence of bluetongue in ruminants of Jharkhand. *Veterinary World* **8**(3): 346–49.
- Waseem A, Pawaiya R V S, Singh R, Gupta V K, Rajukumar K, Mir M S and Aamir S. 2015. Seroprevalence of caprine arthritis encephalitis virus infection (CAEV) in Indian goats. *Indian Journal of Veterinary Pathology* **39**(1): 15–19.