



## Influence of different culture media on embryo development following *in vitro* fertilization of caprine oocytes

PARUL YADAV<sup>1</sup>, S D KHARCHE<sup>2</sup>, A K GOEL<sup>3</sup> and S K JINDAL<sup>4</sup>

ICAR-Central Institute for Research on Goats, Mathura, Uttar Pradesh 281 122 India

Received: 31 January 2017; Accepted: 21 July 2017

**Key words:** Caprine oocytes, Cleavage, *In vitro* fertilization, *In vitro* maturation, Morula

Major developmentally important events take place during development of embryo from post fertilization to the blastocyst stage. Embryo yield and survival usually differ between the different culture systems and culture media (Camargo *et al.* 2006). *In vitro*, the pre-implantation development of the embryo occurs in close relation with the culture medium. This demands specific balanced conditions in the medium to allow embryonic metabolism (Garcia *et al.* 2016). A number of different defined culture media have been extensively evaluated in an effort to improve the quality of embryos produced *in vitro* and to mimic the physiological conditions that embryos would have *in vivo*. The aim of preparing defined media is to minimize the variation in unknown components that occurs naturally in serum. This has the advantage of providing a simplified technique that enhances cryoresistance of embryos. This is due to avoiding the supplementation of biological components such as FCS and bovine serum albumin (BSA) that may increase the potential risks of contamination (Neira *et al.* 2010). The most common defined media used in *in vitro* culture systems are SOF (synthetic oviductal fluid), KSOM, and tissue culture medium (TCM199) (Licola *et al.* 2015).

Keeping this in mind, the objective of the present study was to compare three different defined culture medium to evaluate the developmental competence of embryos. Ovaries were collected from the local abattoir and transported within 4 h to the laboratory in warm saline (37°C) containing 100 IU penicillin-G and 100 µg streptomycin sulfate per ml. The oocytes were collected from each goat ovary and graded as per the method of Kharche *et al.* (2008). Selected oocytes were then washed three times with Oocyte Maturation Medium (OMM) containing TCM-199 supplemented with 10% FBS, hormones (5 µg/ml FSH, 5 µg/ml LH and 1 µg/ml estradiol-17β), 10 ng/ml EGF, Sodium Pyruvate 0.25 mM, Glutamine 100 µg/ml, LH 10 µg/ml, FSH 5 µg/ml, BSA 3 mg/ml and

Gentamycin 50 µg/ml). Washed Cumulus Oocyte Complexes (COC's) were then allowed for *in vitro* maturation at 38.5°C in 5% CO<sub>2</sub> in air for 27 h.

After the maturation, oocytes were separated from cumulus and corona cells by treatment with 0.1% hyaluronidase. The oocytes were washed 8–10 times in Fertilization medium (TALP solution containing 20% oestrous goat serum and 10 µg/ml heparin) and kept in 50 µl drop in culture petriplate.

Fresh semen was collected by artificial vagina (AV), from a fertile purebred Sirohi buck. Semen was evaluated for volume, colour, and consistency, gross and progressive motility. An amount of 50 µl neat semen was diluted with 5 ml of sperm TALP containing 20% EGS and 10 µg/ml heparin and washed 2 - 3 times by centrifugation at 1200 rpm for 5 min. The supernatant was discarded in each washing. Finally, 50 µl pellet of sperm was diluted with 1 ml of fertilization media. Fertilization drop containing oocytes were inseminated with 25 to 50 µl of finally diluted semen to obtain a sperm concentration of 1–2×10<sup>6</sup> sperm/ml. Gametes were co incubated for 18 h in a CO<sub>2</sub> incubator. Oocytes-sperm after 18 h of co-incubation were washed in embryo development medium for 15–20 times and randomly divided in to three groups. Group 1 (n=1511) fertilized oocytes were cultured in TCM-199 containing 10% NCS and 4 mg/ml bovine serum albumin (BSA) for 10 days; Group 2 (n=1333) fertilized oocytes were cultured in synthetic oviductal fluid (SOF) containing 10% NCS and 4 mg/ml BSA for 10 days and Group 3 (n=1000) fertilized oocytes were cultured in potassium simplex optimization medium (KSOM) containing 10% NCS and 4 mg/ml BSA for 10 days. Presumptive zygotes were cultured under mineral oil in 5% CO<sub>2</sub> in air with high humidity at 38.5°C. Presumptive zygotes were observed for cleavage and further *in vitro* embryo development. Developing embryos were evaluated under inverted phase contrast microscope (Fig. 1).

All data were analysed by one-way ANOVA, developmental data were arc sin transformed. The level of significance for cleavage and embryo development was recorded at 5% level of confidence (Snedecor and Cochran 1989).

Present address: <sup>1</sup>Scientific Officer (parul\_biochem@rediffmail.com), AUSIC, Amity University, Jaipur. <sup>2,3,4</sup>Principal Scientist (kharche62@gmail.com, anil@cirg.res.in, jindal@cirg.res.in), Animal Physiology and Reproduction Division.




Fig. 1. 1. Immature oocyte 2. Mature oocytes 3. MII oocyte with polar body 4. MII oocyte with two shiny chromatin spots 5. 2–8 cell stage embryos 6. Orulae.

The cleavage rate, morulae and blastocyst percentage for groups 1, 2, and 3 were 13.6, 10.14 and 2.41%; 11.7, 7.6 and 3.84% and 31.4, 12.73 and 6.05%, respectively. The cleavage rate was significantly higher ( $P < 0.01$ ) for group 3 than for groups 1 and 2. Similarly, embryo development up to morula stage was higher ( $P < 0.05$ ) in KSOM compared with TCM-199 and SOF medium. In our previous studies, we obtained cleavage rate, 2 cell, 4 cell, 8–16 cell, morulae and blastocysts as 44%, 17%, 25%, 33.11%, 16% and 2%, respectively in defined KSOM medium as compared to complex embryo development media TCM 199 (Kharche *et al.* 2011). The effect of three different media RVCL, mSOF, G1.G2 for culture of intraspecies and interspecies cloned goat embryos was nonsignificant (Khan *et al.* 2013).

The effects of supplementation of synthetic oviduct fluid (SOF) with glucose and Minimal Essential Medium (MEM) vitamins was assessed during the first step of the *in vitro* culture (SOF C1) of ovine zygotes and found vitamins must be present for optimal numbers of embryos to reach the blastocyst stage (Karami *et al.* 2012). There are some research teams using synthetic oviductal fluid (SOF) medium instead of TCM199 for sheep (Shabankareh and Akhondi 2012, Manacha *et al.* 2016) and for goat embryos (Herrick *et al.* 2004).

The following study showed that the use of KSOM medium supplemented with 5% FBS after 3 days of culture can improve the blastocyst formation rate of SCNT (Somatic Cell Nuclear Transfer) embryos and reduce the incidence of apoptosis compared to other two-step culture media supplemented with Bovine Serum Albumin (Aries *et al.* 2013). In another study, KSOM medium was found more appropriate than TCM 199 medium to culture the SCNT (Somatic Cell Nuclear Transfer) embryos from the pronuclear to the cleavage stage (Wan *et al.* 2013). The most common media used in bovine *in vitro* culture systems are SOF (synthetic oviduct fluid), KSOM, and CR1aa (Kocyigit *et al.* 2016).

So the results indicated that the KSOM medium is better culture medium as compared to TCM- 199 and SOF medium as far as cleavage rate and embryo development is concerned.

#### SUMMARY

The present study was designed to compare the developmental competence of *in vitro* fertilized caprine embryos at different cell stages in three different culture medium (TCM-199, mSOF, KSOM). Oocytes were collected from ovaries and matured in TCM-199 medium

containing FSH (5 mg/ml), LH (5 mg/ml) and estradiol-17b (1 mg/ml), supplemented with 20% fetal bovine serum at 38.5°C and 5% CO<sub>2</sub> in an incubator under humidified air for 27 h. After 27 h of *in vitro* maturation (IVM), oocytes were denuded, washed and co-incubated with 50 µl neat semen diluted in 5 ml of sperm TALP containing 20% EGS and 10 µg/ml heparin. Oocytes-sperm after 18 h of co-incubation were randomly divided into three groups: Group 1 (TCM-199 containing 10% NCS and 4 mg/ml bovine serum albumin), Group 2 (Synthetic oviductal fluid containing 10% NCS and 4 mg/ml BSA) and Group 3 (Potassium simplex optimization medium (KSOM) containing 10% NCS and 4 mg/ml BSA) and found the cleavage rate, morulae and blastocyst percentage for groups 1, 2, and 3 were 13.6, 10.14 and 2.41%; 11.7, 7.6 and 3.84% and 31.4, 12.73 and 6.05%, respectively. Statistical analysis revealed that the cleavage rate and embryo development up to morula and blastocyst stage in group 3 (KSOM) was significantly higher as compared to group 1 and group 2. The results indicated that the KSOM medium was better among all three medium for embryo development.

#### ACKNOWLEDGEMENT

The authors wish to thank the Director ICAR-CIRG, Makhdoom, Farah, Mathura, UP, India for providing the facilities needed.

#### REFERENCES

- Arias E Maria, Ross J Pablo and Felmer N Ricardo. 2013. Culture medium composition affects the gene expression pattern and *in vitro* development potential of bovine somatic cell nuclear transfer (SCNT) embryos. *Biological Research* **46**: 452–62.
- Camargo L S A, Viana J H M, Ferreira S A W F, Ramos A M and Filho Vale V R. 2006. Factors influencing *in vitro* embryo production. *Animal Reproduction* **3**: 19–28.
- Garcia Moreno Diego, Nera Alberto, Briand Lamia, Topie Emmanuel, Bencharif Djemil and Tainturier Daniel. 2016. Recent advances on *in vitro* fertilization. Chapter 2. *In Vitro Fertilization (IVF)*. ISBN 978–93–86337–13–9. Avid Science Publishers.
- Herrick J R, Behboodi E, Memili E, Blash S, Echelard Y and Krisher R L. 2004. Effect of macromolecule supplementation during *in vitro* maturation of goat oocytes on developmental potential. *Molecular Reproduction and Development* **69**: 338–46.
- Karami S H and Akhondi M. 2012. Effect of glucose levels and MEM vitamins during the first step of IVC in synthetic oviduct fluid medium on *in vitro* development of ovine zygotes. *Small Ruminant Research* **106**: 54–58.
- Khan F A, Bhat M H, Yaqoob S H, Waheed S M, Naykoo N A,

- Athar H, Khan H M, Fazili M R, Ganai N A, Singla S K and Shah R A. 2014. *In vitro* development of goat-sheep and goat-goat zona-free cloned embryos in different culture media. *Theriogenology* **81**(3): 419–23.
- Kharche S D, Goel A K, Jindal S K, Sinha N K and Yadav P. 2008. Effect of somatic cells co-culture on cleavage and development of *in vitro* fertilized caprine embryos. *Indian Journal of Animal Sciences* **78**(7): 686–92.
- Kharche S D, Goel A K, Jindal S K, Goel P and Jha B K. 2011. Birth of twin kids following transfer of *in-vitro* produced goat embryos. *Indian Journal of Animal Sciences* **81**(11): 1132–34.
- Kocyigit Alper. 2016. A review of *in vitro* culture systems in bovine reproductive biotechnologies. *Journal of Veterinary Research and Animal Husbandry* **1**(1): 101.
- Lekola Molvia Podile Khomotso. 2015. Improvement of cattle oocyte retrieval techniques and hormonal influence on *in vitro* embryonic development. A mini-dissertation submitted in partial fulfillment of the requirements for the degree of Master in Science to Agriculture University of Limpopo, South Africa.
- Menchaca A, Barrera N, Dos PC Santos Neto, Cuadro F and Crispo M. 2016. Advances and limitations of *in vitro* embryo production in sheep and goats. *Animal Reproduction* **13**(3): 273–78.
- Neira J A, Tenturier D, Pena M A and Martal J. 2010. Effect of association of IGF, IGF-2, bFGF, TGF-B1, GM-CSF and LIF on the development of bovine embryos produced *in vitro*. *Theriogenology* **3**: 95–604.
- Shabankareh H K and Akhondi M. 2012. Effect of glucose levels and MEM vitamins during the first step of IVC in synthetic oviduct fluid medium on *in vitro* development of ovine zygotes. *Small Ruminant Research* **106**: 54–58.
- Snedecor G W and Cochran W G. 1989. *Statistical Methods*, 6<sup>th</sup> edition. The Iowa State University Press, USA.
- Wan YongJie, Zhang Yanli, Zhou Zhengrong, Jia Ruoxin, Meng LI, Huang Rong, Wang Ziyu, Yan Yibo, You Jihao, Zhang Guomin and Wang Feng. 2013. Effect of potassium simplex optimization medium KSOM and embryo screening on the production of human lactoferrin transgenic cloned dairy goats. *African Journal Of Biotechnology* **12**(49): 6887–93.