



Effect of certain factors on first lactation peak yield and days to attain peak yield in Frieswal cattle

T V RAJA¹, R KUMAR², S K RATHEE³, B PRAKASH⁴ and U SINGH⁵

ICAR-Central Institute for Research on Cattle, Meerut Cantt, Meerut, Uttar Pradesh 250 001 India

Received: 20 March 2017; Accepted: 21 July 2017

Key words: Days to attain peak yield, First lactation milk yield, Frieswal cattle, Peak yield

In India, crossbreeding has been practiced since 1875 by breeding the indigenous females with the temperate dairy breeds for increasing the milk production. Frieswal (Holstein Friesian × Sahiwal) is one of the crossbred dairy cattle being evolved by the ICAR- Central Institute for Research on Cattle (Formerly Project Directorate on Cattle), Meerut, India in collaboration with Ministry of Defence for higher milk production. The crossbred cattle maintained at military farms located in various agro-climatic regions of the country having 62.50±5.00% of exotic blood level is known as Frieswal cattle. The genetic evaluation of Frieswal cattle is mainly done on the basis of their milk production and superior animals are selected for improving the production. The peak yield (PY) and days to attain peak yield (DAPY) are two economically important traits which provide an early proof on the true milk production potential of dairy animals as their genetic correlation with the lactational milk yield is always found to be high. The PY and DAPY also play an important role in determining the persistency of milk production and the shape of lactation curve. Early selection of animals for milk production on the basis of these traits would also reduce the generation interval thereby increase the genetic gain per unit of time. Perusal of available literature revealed no scientific information on the PY and DAPY in Frieswal cattle. In view of the above facts, the present study was undertaken to estimate the effect of certain factors like farm, period, season and sire on the PY and DAPY in Frieswal cattle.

The data for the study was collected from two different military dairy farms viz., Meerut and Ambala. A total of 9957 daily milk yield records of 1073 Frieswal cows calved during 2005 to 2014 were collected and utilized for the study. Additional information on the date of birth, sire, dam, date of calving, date of drying etc. were also collected from the history cum pedigree sheets of the animals. The information on peak yield and days to attain peak yield were generated from the daily milk yield records. In order to

study the effect of different factors, the recorded data were classified according to the farm, sire, season and period of calving. The data included the information on cows maintained at Meerut and Ambala farms sired by 90 bulls. On the basis of temperature, humidity and rainfall, the year of calving was divided into three seasons viz., winter (Nov-Feb), summer (Mar-Jun) and rainy (Jul-Oct). The 10 years of study period was divided into five groups of two consecutive years viz., 2005–06, 2007–08, 2009–10, 2011–12 and 2013–14. The influence of various factors on the PY and DAPY was estimated by least squares analysis of variance for unequal non-orthogonal data using the GLM model of SPSS (Version 20). The statistical model used is as follows:

$$Y_{ijklm} = \mu + F_i + P_j + S_k + B_l + e_{ijklm}$$

where, Y_{ijklm} , dependent variables i.e. PY or DAPY; μ , overall mean; F_i , effect of i^{th} farm; P_j , effect of j^{th} period of calving; S_k , effect of k^{th} season of calving; B_l , random effect of l^{th} sire; e_{ijklm} , random error, assumed to be normally and independently distributed with mean zero and constant variance i.e. NID (0, σ_e^2).

Duncan's multiple range test as modified by Kramer (1957) was used for testing the statistical significance of differences among least squares means (using the inverse coefficient matrix).

The results of least squares analysis of variance on the effect of various factors on PY and DAPY are presented in Table 1. The factors included in the models for PY and DAPY accounted for 10.30 and 12.00% of the total variations, respectively.

Peak yield is defined as the maximum total yield obtained at any day of the lactation. The least squares analysis of variance revealed that the period of calving had highly significant ($P < 0.01$) effect on the first lactation peak yield while the differences due to other factors viz., farm, sire and season of calving were statistically non-significant. The variation in PY due to farm was the least significant followed by the sire and season of calving (Table 1). The significant effect of period of calving on PY was also reported by Subha Lakshmi *et al.* (2009) in Holstein Friesian × Sahiwal cattle, Patond *et al.* (2013) in Jersey cattle.

Present address:^{1,2}Senior Scientist (AG&B) (venkatesanraja09@gmail.com, ravinder774@gmail.com), ³Scientist (AG&B) (skratheecirc@gmail.com), ⁴Director (birhamp@gmail.com), ⁵Principal Scientist (AG&B) (usinghas@gmail.com).

Table 1. Least squares analysis of variance (mean squares only) for peak yield (kg) and days to attain peak yield in Frieswal cattle

Trait	Farm (1)	Period (4)	Season (2)	Sire	Error	R ² value (%)
Peak yield	0.008	61.554**	32.523	12.538 (90)	13.936 (973)	10.30
Days to attain peak yield	2325.984	709.235	8128.314**	841.384 (82)	905.960 (799)	12.00

Figures in parentheses indicate the degrees of freedom; **Significant at 1% level (P≤0.01)

Contrary to the present findings, Subha Lakshmi *et al.* (2009) in Holstein Friesian × Sahiwal cattle and Taher (2012) in Friesian cattle reported significant effect of season of calving on peak yield. However, Rao and Sundaresan (1982) in HF × Sahiwal cattle and Patond *et al.* (2013) in Jersey cattle reported non-significant effect of season of calving on peak yield which corroborated the results obtained in the present study. The non-significant effect of season of calving on peak yield suggested that the Frieswal cows express their full genetic potential for daily yield during the lactation irrespective of the season of calving. The study also revealed that the peak yield of animals maintained in two different farms viz., Ambala and Meerut were almost similar. The effect of sire on peak yield was found to be non-significant indicating that all the sires had similar genetic potential for peak yield; in other words, all the sires had similar breeding values for peak yield.

The overall least squares mean for PY in Frieswal cattle was estimated as 13.86±0.27 kg which was slightly higher than the average PY estimate of 13.30 kg reported by Subha Lakshmi *et al.* (2009) in a limited number of Holstein Friesian × Sahiwal crossbred cattle. The average PY of Frieswal cows was also superior to the estimates of 13.10 and 13.36 kg reported by Patond *et al.* (2013) and Ramchandraiah *et al.* (1990) in Jersey cattle, respectively. Roy and Katpatal (1998) reported even lower estimate of 11.88 kg in Jersey cattle while Taher (2012) reported a slightly higher average PY of 14.25 kg in Friesian breed of cattle. The average peak yields of Frieswal cows maintained at Meerut (13.85 kg) and Ambala (13.86 kg) farms were nearly same and comparable with that of pure Friesian as reported by Taher (2012). In general, the animals calved during winter season had the highest peak yield of 14.14 kg, while the rainy calvers had the lowest yield of 13.46 kg; however, the differences due to season were not statistically significant. The least squares averages of peak yield during different periods showed that the animals calved during 2009–10 had the highest peak yield of 14.87 kg, while the animals calved during 2007–08 had the lowest peak yield of 12.94 kg.

The overall average days to attain peak yield was estimated as 46.60±2.29 days (Table 2). Similar to the present findings, Taher (2012) reported average DAPY estimate of 47.53 days in Friesian cattle. Contrary to the present findings, Patond *et al.* (2013) reported a lower estimate of 21.41 days in Jersey cattle. The least squares analysis of variance revealed that only the season of calving had highly significant (P<0.01) effect on DAPY, while

period of calving had the lowest non-significant effect followed by sire and farm. The significant effect of season of calving on DAPY was also reported by Singh *et al.* (1989) in Haryana halfbreds with HF, Jersey and BS cattle; Taher (2012) in Friesian cattle and Patond *et al.* (2013) in Jersey cattle. The average DAPY of rainy (52.40 days) calvers were highest followed by winter (47.18 days) and summer (40.24 days) calvers. These results suggested that the animals calved during rainy season had taken more time to express their full genetic potential for milk production than the summer and winter calvers. This might be due to the fact that during the rainy season (July to October), the Frieswal cows were under thermal stress due to increased ambient temperature and humidity and the cows calved during rainy season might have diverted their energy to combat the thermal stress and so could not produce the maximum daily yield but once they enter the winter season (November-February), they express their full genetic potential for milk yield as the environment is favourable. However, by the time the rainy calvers enter into the winter season, their peak stage of production might have ceased due to the normal physiological process of lactation and so could not produce the peak yield as higher as the winter calvers. On the contrary, the summer calvers reached their peak yield at the earliest of 40.24 days and this result was

Table 2. Least squares means for peak yield (kg) and days to attain peak yield in Frieswal cattle

	Peak yield		Days to attain peak yield	
	No.	Mean ± SE	No.	Mean ± SE
Overall	1071	13.86±0.27	889	46.60±2.29
<i>Farm</i>				
Ambala	300	13.86±0.37	231	42.96±3.32
Meerut	771	13.85±0.38	658	50.25±3.13
<i>Period of calving</i>				
2005-06	42	13.81±0.73 ^{bc}	37	49.66±6.37
2007-08	185	12.94±0.46 ^a	156	44.90±3.94
2009-10	299	14.87±0.34 ^c	229	49.12±2.97
2011-12	267	14.10±0.39 ^{bc}	234	46.81±3.37
2013-14	278	13.56±0.47 ^{ab}	233	42.54±3.97
<i>Season of calving</i>				
Winter (Nov-Feb)	470	14.14±0.30	405	47.18±2.51 ^b
Summer (Mar-Jun)	378	13.97±0.31	307	40.24±2.66 ^a
Rainy (Jul – Oct)	223	13.46±0.36	177	52.40±3.16 ^c

Means bearing same superscripts did not differ significantly

as expected because, the animals calved during summer next entered in to the rainy season during which the animals were exposed to extreme climatic conditions which reduced the production potential of the animals. So, the summer calvers reached their peak during the summer season itself which resulted in the lowest DAPY as compared to the rainy calvers. The winter calvers produced the highest peak yield and had medium DAPY indicating that they are in their comfortable zone during the early and mid-stage of lactation i.e. winter and summer seasons.

The period of calving had non-significant effect on DAPY in Frieswal cattle. Similar to the present findings, Patil (1989) in HF × Gir cattle, Patond *et al.* (2013) in Jersey cattle also reported non-significant effect of period of calving on DAPY. Perusal of Table 2 revealed that the DAPY had decreased over the years from 49.66 days during 2005–06 to 42.54 days during 2013–14. The effect of sire and farm on the DAPY was nonsignificant.

Based on the results obtained in the present study, it may be concluded that the period of calving and season of calving significantly ($P < 0.01$) altered the PY and DAPY, respectively in Frieswal cattle. However, the farm and sire effects were statistically nonsignificant in altering the two traits in Frieswal cattle. The winter calvers had the lowest PY and longest DAPY while the summer calvers had the shortest DAPY and winter calvers had the highest PY in Frieswal cattle. The study also revealed that the thermal stress due to increased ambient temperature and humidity during the rainy season severely affected milk yield and the Frieswal animals were comfortable during the winter season for higher milk production.

SUMMARY

The effect of certain factors, viz. farm, sire, season and period of calving on first lactation peak yield (PY) and days to attain peak yield (DAPY) was assessed in Frieswal cattle. A total of 9957 daily milk yield records of 1073 Frieswal cows sired by 90 bulls maintained at Meerut and Ambala military farms and calved during the period from 2005 to 2014 were utilized for the study. The data were analysed using the least squares analysis of variance for unequal non-orthogonal data using the GLM of SPSS (Version 20). The overall least squares averages for PY and DAPY were estimated as 13.86 ± 0.27 kg and 46.60 days, respectively in Frieswal cattle and the factors included in the models accounted for 10.30 and 12.00% of the total variations, respectively. The least squares analysis revealed highly

significant effect of period of calving on PY and season of calving on DAPY. The rainy calvers had the lowest PY and longest DAPY while the summer calvers had the shortest DAPY and winter calvers had the highest PY in Frieswal cattle. The study also revealed that the thermal stress due to increased ambient temperature and humidity during the rainy season severely affected milk yield and the Frieswal animals were comfortable during the winter season and produced higher milk yield.

ACKNOWLEDGEMENTS

The authors express their sincere thanks to the DDG, Military farms; Director, Frieswal Project and the Officers in-charge of Ambala and Meerut farms for granting permission to collect data for the present study. The authors are also thankful to the Director, ICAR-CIRC for providing necessary facilities to conduct the present study.

REFERENCES

- Kramer C Y. 1957. Extension of multiple range test to group correlated adjusted means. *Biometrics* **13**: 13–17.
- Patil V S. 1989. Studies on peak yield and total milk yield in Friesian × Gir. M.Sc. (Ag.). Thesis, Mahatma Phule Krishi Vidyapeeth, Rahuri, Ahmednagar, M.S. (India).
- Patond M N, Khutal BB, Pachpute S T and Ramod S S. 2013. Studies on peak yield and days to attain peak yield in jersey cattle. *Research Journal of Animal Husbandry and Dairy Science* **4**(1): 4–6.
- Ramachandraiah K, Kumar K S and Srocmanoaryana O. 1990. A study of lactation persistency in relation to certain economic traits in purebred Jersey cows. *Indian Journal of Dairy Science* **43**: 270–73.
- Rao M K and Sundaresan D. 1982. Factors affecting shape of lactation curve in Friesian × Sahiwal crossbred cows. *Indian Journal of Dairy Science* **85**(2): 160–67.
- Roy N C and Katpatal B G. 1988. Genetic studies on first lactation peak yield and days to attain peak yield in Jersey cattle. *Indian Veterinary Journal* **65**(12): 1103–08.
- Singh V K, Nivsarkar A E, Nantyal A P and Singh B P. 1989. Studies on peak yield and days to attain peak yield in crossbred cattle. *Indian Journal of Animal Production and Management* **2**: 56–60.
- Shubha Lakshmi B, Gupta R, Sudhakar K, Gnana Prakash M and Sharma S. 2009. Genetic analysis of production performance of Holstein Friesian × Sahiwal cows. *Tamil Nadu Journal of Veterinary and Animal Sciences* **5**(4): 143–48.
- Taher K N. 2012. Some factors influencing peak yield and days attain to peak yield in Friesian cattle in the central region of Iraq. *AL-Qadisiya Journal of Veterinary Medicine Science* **11**(2): 117–20.