

Biology of the cleftbelly trevally *Atropus atropus* (Bloch and Schneider, 1801) from Ratnagiri coast of India

Akanksha V. Rakhunde¹, Vivek H. Nirmale^{1*}, Ravindra A. Pawar¹, Santosh Y. Metar² and Bhaskar P. Bhosale¹

¹College of Fisheries, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Ratnagiri-415 629, Maharashtra, India

²Marine Biological Research Station, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Zadgaon, Ratnagiri-415 612, Maharashtra, India

Abstract

The cleftbelly trevally *Atropus atropus* (Bloch and Schneider, 1801) is one of the important carangid species landed mostly as trawl bycatch along Ratnagiri coast. Investigations were carried out on *A. atropus* with respect to length-weight relationship, condition factor, morphometrics, feeding and reproductive biology. The length-weight relationship indicated negative allometric growth in *A. atropus*. Morphometric measurements showed significant correlation with total length. The gut content analysis revealed *A. atropus* to be a carnivore fish feeding mainly on *Acetes* spp., small teleostean fishes, cephalopods, and squilla. Sex ratio showed dominance of males. Gonadosomatic index and maturity studies indicated that *A. atropus* has a protracted spawning season extending from May to December. Length at sexual maturity for females was estimated to be 171 mm. Fecundity ranged from 16458 to 138688 eggs with an average of 70097 eggs. Ova diameter studies showed that *A. atropus* spawns twice in a protracted spawning season.

Introduction

Family Carangidae is a highly diverse group of important food fishes including jacks, trevallies, scads, queenfishes, runners, pompanos and amberjacks. The family is made up of four sub-families represented by 32 genera and 140 species (Nelson, 2006). Carangids can be found in all tropical as well as subtropical marine seas worldwide, and occasionally in temperate regions (Nair *et al.*, 2018). They are extensively spread in the Atlantic, Pacific, and Indian oceans (Nelson, 2006). Carangids occur along both the east and west coasts of India. They are mostly confined in their distribution to shallow waters upto a depth of 60 m (Nair, 2000). The carangid catches are dominant during the monsoon season, especially along the north-east and south-west coasts of India (Sivakami *et al.*, 1996). About 62 species from 20 genera have been reported from Indian waters (Joshi *et al.*, 2011). Carangids are mostly

caught by trawls, purse-seines and drift gillnets apart from boat seines, gillnets and hooks and line (Bandkar *et al.*, 2020,2022). In India, the total production of carangids stood at 1.8 lakh t contributing about 4.83% to the total marine fish landings of the country (Anon., 2020). The cleft belly trevally, *Atropus Atropus* (Bloch and Schneider, 1801) is an important species belonging to family Carangidae. *A. atropus* is found in tropical and subtropical waters of the Indo-Pacific, from the south-western to the southern regions of Taiwan (Lin and Shao, 1999). In India, *A. atropus* is distributed along the entire coastline, however it is abundant in the states of Gujarat, Maharashtra, Karnataka, Kerala and Tamil Nadu. Studies on length-weight relationship of *A. atropus* were carried out by Reuben *et al.* (1992), Raje (1994), Kasim (1999), Qamar and Panhwar (2017), Rajesh *et al.* (2017b) and Rahman *et al.* (2021); fishery and biology was studied from Gujarat coast by Raje (1994); maturation and



*Correspondence e-mail:

viveknirmale416@gmail.com

Keywords:

Bycatch, Carangidae, Carnivore, Reproductive biology, Spawning season

Received : 14.12.2022

Accepted : 01.06.2023

spawning studies by Rajesh et al. (2018); biology and stock assessment by Reuben et al. (1992); Kasim (2003) and food and feeding habits by Rajesh et al. (2017a). Reports on the biology of *Atropus atropus* from Indian waters are very few and no work has been carried out on the species from Ratnagiri coast of India. Hence this study on the length-weight relationships, morphometrics, reproductive biology and trophodynamics of *A. atropus* from Ratnagiri was undertaken. The biological studies on the species will yield vital information on growth, feeding habits, reproductive potential, maturity and spawning, which can form useful inputs for promulgating management plans for sustainable fisheries of the species from the region.

Materials and methods

A total of 589 samples of *Atropus atropus* were collected from commercial trawl bycatch landed at Mirkarwada fish landing centre (16.9800° N; 73.3000° E) in Ratnagiri, Maharashtra, at weekly intervals from December 2021 to October 2022. Species identification was carried out as per Fischer and Bianchi (1984). Total length (cm) and body weight (g) for each individual were recorded. Nineteen morphometric measures were regressed on the total length (Zar, 2005). The length-weight relationship (LWR) was calculated by linear regression on the transformed equation: $\log W = \log a + b \log L$, where W is body weight (g), L is length (cm), a is intercept and b is slope of linear relationship analysis (Froese, 2006). t-test was employed to check whether estimated 'b' values differed significantly from 3, which would indicate the type of growth to be isometric, negative or positive allometric. Relative condition factor (Kn) was calculated as per Le Cren (1951):

$$Kn = \frac{W}{Wc}$$

where, Kn is relative condition factor, W is observed weight of the fish and Wc is estimated weight of the fish.

The fishes were dissected and the sex and the stage of maturity were noted. Data on sex ratio were analysed by χ^2 (chi square) test to find out if either sex was dominant. Gonads were excised out and weighed using analytical balance to the nearest 0.001g, and preserved in 4% neutral formalin for further studies. The gut content analysis was carried out as per Hynes (1950) using Frequency of Occurrence and Points (volumetric) methods. Gonadosomatic index (GSI) was estimated as per Bal and Rao (1984). To define the different

maturity stages, colour and size of the ovary and size of the ova were taken into account. The female maturity stages of *A. atropus* were classified using a five-point maturity scale used in the case of multiple spawners (Holden and Raitt, 1974) as immature (I), maturing (II), matured (III), gravid (IV) and spent (V) based on the observations made at the laboratory on physical appearance such as colour and abdominal cavity occupied by gonads as well as ova diameter. The mean length at sexual maturity (L_m) was estimated as per King (1995). Fecundity was determined as per Sinha (1995). Ova diameter of intra ovarian eggs was measured from subsamples taken from anterior, middle and posterior regions of the ovary using ocular micrometer. Frequency polygons were drawn for all stages of maturity for ascertaining spawning frequency.

Results and discussion

Length-weight relationship and morphometrics

Estimated parameters of the length-weight relationship including total length (TL) and total weight (TW), number of specimens (n), values of parameters 'a' and 'b' and the coefficient of determination (r^2) are given in Table 1. The generalised length-weight relationship of *A. atropus* along the Ratnagiri coast was determined as $W = 0.0358 L^{2.7234}$. The estimated 'b' values in the present study indicated negative allometric growth ($p < 0.05$) which is similar to the findings of Reuben et al. (1992), Raje (1994), Kasim (1999), Qamar and Panhwar (2017) and Rajesh et al. (2017b). Rahman et al. (2021) reported that *A. atropus* from the Bay of Bengal, off Bangladesh exhibited positive allometric growth based on estimated b values. The 'b' values in the present study are found to be in the expected range of 2.5 to 3.5 (Froese, 2006). Variability in the length-weight relationship can be attributed to several factors such as sex, gonad maturity, stomach fullness, sample size, size range cover and habitat (Krishnamoorthi, 1971; Froese, 2006; Bandkar et al., 2022).

Comparison of the nineteen morphometric measurements with the total length (TL) of the species yielded a moderate to high degree of correlation ranging from 0.7131 to 0.997 (Table 2). The morphometric characters compared in the present study would prove helpful for comparison with *A. atropus* in other geographical areas. Morphological variability among different geographical populations could

Table 1. Estimated length-weight relationship for *A. atropus* off Ratnagiri coast

Group	N	TL range (cm)	Body weight range (g)	W= aL ^b parameters		
				a	b	r ²
Males	300	11.2-25.4	23.9-222.5	0.0361	2.7186	0.9816
Females	247	11.4-25	26.1-221.6	0.0309	2.7770	0.9862
Indeterminates	42	10-12.1	20.9-35.2	0.0460	2.6318	0.6121
Total (combined)	589	10-25.4	20.9-222.5	0.0358	2.7234	0.9848

N-Number of specimens; a-Intercept; b-Slope of linear relationship; r²-Coefficient of determination

be attributed to the different genetic structure of populations and to different environmental conditions prevailing in each geographical area (Mamuris *et al.*, 1998). Therefore, the animals, with the same morphometric measurements are often assumed to constitute a stock (Waldman *et al.*, 1988) and has been used widely in fishery stock differentiation studies (Avsar, 1994; Natarajan *et al.*, 2011).

Condition factor

Condition factor was observed to be minimum in February and maximum in August (Fig. 1). The values of condition factor were higher during the spawning season. Raje (1994) observed that the variation in Kn values in females

Table 2. Comparison between various morphometric characters of *A. atropus*

Regressions tested	Observations (n)	Coefficient of Correlation (r)
TL vs Standard length	589	0.9925
TL vs Fork length	589	0.997
TL vs Head length	589	0.9717
TL vs Snout length	589	0.8746
TL vs Eye diameter	589	0.9492
TL vs First dorsal base length	589	0.9078
TL vs Second dorsal base length	589	0.7131
TL vs Anal base length	589	0.9849
TL vs First pre-dorsal length	589	0.8833
TL vs Second pre-dorsal length	589	0.9787
TL vs Pre-anal length	589	0.9822
TL vs Pre-pelvic length	589	0.9423
TL vs Pre-pectoral length	589	0.9747
TL vs Body depth	589	0.9867
TL vs Pectoral fin length	589	0.9682
TL vs First dorsal height	589	0.8124
TL vs Second dorsal height	589	0.9013
TL vs Anal height	589	0.9211
TL vs Pelvic height	589	0.9467

of *A. atropus* was more strongly associated with gonad maturity. Rajesh *et al.* (2017b) observed monthly variation in condition factor (Kn) of *A. atropus* from Mangalore coast and reported that Kn values were roughly comparable in both sexes. Alteration in the general condition of the fish can be due to changes in the gonad cycle, including the state of fullness of the alimentary canal (Weatherly, 1979). The




Fig. 1. Month-wise relative condition factor in *A. atropus*

Kn value is influenced by various factors like environment, feeding intensity, food type, amount of fat or muscular developments, breeding cycle, age, sex and development stage of gonads of fish (Abowei, 2010; Pawase *et al.*, 2020).

Food and feeding biology

The results of gut content analysis by Frequency of Occurrence and Points (volumetric) method is given in Fig. 2 and 3 respectively. The Index of Preponderance is presented in Table 3. Table 4 gives a comparative account on the feeding habits of *A. atropus* and other species of trevallies reported by different authors. The carnivorous feeding habit of *A. atropus* from Ratnagiri coast is in conformation with the earlier reports of Fischer and Bianchi (1984), Raje




Fig. 2. Percentage composition of food items of *A. atropus* during different months by Frequency of Occurrence method




Fig. 3. Percentage composition of food items of *A. atropus* during different months by Points (volumetric) method

(1994) and Rajesh et al. (2017a). The variations in food items consumed by a fish during different months may be due to the dominance of respective food items in different regions (Bandkar et al., 2022). Smaller and moderate size fishes of *A. atropos* from Ratnagiri in the present study had *Acetes* spp., semi-digested matter, fish scales and cephalopods in their stomach, whereas the bigger individuals mainly had small teleostean fishes and squilla. Carangids feed on planktonic crustaceans in the early stages of their life and on fish juveniles in the later stages (Sivakami, 1996). Food preference observed in individuals of different size groups can be attributed to the size and availability of prey organisms, mouth size and life history traits (Garcia et al., 2018; Batool and Siddiqui, 2020).

The feeding intensity in *A. atropos* was generally observed to be higher in maturing and spent individuals and lower in spawners in the present study. Sivakami (1996) noted that the feeding intensity in carangids was more in immature and juvenile fish and less in breeders. Venkataramani and Natarajan (1988) observed that in *Seriolina leptolepis*, maturing and spent recovering fishes fed more than juveniles and matured groups. The highest percentage of full stomachs in *A. atropos* was observed during post-spawning months and empty stomachs were noted during spawning season.

However, individuals with full stomachs were also noted in August which possibly included spent and immature individuals. Similarly, low feeding intensity was also observed in maturing pre-spawners during April, contradicting the general observation that increased feeding takes place in maturing individuals. This may be related to the sparse availability of prey food in the environment. A similar condition with respect to spawning and feeding intensity was observed by Kagwade (1968) in *Caranx kalla* from Calicut coast and Kende (2016) in *Thryssa mystax* from Ratnagiri coast. Raje (1994) reported that the low feeding intensity was associated with the spawning period only during certain months in which females were both in maturing and matured condition.

Reproductive biology

In the present study, the average sex ratio was found to be 1:0.82. However, significant difference ($p < 0.05$) was noted in the sex ratio during December and October only (Fig. 4). The results confirmed the findings of Raje (1994). Rajesh et al. (2018) reported the dominance of females from Mangalore coast in *A. atropos*. The change in the sex ratio from the Mendelian ratio of 1:1 may be caused by partial segregation of mature forms through habitat preferences (Reynolds,

Table 3. Index of Preponderance (IP) of food items in the stomach of *A. atropos*

Food items	Percentage of occurrence (Oi)	Percentage of volume (Vi)	Oi Vi	IP	Grade
<i>Acetes</i> spp.	42.58	30.28	1289.36	33.99	I
Small teleostean fishes	36.62	28.91	1058.65	27.91	II
Semi-digested matter	48.76	21.13	1030.05	27.15	III
Fish scales	25.61	15.58	399.075	10.52	IV
Cephalopod	4.73	3.22	15.2246	0.40	V
Squilla	1.26	0.88	1.11747	0.03	VI

Table 4. Food composition of *A. atropos* and other species of trevallies from different localities

Species	Locality	Main food item	Feeding habit	Author
<i>A. atropos</i>	Western Indian Ocean	Shrimps, copepods, decapod crustaceans and small fish	Carnivorous	Fischer and Bianchi (1984)
<i>A. atropos</i>	Veraval, Gujarat	<i>Acetes</i> spp., cephalopods, teleosts, squilla and copepods	Carnivorous	Raje (1994)
<i>Caranx ignobilis</i>	Tamil Nadu	Juveniles of sardines, anchovies, and other finfishes, shrimps, crabs instars, amphipods, <i>Decapterus</i> sp., other carangids, threadfin breams, lizard fishes, silverbellies, goatfishes and crabs	Carnivorous	Abdussamad et al. (2008)
<i>A. atropos</i>	Mangalore, Karnataka	<i>Acetes</i> spp., semi-digested matter, fish juveniles, teleosts, cephalopods, and copepods	Carnivorous	Rajesh et al. (2017a)
<i>Seriolina nigrofasciata</i>	South-west coast	Teleosts, cephalopods and crustaceans	Carnivorous	Rajesh et al. (2019)
<i>A. atropos</i>	Ratnagiri	<i>Acetes</i> spp., small teleostean fishes, cephalopods, fish scales, squilla and semi-digested matter	Carnivorous	Present study

1974), due to migration (Collignon, 1960) or behavioural differences between sexes (Polonsky and Tormosova, 1969) thus rendering one sex to be more easily caught than another. The dominance of either sex might be possible due to selectivity of gear, vulnerability of either sex to capture and variation in size at first capture (Pawase *et al.*, 2020).

GSI was noted to be higher during the period May to December indicating the occurrence of more ripe individuals (Fig. 5). Individuals with ovaries in Stage III (Matured) and Stage IV (Gravid) were observed to be maximum during this period (Fig. 6). It can be inferred from the maturity studies and GSI studies that *A. atropus* has a prolonged spawning season extending from May to December along Ratnagiri coast. Reuben *et al.* (1992) observed the spawning period of *A. atropus* from north-west coast to be around November or December. Raje (1994) noted that *A. atropus* from Veraval coast had a prolonged spawning period and spawning took place twice in the spawning season. The spawning season of *A. atropus* from the Gujarat coast was reported to be from May to September by Kasim (1999). Spawning of *A. atropus* along the north-west coast took place in November and December (Kasim, 2003). Based on the GSI and maturity studies on *A. atropus* from Mangalore coast, Rajesh *et al.*




Fig. 4. Monthly variation in numbers of male and female fish in *A. Atropus* samples collected during the study




Fig. 5. Monthly GSI trend in males and females of *A. atropus*




Fig. 6. Monthly variation in maturity stages (%) in females of *A. atropus*

(2018) observed that *A. atropus* is a prolonged spawner spawning twice in a season from September to December and February to May. The variation in the spawning season of a fish could be attributed to spatial, environmental and stock level differences (Bandkar *et al.*, 2022).

A. atropus is found to be a moderately fecund fish with an average fecundity of 70097 eggs per individual. Raje (1994) reported that the fecundity of *A. atropus* from the Veraval coast ranged between 31432 to 372344 eggs per fish, with an average of 141515 eggs in sizes ranging from 165-253 mm. It was observed that the fecundity increased with an increase in the size of the fish. Similar trend in fecundity was reported by Kagwade (1968) for *Caranx kalla*. Rajesh *et al.* (2018) reported that the fecundity of *A. atropus* from Mangalore coast ranged from 33298 to 188675 eggs with an average of 94083 eggs. They observed the variation in fecundity with respect to both length and weight. The fecundity varies from species to species and within a species from one region to another in accordance with the reproductive potential of the stocks (Kende, 2016; Gurjar *et al.*, 2018; Pawase *et al.*, 2020; Bandkar *et al.*, 2022).

The appearance of two modes in Stage IV ovaries (Fig. 7) shows the possibility of *A. atropus* spawning two times in a spawning season along the Ratnagiri coast. Rajesh *et al.* (2018) reported the presence of two modes of mature ova in Stage VI of *A. atropus* from Mangalore coast and stated that *A. atropus* spawned twice in a spawning season.

The size at first maturity for females of *A. atropus* from Ratnagiri coast was estimated to be 17.1 cm (Fig. 8). Length at first maturity reported by different workers (Reuben *et al.*, 1992; Raje, 1994; Rajesh *et al.*, 2018) for *A. atropus* ranged from 16 to 21 cm. Difference in size at first maturity within a species points possibly to the presence of distinct stock, variation in growth rate and maximum size reached by different species under the influence of environmental factors including availability of food (Sawant *et al.*, 2017; Mahadwala, 2019; Pawase *et al.*, 2020; Bandkar *et al.*, 2022).

The present study provides vital information on morphometrics and length-weight relationship, feeding and reproductive biology of *A. atropus* from Ratnagiri coast. *A. atropus* was




Fig. 7. Ova diameter trend in *A. atropos*. (a) Stage I, (b) Stage II, (c) Stage III, (d) Stage IV, (e) Stage V




Fig. 8. Mean length at sexual maturity in females of *A. atropos* along Ratnagiri coast

found to be moderately fecund fish with a spawning season from May to December in the study region. The size at first maturity for females was estimated to be 17.1 cm. The morphometric results of this study can be used for stock identification of the species in Indian waters. This study provides an insight about the biology of *A. atropos* that can be effectively used to frame management policies for sustaining the fisheries of this species in the region.

Acknowledgements

The authors are thankful to the authorities of Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli for providing the facilities at College of Fisheries, Ratnagiri during the entire period of research work.

References

- Abdussamad, E. M., Kasim, H. M. and Balasubramanian, T. S. 2008. Distribution, biology and behaviour of the giant trevally, *Caranx ignobilis*-A candidate species for mariculture. *Bangladesh J. Fish. Res.*, 12(1): 89-94.
- Abowei, J. F. N. 2010. The condition factor, length-weight relationship and abundance of *Ilisha africana* (Block, 1795) from Nkoro River Niger Delta, Nigeria. *Adv. J. Food. Sci. Technol.*, 2(1): 6-11.
- Anon. 2020. *Handbook of fisheries statistics*. Department of Fisheries, Ministry of Fisheries, Animal Husbandry and Dairying, Government of India, New Delhi, India.
- Avsar, D. 1994. A stock differentiation study of the sprat (*Sprattus sprattus phalericus* Risso) off the southern coast of the Black Sea. *Fish. Res.*, 19(3-4): 363-378. [https://doi.org/10.1016/0165-7836\(94\)90050-7](https://doi.org/10.1016/0165-7836(94)90050-7)
- Bal, D. V. and Rao, K. V. 1984. *Marine fisheries of India*. Tata McGraw-Hill Publishing Co. Ltd., New Delhi, India, 470 pp.
- Bandkar, D. S., Nirmale, V. H., Metar S. Y. and Pawar R. A. 2020. Estimation of population parameters of shrimp scad, *Alepes djedaba* (Forsskal, 1775) along the Ratnagiri coast of Maharashtra, India. *J. Indian Fish. Assoc.*, 46(1): 67-73.
- Bandkar, D., Nirmale, V. H., Gurjar, U. R., Metar, S. Y., Pawar, R. A. and Chogale, N. D. 2022. Reproductive biology and feeding dynamics of the shrimp scad *Alepes djedaba* (Forsskal, 1775) from Ratnagiri, central west coast of India. *Indian J. Fish.*, 69(1): 37-44. <https://doi.org/10.21077/ijf.2022.69.1.109595-04>
- Batool, T. and Siddiqui, G. 2020. The seasonal variation in diet composition and feeding strategy of *Grammoplites suppositus* (Troschel, 1840) (Family: Platycephalidae) in relation to its sex and size. *Indian J. Mar. Sci.*, 49(12): 1848-1855.
- Collignon, J. 1960. Contribution to the knowledge of the *Otolithus* of the coasts of Equatorial Africa. *Bulletin of the Institute of Central African Studies*, 19-20: 55-84 (In French).
- Fischer, W. and Bianchi, G. 1984. *FAO species identification sheets for fishery purposes. Western Indian Ocean; Fishing Area 51*. Food and Agricultural Organisation of the United Nations, Rome, Italy.
- Froese, R. 2006. Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *J. Appl. Ichthyol.*, 22(4): 241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>
- Garcia, D. A. Z., Vidotto-Magnoni, A. P. and Orsi, M. L. 2018. Diet and feeding ecology of non-native fishes in lentic and lotic freshwater habitats. *Aquat. Invasions*, 13: 565-573.
- Gurjar, U. R., Sawant, M. S., Pawar, R. A., Nirmale, V. H. and Pawase, A. S. 2018. Reproductive biology and fishery of the white sardine, *Escualosa thoracata* (Valenciennes, 1847) from the Ratnagiri coast, Maharashtra. *Indian J. Mar. Sci.*, 47(12): 2485-2491.
- Holden, M. J. and Raitt, D. F. S. 1974. Manual of fisheries science, Part 2: Methods of resource investigation and their application. *FAO Fisheries Technical Paper No. 115. Rev. 1*. Food and Agricultural Organisation of the United Nations, Rome, Italy.
- Hynes, H. B. N. 1950. The food of freshwater sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*), with a review of methods used in studies of the food of fishes. *J. Anim. Ecol.*, 19(1): 36-58. <https://doi.org/10.2307/1570>
- Joshi, K. K., Nair, R. J., Thomas, E. M. A., Kakati, V. S., Varghese, S. J. M., Paul, M., Gorge, S. S. and Manisseri, M. K. 2011. *The Carangids of India-A Monograph*. ICAR-Central Marine Fisheries Research Institute, Kochi, India, 437 p.
- Kagwade, V. N. 1968. Maturation and spawning in horse mackerel, *Caranx kalla* (Cuv. & Val.). *Indian J. Fish.*, 15: 207-220.
- Kasim, M. H. 1999. Age growth mortality yield per recruit and stock assessment of *Atropus atropus* (Bloch and schneider). *The Fourth Indian Fisheries Forum Proceedings*, Asian Fisheries Society Indian Branch, Mangaluru, Karnataka, India, p. 373-375.
- Kasim, M. H. 2003. Carangids. In: Mohan Joseph, M. and Jayaprakash, A. (Eds.), *Status of exploited marine fishery resources of India*, ICAR-Central Marine Fisheries Research Institute, Kochi, India, pp.166-175.
- Kende, D. R. 2016. *Biological studies on moustached thryssa, Thryssa mystax along the Ratnagiri coast of Maharashtra*. M. F. Sc. Thesis, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Maharashtra, India, pp.39-44.
- King, M. 1995. *Fisheries biology, assessment and management*. Fishing news Books Ltd., Surrey, UK, pp. 158-160.
- Krishnamoorthi, B. 1971. Biology of the threadfin bream, *Nemipterus japonicus* (Block). *Indian J. Fish.*, 18(1&2): 1-21.
- Le Cren, E. D. 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20: 2-19. <http://dx.doi.org/10.2307/1540>
- Lin, P. L. and Shao, K. T. 1999. A review of the carangid fishes (Family Carangidae) from Taiwan with descriptions of four new records. *Zool. Stud.*, 38(1): 33-68.
- Mahadwala, P. L. 2019. *Biological studies on the spineless cuttlefish Sepiella inermis (Orbigny, 1848) along the Ratnagiri coast of Maharashtra*. M. F. Sc. Thesis, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Maharashtra, India, 40 p.
- Mamuris, Z., Apostolidis, P., Panagiotaki, P., Theodorou, A. J. and Triantaphyllidis, C. 1998. Morphological variation between red mullet populations in Greece. *J. Fish Biol.*, 52(1), 107-117. <https://doi.org/10.1111/j.1095-8649.1998.tb01556.x>
- Nair, P. 2000. Carangid resources of India. In: *Marine fisheries research and management*. ICAR-Central Marine Fisheries Research Institute, Kochi, India, pp. 317-348.
- Nair, R., Joshi, K. K., Kuriakose, S. and Geetha, P. M. 2018. *A study on the diversity of carangid resources off Kochi, Kerala*. <https://doi.org/10.13140/RG.2.2.14328.96003>
- Natarajan, S., Subrahmanyam, S., Santhanam, R. and Thangavel, B. 2011. Morphometric studies on wild caught and cultured shrimp *Penaeus monodon* (Fabricius, 1798) from Parangipettai, India. *Adv. Appl. Sci. Res.*, 2: 490-507.
- Nelson, J. S. 2006. *Fishes of the world*, 4th edn. John Wiley and Sons Inc., Hoboken, New Jersey, USA, 362 p.
- Pawase, S. V., Nirmale, V. H., Bhosale, B. P., Pawar, R. A., Sawant, M. S. and Kende, D. R. 2020. Study on biology of *Thryssa dussumieri* (Valenciennes, 1848) from the coast of Ratnagiri, Maharashtra, India. *Indian J. Mar. Sci.*, 49 (01): 87-94.
- Polonsky, A. S. and Tormosova, I. D. 1969. The spawning of the horse mackerel of the North-east Atlantic and the distribution of its eggs and larvae. *Trudy Atlant. NIRO*, 23: 27-48.
- Qamar, N. and Panhwar, S. K. 2017. Length-weight relationships of nine rarely occurring carangids in the northern Arabian Sea coast of Pakistan. *J. Appl. Ichthyol.*, 34(1), 221-223. <https://doi.org/10.1111/jai.13540>
- Rahman, M. A., Hossain, M. Y., Tanjin, S., Mawa, Z., Hasan, M. R., Habib, K. A. and Ohtomi, J. 2021. Length weight relationships of five marine fishes from the Bay of Bengal. *J. Appl. Ichthyol.*, 37(2): 364-366. <https://doi.org/10.1111/jai.14176>
- Raje, S. G. 1994. Fishery and biology of *Atropus atropus* (Bloch and Schneider) from Veraval. *J. Mar. Biol. Assoc. India*, 36(2): 181-187.
- Rajesh, D. P., Anjanayappa, H. N. and Benakappa, S. 2018. Maturation and spawning frequencies of *Atropus atropus* from Mangalore coast, Karnataka, India. *Journal of Fisheries and Life Science*, 3: 46-53.
- Rajesh, D. P., Anjanayappa, H. N., Benakappa, S., Naik, A. S. and Prakash, D. P. 2017a. Food and feeding habits of *Atropus atropus* from Mangalore coast. *J. Exp. Zool. India*, 20(1): 1309-1312.
- Rajesh, D. P., Anjanayappa, H. N., Nayana, P. and Benakappa, S. 2017b. Length-weight relationship and relative condition factor of *Atropus atropus* (Bloch & Schneider, 1801) from Mangalore coast. *J. Aquac. Res. Dev.*, 8: 2 (Suppl).
- Rajesh, K. M., Rohit, P., Mini, K. G., Sathyavathi, T. and Abdul Hakeem, M. M. 2019. *Seriolina nigrofasciata* (Ruppel, 1829), its fishery and biological aspects off south-west coast of India. *Indian J. Fish.*, 66(1): 9-16. <https://doi.org/10.21077/ijf.2019.66.1.82727-02>

- Reuben, S., Mohamad Kasim, H., Sivakami, S., Nair, P. N., Kurup, K. N., Sivadas, M., Noble, A., Nair, K. V. and Raje, S. G. 1992. Fishery, biology and stock assessment of carangid resources from the Indian seas. *Indian J. Fish.*, 39 (3&4): 195-234.
- Reynolds, J. D. 1974. Biology of the small pelagic fishes in the New Volta Lake in Ghana, Part III: Sex and reproduction. *Hydrobiologia*, 45(4): 489-508.
- Sawant, P. P., Nirmale, V. H., Metar, S. Y., Bhosale, B. P. and Chogale, N. D. 2017. Biology of Indian sand whiting, *Sillago sihama* (Forsskal) along the Ratnagiri coast. *Indian J. Mar. Sci.*, 46(9): 1899-1907.
- Sinha, R. K. 1995. *Some aspects of biology of freshwater catfish Clarius batrachus (Linn. 1758) of the Bombay region*. M. Sc. Thesis, ICAR-Central Institute of Fisheries Education, Mumbai, India, 74 p.
- Sivakami, S. 1996. On the food habits of the fishes of the family carangidae-A review. *J. Mar. Biol. Assoc. India*, 38(1 & 2): 118-123.
- Sivakami, S., Nair, P. N., Jayaprakash, A. A., Mohamad Kasim, H., Yohannan, T. M., Sivadas, M., Koya, K. P. and Seetha, P. K. 1996. Distribution and abundance of carangids along the EEZ India. In: *Proceedings of the Second Workshop on Scientific Results of FORV Sagar Sampada*, Department of Ocean Development, New Delhi, India, pp.347-361.
- Venkataramani, V. K. and Natarajan, R. 1988. Food and feeding habits of *Selaroides leptolepis* Val. Off Tuticorin coast. *Matsya*, 14: pp. 53-63.
- Waldman, J. R., Grossfield, J. and Wrigin, I. 1988. Review of stock discrimination techniques for striped bass. *N. Am. J. Fish. Manag.*, 8 (4): 410-425.
- Weatherly, A. H. 1979. *Growth and ecology of fish population*. Academic press Inc. Ltd., London, UK, 75 p.
- Zar, J. H. 2005. *Biostatistical analysis*, 4th edn. Pearson Education (Singapore) Pvt. Ltd., New Delhi, India, 633 p.