

## Evaluation of genetic variation in *Labeo fimbriatus* (Bloch, 1795) populations using heterologous primers

S. SWAIN<sup>1</sup>, S. P. DAS<sup>1</sup>, D. BEJ<sup>1</sup>, A. PATEL<sup>1</sup>, P. JAYASANKAR<sup>1</sup>,  
B. K. CHAUDHARY<sup>2</sup>, J. K. JENA<sup>3</sup>, P. ROUTRAY<sup>1</sup>, S. K. SWAIN<sup>1</sup>,  
P. C. DAS<sup>1</sup>, W. S. LAKRA<sup>4</sup> AND P. DAS<sup>1</sup>

<sup>1</sup>Fish Genetics and Biotechnology Division, Central Institute of Freshwater Aquaculture, Kausalyaganga  
Bhubaneswar - 751 002, Odisha, India

<sup>2</sup>ICAR Research Complex for Eastern Region, Patna - 801 505, Bihar, India

<sup>3</sup>National Bureau of Fish Genetic Resources, Canal Ring Road, P. O. Dilkusha, Lucknow - 226 002, Uttar Pradesh, India

<sup>4</sup>Central Institute of Fisheries Education, Mumbai - 400 061, Maharashtra, India

e-mail: pdas77@hotmail.com

### ABSTRACT

The fringe lipped peninsula carp, *Labeo fimbriatus*, is native to lower peninsular region of India. It is a commercially important species widely fished and consumed throughout much of the tropical and subtropical region of south India. Knowledge of genetic diversity of this species is important to support management and conservation programmes which will subsequently help in sustainable production of this species. DNA markers, mostly microsatellites are excellent tools for evaluation of genetic variation of populations. However, development of these markers from each species is time consuming and costly. Therefore, we studied the usefulness of heterologous markers by assessing the genetic variation of three populations of *L. fimbriatus*. We examined population structure of *L. fimbriatus* from different riverine locations in India using rohu microsatellite loci. Out of 30 rohu microsatellite loci, 15 produced amplified products having unambiguous band pattern with clarity in allele scoring. Therefore, these 15 loci were used in 110 samples of *L. fimbriatus* collected from river Krishna (Vijayawada; n= 30), river Kaveri (Bengaluru; n=50) and river Mahanadi (Sambalpur; n=30). Out of 15 loci, only 3 loci were polymorphic in all the populations and remaining 12 loci were found to be monomorphic. The range of expected heterozygosity values for these polymorphic loci was 0.597 to 0.602. One locus did not satisfy Hardy-Weinberg equilibrium ( $p=.004$ ) in Kaveri population. Negative inbreeding coefficients ( $F_{is}$ ) were found across populations with no significant population structuring ( $F_{st}$ ) indicating existence of panmictic populations. As only 3 loci (20%) out of 15 were polymorphic, use of heterologous primers may not be a choice for population studies in *L. fimbriatus*.

Keywords: Heterologous primers, *Labeo fimbriatus*, Microsatellite markers

### Introduction

Aquaculture is the fastest growing primary production sector in the world. Being home for more than 10% of global fish biodiversity, India ranks third in the world in total fish production (Ayyappan *et al.*, 2009). Indian aquaculture production comprises mainly three Indian major carps (*Labeo rohita*, *Catla catla* and *Cirrhinus mrigala*). Common carp and other medium and minor carps constitute about 5% of the total freshwater aquaculture production. With the increase in demand for aquaculture foods, there is need for more efficient production systems. Therefore, emphasis is being given to species diversification. Several species of *Labeo*, other than rohu such as *L. fimbriatus*,

*L. calbasu*, *L. bata* and *L. gonius* are being introduced in the culture system as potential candidate species (Jena *et al.*, 2011). *L. fimbriatus*, locally known as 'podosi' is gaining importance for commercial aquaculture. This species is widely distributed in South-East Asia including Indian River systems. Published work on *L. fimbriatus* include evaluation of growth in tanks provided with solid substrates (Mridula *et al.*, 2003), impact of endosulfan (Saravanan *et al.*, 2011) and on life cycle trait pattern like age, growth, maturity and fecundity (Rao, 2011). Limited information is available on the genetic aspect of this species, except for preliminary studies on karyotyping (Biswal *et al.*, 2010).

Knowledge of genetic parameters of the species cultured is vital for sustainability of the production system. More emphasis is being given on genetic improvement of aquacultured species, following the success story of salmon, tilapia and rohu selective breeding programmes (Gjedrem and Baranski, 2009). Owing to high market value, excellent flavor and meat quality, *L. fimbriatus* is being targeted for its genetic improvement in order to increase production efficiency which necessitates proper evaluation of genetic stock that will greatly help to create base population.

Microsatellite markers have been developed for selected Indian fish species such as rohu (Das *et al.*, 2005; Patel *et al.*, 2009), catla (McConnell *et al.*, 2001), chitala (Punia *et al.*, 2006) and mrigala (Lal *et al.*, 2011). Simple sequence repeat (SSR) markers are preferable because they are codominant and highly polymorphic. In addition, microsatellites have a wide distribution in the genome and can be efficiently identified, which is essential in studies on the genetic variability of populations (Balloux and Lugonae, 2002). However, the use of microsatellite markers requires the development of specific primers, *i.e.*, the unique DNA sequences flanking SSR loci. Given that the development of primers can be costly, several studies have tested for possible heterologous amplifications in a target species using primers already described for evolutionarily related species or even for evolutionarily related genera (Zhan and Fu, 2008). Many studies have successfully demonstrated heterologous amplification (Barbosa *et al.*, 2006) thereby lowering the cost of future projects. However, the loci detected by successful heterologous amplification should also be tested for their ability to assess polymorphism in the genome of the target species. Therefore, the aim of this study was to investigate the heterologous amplification of microsatellite loci originally developed from rohu and to evaluate genetic variability in 3 riverine stocks of *L. fimbriatus*.

## Materials and methods

Individuals of *L. fimbriatus* were collected from river Krishna (n=30), Mahanadi (n=30) and Kaveri (n=50) between 2009 to 2011 (Fig. 1). DNA was isolated from fin tissues following standard phenol chloroform method (Sambrook *et al.*, 2001) and quantified using 0.8% agarose gel. PCR amplification was done employing heterologous primers (n=30) which were originally isolated from rohu, taking *L. fimbriatus* DNA (25 ng  $\mu\text{L}^{-1}$ , each) as template. A touchdown PCR protocol was adopted and amplifications performed in a GeneAmp 9700 thermocycler (Applied Biosystems, USA). PCR was performed in a 10  $\mu\text{L}$  reaction volume containing 25 ng  $\mu\text{L}^{-1}$  genomic DNA, 2.5 pmol of each forward and reverse primers, 200  $\mu\text{M}$  of each dNTP, 2 mM  $\text{MgCl}_2$  and 0.2 units *Taq* DNA polymerase (Bangalore Genei, India). Amplification was carried out with the

following temperature profiles: one cycle at 94 °C for 5 min, 35 cycles at 94 °C for 45 sec, 55 °C for 1.5 min and 72 °C for 2 min and a final extension of 72 °C for 7 min. The PCR products were concentrated on a vacuum concentrator and then added with 5  $\mu\text{L}$  loading dye each (98% deionised formamide, 0.05% bromophenol blue, 0.05% xylene cyanol) and heated at 95 °C for 5 min and immediately chilled on ice. 4  $\mu\text{L}$  aliquots of each PCR products were analysed on a 6% denaturing PAGE Gel. The samples along with DNA standard,  $\phi\text{x}174/\text{HindIII}$  (Finnzymes, Finland) were subjected to electrophoresis at constant power supply of 25 watt for 2 h approximately at a constant temperature of 50 °C. Gels were silver stained and documented by visual counting.

Descriptive statistics like average number of alleles per locus ( $A$ ), expected proportion of heterozygotes ( $H_e$ ), observed proportion of heterozygotes ( $H_o$ ) and test of Hardy-Weinberg equilibrium (HWE) for each population were estimated using the software GDA, version 1.0, (Lewis and Zaykin, 1999). The estimates of  $H_e$  was unbiased for sample size (Weir and Ott, 1996). Pairwise linkage disequilibrium (LD) among loci was tested using the allele frequency for loci that were in HWE and genotypic frequencies were used to calculate LD of loci that deviated from HWE to prevent interference from within-locus disequilibrium. Assessment of genetic structure was done using the infinite allele model (Kimura and Crow, 1964) and  $F_{st}$  values were estimated using co-ancestry identity (Weir and Cockerham, 1984). F-statistics were estimated by bootstrap analysis using 1000 replicates, with a 95% confidence interval, using same software.



Fig. 1. Map showing the rivers and the spots from where the samples were collected (star marks indicate the sampling sites).

## Results and discussion

In *L. fimbriatus* only 15 out of the 30 microsatellite markers of rohu were successfully cross amplified (Table 1) and 3 markers showed polymorphism in all the 3 riverine populations. In total, only 9 alleles were found in 110 individuals representing 3 natural populations of *L. fimbriatus*. The average number of alleles, observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ) and p value for HWE per locus per population are presented (Table 2). Further, populationwise genetic diversity measures are also presented (Table 3). For each population, expected heterozygosity ( $H_e$ ) was generally lower than the observed heterozygosity ( $H_o$ ) leading to negative inbreeding coefficients ( $F_{is}$ ). Exact test for Hardy Weinberg Equilibrium (HWE) showed Lro-26 ( $p=0.004$ ) in Kaveri population to be significantly deviating from HWE at 5% significance level (Table 2). No significant ( $p>0.05$ )

linkage disequilibrium was detected between different genotypes at each of the different microsatellite loci. Bootstrapping over loci to obtain confidence intervals for F statistic analysis showed no significant levels (the 95% confidence region of these estimates span 0.0) of these estimates (Table 4).

In order to know how the genetic variation is distributed within and among populations, the following hypotheses were tested. The first two null hypotheses tested were that the allelic distribution is identical across populations and between population pairs for each locus. The second two null hypotheses tested were that the genotypic distribution is identical across populations and between population pairs. Though, Mahanadi population was found to be genetically more different from Kaveri and Krishna (Fig. 2; Table 5) the  $F_{st}$  values among populations were not significantly different from 0.0.

Table 1. List of rohu microsatellite loci that are amplified in *L. fimbriatus*.

| Locus name | Accession no. | Repeat types | Primer 5'-3'                                                        | <i>L. rohita</i><br>(product sizes) | <i>L. fimbriatus</i><br>(product sizes) |
|------------|---------------|--------------|---------------------------------------------------------------------|-------------------------------------|-----------------------------------------|
| Lr-03      | AJ507520      | (TG)19       | F ATC TGG CTG CCT ATT CAC C<br>R CAT CGG CGA CTG CAC TGG A          | 155                                 | 145                                     |
| Lr-10      | AJ507523      | (CA)13       | F GAT CTT CAG CGC CAG CGT G<br>R GAG GAC CTG CCC AGC ATG            | 245                                 | 240                                     |
| Lr-24      | AJ831438      | (TG)17       | F CAA GGC GAA AAG TGT CCA T<br>R AGG AAA TTG GTA AAG TGT TTC        | 165                                 | 160                                     |
| Lr-26      | AJ831439      | (TG)8        | F CCA GGG AGC TGC TAA GAA T<br>R AGC GCT TCA TGC AGT CTA C          | 145                                 | 140                                     |
| Lr-28      | AM231177      | (AC)18       | F TTCACGGACAGATTTGACCCAG<br>R AGTCTTTTCAGGAGATTAGCAG                | 180                                 | 170                                     |
| Lr-30      | AM231179      | (AC)15       | F ACGCGCTAGGGTCGTACAGTG<br>R CAGCATCATGTTAAGCGCTGTC                 | 185                                 | 180                                     |
| Lr-35      | AM269525      | (CA)13       | F TGT GAA CAT GCA AGC TCT CAG<br>R CTA GTC CCA CTC TAG TCA GCA      | 155                                 | 145                                     |
| Lr-36      | AM269526      | (CA)10       | F AGC GTG TCT GAT GTG TGA AAG G<br>R TCA GAT GCC TCC TGC ATT CTG    | 175                                 | 170                                     |
| Lr-38      | AM269528      | (GT)12       | F ATA GCA TCA CCA TCT GTT GGT G<br>R TCT GCT TCA GTC ACT CAG CAC    | 150                                 | 145                                     |
| Lr-39      | AM269529      | (GT)11       | F GTT TTT ATT CAG ACA GTC AGA C<br>R GTG AGA ATC CAG AGT GTC AC     | 160                                 | 155                                     |
| Lro-6      | AM184133      | (GT)29       | F TCT CCA CCC TCC ATA CCA CGA<br>R TGC CCT GCA TTT CTC CCA TCC      | 185                                 | 180                                     |
| Lro-26     | AM184144      | (GT)17       | F AGA TCA TTG CTG GGG AGT GTT TAT<br>R GAC CTG CCT GTG CCA TCT GTA  | 205                                 | 200                                     |
| Lro-31     | AM184147      | (GT)22       | F CAT AAT AGC AGT GGC GAG CAG<br>R AAC CAC CAG CAC ACC TTT CAC      | 180                                 | 185                                     |
| Lro-37     | AM184153      | (CA)16       | F ATG TTG TGG TCA TCA TGT AAA TC<br>R C AGT TTC CTC CCT TCA TAG TTT | 170                                 | 175                                     |

Table 2. No. of individuals (n), no. of alleles (A), alleles per locus ( $A_p$ ), expected heterozygosity ( $H_e$ ), observed heterozygosity ( $H_o$ ),  $F_{is}$  value (f) and p value for testing Hardy–Weinberg equilibrium in three populations of *L. fimbriatus*.

| Locus name | Parameters | Krishna | Mahanadi | Kaveri |
|------------|------------|---------|----------|--------|
| Lr-36      | n          | 24      | 30       | 27     |
|            | A          | 3       | 3        | 3      |
|            | $A_p$      | 3       | 3        | 3      |
|            | $H_e$      | 0.642   | 0.472    | 0.645  |
|            | $H_o$      | 0.833   | 0.433    | 0.814  |
|            | f          | -0.304  | 0.083    | -0.268 |
|            | p          | 0.115   | 0.079    | 0.265  |
| Lro-26     | n          | 25      | 27       | 28     |
|            | A          | 3       | 3        | 3      |
|            | $A_p$      | 3       | 3        | 3      |
|            | $H_e$      | 0.649   | 0.561    | 0.592  |
|            | $H_o$      | 0.520   | 0.370    | 0.214  |
|            | f          | 0.203   | 0.344    | 0.642  |
|            | p          | 0.064   | 0.060    | 0.004  |
| Lr-30      | n          | 28      | 28       | 26     |
|            | A          | 3       | 3        | 3      |
|            | $A_p$      | 3       | 3        | 3      |
|            | $H_e$      | 0.575   | 0.607    | 0.631  |
|            | $H_o$      | 0.678   | 0.892    | 0.846  |
|            | f          | -0.182  | -0.483   | -0.349 |
|            | p          | 0.500   | 0.078    | 0.080  |


Table 3. Descriptive statistics (by population) of *L. fimbriatus*, where n= no. of individuals, A = no. of alleles,  $A_p$  = alleles per locus,  $H_e$  = expected heterozygosity,  $H_o$  = observed heterozygosity, f =  $F_{is}$  value (inbreeding coefficient)

| River    | n  | A | $A_p$ | $H_e$ | $H_o$ | f      | Bootstrap confidence intervals for f<br>(1000 replications, at nominal<br>confidence interval = 95%) |        |
|----------|----|---|-------|-------|-------|--------|------------------------------------------------------------------------------------------------------|--------|
| Krishna  | 30 | 3 | 3     | 0.622 | 0.677 | -0.089 | Upper                                                                                                | 0.203  |
|          |    |   |       |       |       |        | Lower                                                                                                | -0.304 |
| Mahanadi | 30 | 3 | 3     | 0.546 | 0.565 | -0.034 | Upper                                                                                                | 0.344  |
|          |    |   |       |       |       |        | Lower                                                                                                | -0.483 |
| Kaveri   | 50 | 3 | 3     | 0.623 | 0.625 | -0.002 | Upper                                                                                                | 0.642  |
|          |    |   |       |       |       |        | Lower                                                                                                | -0.349 |
| Mean     |    | 3 | 3     | 0.597 | 0.622 | -0.042 |                                                                                                      |        |

Table 4. F-statistics and its bootstrap values of overall populations and loci; where  $f=F_{is}$ 

| Locus name                                                                                                    | f                           | $F_{it}$                    | $F_{st}$                    |
|---------------------------------------------------------------------------------------------------------------|-----------------------------|-----------------------------|-----------------------------|
| Lr-36                                                                                                         | -0.173562                   | -0.106508                   | 0.057137                    |
| Lro-26                                                                                                        | 0.400378                    | 0.392183                    | -0.013668                   |
| Lr-30                                                                                                         | -0.340769                   | -0.349937                   | -0.006838                   |
| Mean                                                                                                          | -0.035292                   | -0.022204                   | 0.012641                    |
| Bootstrap confidence intervals for F- statistics<br>(1000 replications, at nominal confidence interval = 95%) | Upper 0.400<br>Lower -0.340 | Upper 0.392<br>Lower -0.349 | Upper 0.057<br>Lower -0.013 |

Despite the popularity of microsatellites in the study of population genetics, their development requires substantial time, financial as well as technical resources.

Fig. 2. Dendrogram based Nei's (1972) genetic distance summarising variations among 3 populations of *L. fimbriatus* using heterologous microsatellite markers

Cross-species amplification is an alternative strategy to extend the utilisation of microsatellites across related species. Microsatellite loci generally show considerable evolutionary conservation, suggesting that primers developed for any one species may often be useful across a

Table 5. Pairwise comparison of  $F_{ST}$  values (below diagonal) and their bootstrap values of 1000 replications at nominal confidence interval = 95% (above diagonal) among three riverine populations of *L. fimbriatus*.

| River    | Krishna | Mahanadi                    | Kaveri                       |
|----------|---------|-----------------------------|------------------------------|
| Krishna  | 0       | Upper 0.066<br>Lower -0.008 | Upper -0.003<br>Lower -0.015 |
| Mahanadi | 0.019   | 0                           | Upper 0.104<br>Lower -0.025  |
| Kaveri   | -0.009  | 0.025                       | 0                            |

wide range of taxa. However, one drawback of heterologous primers is that mutations in the flanking sequences, to which PCR primers are designed to anneal, can result in non-amplifying PCR null alleles (Hoffman and Amos, 2005; Selkoe and Toonen, 2006). Null alleles produce an apparent heterozygote deficiency in a sample due to mis-scoring of heterozygotes as homozygotes. Heterozygote deficiency can also reflect various biological processes such as inbreeding, Wahlund effects and selection (Van Oosterhout *et al.*, 2004). It is very important to be able to determine the source of heterozygote deficiency observed in a sample. However, in the present study, only one locus deviated from HWE at 0.05% significance level (Lro-26; Kaveri population).

Present study was conducted with one of the objectives being, evaluation of usefulness of rohu microsatellite loci in population genetic study of *L. fimbriatus*. Though *L. fimbriatus* belongs to the same genus as *L. rohita*, only 15 loci (50%) out of 30 were successfully cross amplified in this species. This finding is inconsistent with the earlier observation of cross species amplification conducted in same genus where nearly 90% loci were cross amplified (Das *et al.*, 2005). Number of polymorphic loci of 3 (20%) found in this study was also very low as compared to rohu loci when checked in *L. bata* (47%) (Patel *et al.*, 2010). This indicates that genome complexity of *L. fimbriatus* is quite diverse from other species of *Labeo* genus. However, large numbers of loci may be needed to support this.

The second objective of this study was to estimate genetic variability of riverine stocks of this species. Obligate freshwater fishes are expected to display greater levels of genetic differentiation and population subdivision than marine species due to the isolating nature of river systems and small effective population sizes (Ward *et al.*, 1994). In the present study, data reveals that the levels of genetic differentiation among populations of *L. fimbriatus* were not significant (Table 4) and estimates of  $F_{IS}$  for each population was also found to be negative indicating excess heterozygosity. Overall  $F_{ST}$  for all samples combined was

found to be 0.012 indicating approximately 1% genetic variation caused by genetic differentiation of *L. fimbriatus*. Similar finding (2%) was reported in *L. dero* (Chaturvedi *et al.*, 2011) while studying Himalayas at central plateau of riverine populations in India and Queensland population of *Hypseleotris compressa* ( $F_{ST}=0.014$ ) (McGlashan and Hughes, 2001), but significantly different ( $F_{ST}=0.158$ ) from *L. rohita* populations of Indian rivers when analysed using MtDNA cytochrome b region (Luhariya *et al.*, 2011). Negative  $F_{IS}$  value and no significant  $F_{ST}$  among populations of *L. fimbriatus* may be attributed to less aquaculture practice of this species as compared with commonly cultivated species like rohu and catla.

Several evolutionary forces like random genetic drift, migration, mutation and their mutual interactions act on the wild populations and influence the pattern of genetic differentiation (Avice, 1994). Random genetic drift tends to cause genetic differentiation, after subpopulations are fragmented and gene flow between them is either reduced or absent. *L. fimbriatus* is primarily known to be a fringed lip peninsular carp inhabiting parts of rivers in foothill regions that have substrata having rocky bed. It is highly unlikely that in the present geographical scenario, *L. fimbriatus* in the rivers have the chance of intermixing. The rivers studied are more than 200 km apart and no water connection exists among them. Therefore, possible explanation for no significant genetic differentiation among populations, is that *L. fimbriatus* sampled from different river basins in the present study could have common ancestral gene pool. It appears that there has been at least historical connectivity between populations now inhabiting different rivers. Secondly, use of only three polymorphic markers may be highly insufficient to draw valid conclusions. The study suggests that stocks of *L. fimbriatus* are panmictic and genetic variation exists within and between populations.

## Acknowledgments

The work presented here is part of the ongoing program entitled “Outreach activity on fish genetic stocks” with financial support from Indian Council of Agricultural Research, New Delhi. We are thankful to the Director, CIFA for providing facilities to carry out the work. Thanks are due to Dr. A. E. Eknath, former Director, CIFA for his support. We are also thankful to Dr. S. Ayyappan, Secretary DARE and Director General, ICAR for his support and guidance. Help and guidance provided by Dr. V. S. Giri, RRC (CIFA), Vijayawada, Dr. M. R. Raghunath, RRC (CIFA), Bangalore during sample collection are duly acknowledged.



## References

- Avise, J. C. 1994. *Molecular markers, natural history and evolution*, 1<sup>st</sup> edn. Chapman and Hall, New York, USA., 551 pp.
- Ayyappan, S., Gopalakrishnan, A. and Kumar, G. 2009. Species diversification in aquaculture and domestic fish marketing in India. In: *MPEDA Souvenir 2009*, Indaqua 2009, Bhubaneswar, 21-23 January 2009. p.13-22.
- Balloux, F. and Lugonae, M. N. 2002. The estimation of population differentiation with microsatellite markers. *Mol. Ecol.*, 11: 155-165.
- Barbosa, A. C. D. R., Corrasa, T. A., Galzerani, F., Galettijr, P. M. and Hatanaka, T. 2006. Thirteen polymorphic microsatellite loci in the neotropical fish *Prochilodus argenteus* (Characiformes, Prochilodontidae). *Mol. Ecol. Notes*, 6: 936-938.
- Biswal, J. R., Sahoo, P. K. and Mohanty, P. K. 2010. Karyotypic diversity of some freshwater fish species in rural ponds of Orissa, India. *Cytologia*, 75: 237-242.
- Chaturvedi, A., Mohindra, V., Singh, R. K., Lal, K. K., Punia, P., Bhaskar, R., Mandal, A., Narain, L. and Lakra, W. S. 2011. Population genetic structure and phylogeography of cyprinid fish, *Labeo dero* (Hamilton, 1822) inferred from allozyme and microsatellite DNA marker analysis. *Mol. Bio. Rep.*, 38: 3513-3529.
- Das, P., Barat, A., Meher, P. K., Ray, P. P. and Majumdar, D. 2005. Isolation and characterisation of polymorphic microsatellites in *Labeo rohita* and their cross species amplification in related species. *Mol. Ecol. Notes*, 5: 231-233.
- Gjedrem, T. and Baranski, M. 2009. Selective breeding in aquaculture: An introduction, 1<sup>st</sup> edn. Springer Hedenberg, 221 pp.
- Hoffman, J. I. and Amos, W. 2005. Microsatellite genotyping errors: detection approaches, common sources and consequences for paternal exclusion. *Mol. Ecol.*, 14: 599-612.
- Jena, J., Das, P. C., Mitra, G., Patro, B., Mohanta, D. and Mishra, B. 2011. Evaluation of growth performance of *Labeo fimbriatus* (Bloch), *L. gonius* (Hamilton) and *Puntius gonionotus* (Bleeker) in polyculture with *Labeo rohita* (Hamilton) during fingerlings rearing at varied densities. *Aquaculture*, 319:493-496.
- Kimura, M. and Crow, J. F. 1964. The number of alleles that can be maintained in a finite population. *Genetics*, 49: 725.
- Lal, K. K., Chauhan, T., Mandal, A., Singh, R. K., Khulbe, L., Ponniah, A. G. and Mohindra, V. 2004. Identification of microsatellite DNA markers for population structure analysis in Indian major carp, *Cirrhinus mrigala*. *J. Appl. Ichthyol.*, 20: 87-91.
- Lewis, P. O. and Zaykin, D. 2001. Genetic data analysis: Computer program for analysis of allelic data, Ver 1.0. Free program distributed by the authors over the internet available at <http://hydrodyction.eeb.uconn.edu/people/plewis/software.php>
- Luhariya, R. K., Lal, K. K., Singh, R. K., Mohindra, V., Punia, P., Chauhan, U. K., Gupta, A. and Lakra, W. S. 2011. Genetic divergence in wild population of *Labeo rohita* (Hamilton, 1822) from nine Indian rivers, analysed through MtDNA cytochrome b region. *Mol. Biol. Rep.*, DOI:10.1007/s11033-011-1140-4
- McConnell, S. K. J., Leamon, J., Skibinski, D. O. F. and Mair, G. C. 2001. Microsatellite markers from the Indian major carp species, *Catla catla*. *Mol. Ecol. Notes*, 1: 115-116.
- McGlashan, D. J. and Hughes, J. M. 2001. Low levels of genetic differentiation among populations of the freshwater fish *Hypseleotris compressa* (Gobiidae: Eleotridinae): implications for its biology, population connectivity and history. *Heredity*, 86: 222-233.
- Mridula, R. M., Manissery, J. K., Keshavanath, P., Shankar, K. M., Nandeesh, M. C. and Rajesh, K. M. 2003. Water quality, biofilm production and growth of fringe-lipped carp (*Labeo fimbriatus*) in tanks provided with two solid substrates. *Bioresour. Technol.*, 87: 263-267.
- Patel, A., Das, P., Swain, S. K., Meher, P. K., Jayasankar, P. and Sarangi, N. 2009. Development of 21 new microsatellite markers in *Labeo rohita* (rohu). *Anim. Genet.*, 40: 253-254.
- Patel, A., Das, P., Barat, A., Meher, P. K., and Jayasankar, P. 2010. Utility of cross-species amplification of 34 rohu microsatellite loci in *Labeo bata*, and their transferability in six other species of the cyprinidae family. *Aquacult. Res.*, 41: 590-593.
- Punia, P., Gupta, H. S., Singh, R. K., Mohindra, V., Lal, K. K., Chauhan, V. S. and Lakra, W. S. 2006. Polymorphic microsatellite markers isolated from partially enriched genomic library of *Chitala chitala*. *Mol. Ecol. Notes*, 6: 1263-1265.
- Rao, G. R. 2011. Observations on the age and growth, maturity and fecundity of *Labeo fimbriatus* (bloch) of the river Godavari. *Indian J. Fish.*, 21: 427-444.
- Sambrook, J. and Russell, D. W. 2001. *Molecular cloning: A laboratory manual*, 3<sup>rd</sup> edn. Cold Spring Harbor Laboratory Press, New York, USA.
- Saravanan, M., Alora, E. V., Ramesh, M., Malarvizhi, A. and Kavitha, C. 2011. Impact of endosulfan on certain hematological and biochemical parameters of catfish and *Labeo fimbriatus*: sublethal study. *Toxicol. Ind. Health*, 27: 555-562.
- Selkoe, K. A. and Toonen, R. J. 2006. Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecol. Let.*, 9: 615-629.
- Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M. and Shipley, P. 2004. Micro Checker: software for identifying and correcting genotyping errors in microsatellite data. *Mol. Ecol. Notes*, 4: 535-538.

- Ward, R. D., Grewe, P. M. and Smolenski, A. J. 1994. A comparison of allozymes and mitochondrial DNA in Atlantic salmon from Tasmania and from the ancestral population in Canada. *Aquaculture*, 126: 257-264.
- Weir, B. S. and Cockerham, C. C. 1984. Estimating F-statistics for the analysis of population structure. *Evolution*, 38: 1358-1370.
- Weir, B. S. and Weir, B. S. 1996. *Genetic data analysis II: methods for discrete population genetic data*, Sinauer Associates Sunderland, MA, 445 pp.
- Zhan, A. and Fu, J. 2008. Microsatellite DNA markers for the Chinese wood frog *Rana chensinensis* and tests for their cross utility in 15 ranid frog species. *Mol. Ecol. Res.*, 8: 1126-1129.