

Molecular cloning and characterisation of integrin gene in the mud crab *Scylla serrata* (Forsk., 1775)

Thanuja Subramanian, Jeena Kezhedath*, Gayatri Tripathi, Megha Kadam Bedekar and K. V. Rajendran

Aquatic Environment and Health Management Division, ICAR-Central Institute of Fisheries Education, Mumbai-400061, Maharashtra, India



Abstract

The present study reports the first full length sequence of the integrin gene of the mud crab *Scylla serrata* (Ss-integrin). Ss-integrin cDNA comprises an open reading frame (ORF) of 2349 bp encoding a putative protein of 782 amino acids (aa). The sequence showed high similarity with the integrin gene of *Scylla paramamosain* (97.06% nucleotide identity and 95.78% amino acid identity). mRNA expression of Ss-integrin was detected in haemocytes, heart, gills, hepatopancreas, gut and muscle tissues. The response of the gene to white spot syndrome virus (WSSV) was investigated in experimentally infected *S. serrata*. The virus challenge resulted in varying patterns of immune response of the gene in different tissues. Ss-integrin was significantly upregulated in haemocytes, heart, gill, and hepatopancreas post-WSSV challenge, showing a tissue-specific temporal correlation with WSSV infection. The study indicated the role of Ss-Integrin in the immune response against WSSV infection in the mud crab, a relatively resistant host of the virus.

Introduction

Crustaceans primarily exhibit innate immune responses (Zhang *et al.*, 2019), relying on cellular and humoral mechanisms such as encapsulation, phagocytosis, autophagy, melanisation, coagulation, and antimicrobial peptide (AMPs) synthesis (Chen *et al.*, 2019; Tran *et al.*, 2020; Kulkarni *et al.*, 2021). Pattern recognition receptors (PRRs), which are essential in identifying foreign pathogens, stimulate immunological responses in crustaceans (Zhou *et al.*, 2021). Among these receptors, integrin, a transmembrane receptor linking the extracellular matrix to the cytoskeleton, plays a crucial role and was first characterised by Tamkun *et al.* (1986). Integrin can signal bidirectionally across cell membranes, influencing cellular processes through "inside-out" and "outside-in" signaling (Faull and Ginsberg, 1996; Hynes, 2002; Zhu *et al.*, 2007). Studies have revealed the presence of integrin in various crustacean tissues, with varying expression levels (Tang *et al.*, 2012; Lin *et al.*, 2013; Zhong *et al.*, 2013; Shi *et al.*, 2020).

(Holmblad *et al.*, 1997). Notably, the integrin subunit β -integrin has been implicated in pathogen infections, including viral infections. Research on the Chinese white shrimp *Fenneropenaeus chinensis*, demonstrated that white spot syndrome virus (WSSV) induces the expression of Fc-Int (Tang *et al.*, 2012; Zhong *et al.*, 2013). Furthermore, integrin-related signaling pathways during WSSV stimulation may facilitate virus spread (Tang *et al.*, 2017). Shrimp integrin gene silencing was reported to prevent WSSV infection, highlighting the role of integrin as a WSSV receptor (Tang *et al.*, 2017). Lin *et al.* (2013) reported the downregulation of integrin in *Penaeus* (*Litopenaeus*) *vannamei* challenged with WSSV, and integrin characterised in *P. monodon* also showed responses against bacterial pathogens (Shi *et al.*, 2020).

Scylla serrata (Forsk., 1775), also known as the Indo-Pacific swamp crab or giant mud crab, is a commercially important species, contributing 2.2% of the total crustacean production (FAO, 2022). Among the many challenges in mud crab farming, viral infections, particularly WSSV, pose a significant threat to commercial success and crab survival (Lo *et al.*, 1996; Kanchanaphum *et al.*, 1998; Kou *et al.*, 1998;

*Correspondence e-mail:

jeena@cife.edu.in

Keywords:

Host defence mechanism, *Scylla paramamosain*, White spot syndrome virus

Received : 30.05.2026

Accepted : 09.09.2026

The first invertebrate integrin gene was discovered in the haemocytes of the freshwater crayfish *Pacifastacus leniusculus*

Supamattaya *et al.*, 1998; Rajendran *et al.*, 1999; Chen *et al.*, 2000; Chakraborty *et al.*, 2002; Lavilla-Pitogo and de la Pena, 2004). As mud crabs rely on innate immune components to combat pathogens, understanding these innate immune mechanisms is crucial for managing their health. However, studies on innate immune responses, especially pathogen recognition receptors in mud crabs, are limited (Cao *et al.*, 2013; Vidya *et al.*, 2016; Chen *et al.*, 2018; Kong *et al.*, 2018; Yang *et al.*, 2023). While the β -integrin gene has been sequenced in *Scylla paramamosain*, the characterisation of integrin in *S. serrata* is yet to be completed. Furthermore, since *S. serrata* has been reported to be comparatively resistant to WSSV (Somboonna *et al.*, 2010; Pratapa *et al.*, 2023), elucidating the role of integrin in responding to WSSV infection becomes significant. Against this background, the present study aims to characterise the integrin gene in *S. serrata* and to understand the gene's expression in response to WSSV infection. We hypothesise that Ss-integrin contributes to modulating antiviral immune responses in *S. serrata*, a species comparatively resistant to WSSV.

Materials and methods

Experimental animals

Healthy, juvenile wild caught mud crabs (*S. serrata*), weighing 110–145 g were collected from Versova, Mumbai, India and transported to the Aquatic Animal Health Laboratory of ICAR-Central Institute of Fisheries Education (ICAR-CIFE), Mumbai. A total of 81 crabs were used in this study, of which 9 individuals were used for gene sequencing, while the remaining 72 were allocated to the treatment and control groups. Three animals were pooled into a single biological replicate. The animals were reared at an ambient temperature of 28–30°C in rectangular plastic crates (24 l capacity) holding seawater at a salinity of 33‰. Hideouts were provided to minimise cannibalism. The crabs were fed chopped fish meat daily *ad libitum* and were monitored regularly throughout the experimental period.

Identification and characterisation of *S. serrata* integrin gene (Ss-integrin)

cDNA synthesis, PCR amplification and sequencing

The animals were first immobilised on ice, and haemolymph was drawn from the coxa of the swimming legs using a sterile syringe fitted with a 24-gauge needle in sodium citrate solution (4% w/v) at a 1:1 ratio. Haemocytes were collected from pooled haemolymph by centrifugation at 10000 *g* for 5 min at 4°C and were stored at -80°C until RNA isolation. Total RNA was isolated from the haemocytes of *S. serrata* using TRIzol® Reagent (Invitrogen, USA) following the manufacturer's protocol. The quality and quantity of extracted RNA were estimated using a Nanodrop spectrophotometer (Thermo Scientific, USA). The cDNA was synthesised using the First-Strand cDNA Synthesis Kit (Thermo Scientific, USA) following the manufacturer's protocol.

The primers were designed based on the conserved sequences of the β -integrin mRNA of *S. paramamosain* (Accession no. HQ260916.1). PCR was performed using the following cycling

conditions: 1 cycle of initial denaturation at 94°C for 2 min, followed by 35 cycles of denaturation at 94°C for 30 s, annealing at the primer-specific temperature as mentioned in Table 1 (T_m °C) for 1 min, and extension at 72°C for 5 min. DNA concentration was standardised to 100 ng per reaction. The primers used to generate the integrin gene sequences, amplicon size and the annealing temperature are summarised in Table 1. The amplified PCR products were electrophoresed on a 1.2% agarose gel and visualised under a gel documentation system (Vilber Lourmat, Germany). The PCR products were purified using a gel extraction kit (Thermo Scientific, USA). The eluted DNA was sequenced through Eurofins Genomics India Pvt Ltd., Bengaluru, India.

Table 1. Details of primers, amplicon size and annealing temperature (T_A) used in the study

Primer	Sequence 5'–3'	Amplicon size	T_A
Ss-Integrin -1	F-GGAGACCATGAGGAACCTTACAA R-GCACGGCGAAGATGATGTT	500	58
Ss-Integrin -2	F-GACGACGAGGTAACCTTCAACCT R-GACAGTGGACGCCAGAGTATCC	496	60
Ss-Integrin -3	F-CGATCGCCGGGTGTGT R-TGGAGTTGGAGAGGTCCATGA	401	59
Ss-Integrin -4	F-ATGAGGAGGAGAGGCATG R-CTGAATTATGGAGCTGGGTG	595	56
Ss-Integrin -5	F-ACAACGAGTACACGGCCTTT R-GCCGTGATCTCCAGGTGAA	367	56
Ss-Integrin -6	F-ACAACGAGTACACGGCCTTT R-TCAGTGTCTGCGCCTCA	1200	60
Ss-Integrin -7	F-CCTACACGGGCCGCTTC R-TCAGTGTCTGCGCCTCA	518	56

Bioinformatic analysis

Sequences were first assembled using Finch TV 1.4.0 Software, followed by a homology search utilising the NCBI BLAST tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Nucleotide sequence alignments were performed using the Clustal Omega multiple alignment program (<http://www.ebi.ac.uk/Tools/msa/clustalo/>). Open reading frame (ORF) was predicted using ORF Finder (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). Translation was performed by the ExPASy tool (<http://www.expasy.org/tools/>). Simple modular architecture research tool SMART (<http://smart.embl-heidelberg.de/>) was used to analyse the protein domain topology of the deduced amino acid sequence, and NetNGlyc predicted N-glycosylation sites in the coding region of Ss-integrin. The molecular weight and isoelectric point of the deduced Ss-integrin protein were predicted using the ExPASy ProtParam tool. The 3D structure of Ss-integrin was predicted using the SWISS-MODEL tool. Similar gene sequences from various crustacean species were downloaded from NCBI GenBank for phylogenetic tree construction using MEGA 11 software (Tamura *et al.*, 2021).

Functional characterization of mud crab integrin gene

Virus inoculum and experimental infection

White spot syndrome virus (WSSV) inoculum was prepared from WSSV-positive *P. (L.) vannamei* as per the procedure described by

Rajendran *et al.* (1999). Briefly, infected tissues (gills, stomach, and epidermal layer) were homogenised in sterile marine phosphate buffered saline (PBS) at a ratio of 1 g of tissue to 1 ml of PBS. The homogenate was centrifuged at 2000 rpm for 10 min at 4°C, and the resulting supernatant was passed through a 0.22 µm membrane filter to obtain the viral filtrate. Healthy mud crabs (*S. serrata*) that tested negative for WSSV by PCR were selected for the experimental infection. Crabs were intramuscularly injected with 100 µl of WSSV inoculum (1.22×10^9 copies per µl) into the coxa of the swimming legs. A control group received an equivalent volume (100 µl) of sterile PBS via identical route.

WSSV infection in the experimental animals was confirmed using WSSV-specific PCR with 146F2/146R2 primers (146F2: 5'-GTA ACT GCC CCT TCC ATC TCC A-3'; 146R2: 5'-TAC GGC AGC TGC TGC ACC TTG T-3'), and for quality verification of extracted DNA, PCR was conducted using decapod-specific primers, 143F (5'-TGC-CTT-ATC-AGC-TAC-GTT-CGA-TTG-TAG-3') and 143R (5'-TTC-AGA-CGT-TTT-GCA-ACC-ATACTT-CCC-3') (Lo *et al.*, 1996). Thermal cycling parameters for both conventional PCR assays comprised an initial denaturation at 95°C for 5 min; 30 cycles including annealing at 55°C for 1 min; and a final extension at 72°C for 1 min.

Viral load was then quantified using real-time PCR, as described by Deepika *et al.* (2014). Reactions were conducted in a total volume of 25 µl containing 12.5 µl of 2x SYBR Green master mix (Takara Bio Inc., Japan), 2 µl of template DNA (50 ng µl⁻¹), and 0.5 µl of each WSSV-gene-specific primer (wssv366F1 and wssv366R1; stock concentration of 10 µM). Amplification was carried out in 96-well plates using an ABI 7500 Real-Time PCR System (Applied Biosystems, USA) set at its default thermal profile. Viral copy numbers were calculated by extrapolating values from a standard curve generated using a 10-fold serially diluted wssv366 plasmid DNA standard.

Expression profile of integrin gene in healthy and experimentally infected mud crab

To study the normal expression of the gene, haemolymph and tissues such as heart, gill, hepatopancreas, gut and muscle were collected from nine healthy mud crabs, and samples from three animals were pooled into one biological replicate. Similarly, for post-infection gene expression analysis, crabs were divided into treatment (WSSV-injected) and control (PBS-injected) groups, and samples were collected at 0, 24, 48, and 72 h post-infection. A total of 72 crabs (36 control, 36 treatment) were sacrificed, with nine crabs per group at each time point, and tissues including haemolymph, heart, gill, hepatopancreas, gut, and muscle were collected. Samples from three crabs were pooled to form one biological replicate, resulting in three pooled samples per group per time point. Subsequently, RNA was isolated from these samples using TRIzol® Reagent (Invitrogen, USA), and cDNA was prepared as described in section 2.2.1. Primers for the real-time PCR were designed to target the conserved region of the integrin gene sequence obtained from *S. serrata*. Similarly, primers for the housekeeping gene EF-1α were also designed (Table 2). Integrin mRNA expression profile was determined using Light cycler® Real-time PCR system. The comparative Ct method was used to estimate the relative expression of mRNA. The difference in the Ct

Table 2. Details of primers used in the gene expression studies employing real time PCR

Primer	Sequence 5' –3'	Amplicon size	T _A
EF-1α	F-GGTGCTGGACAAGCTGAAGGC R-GGTTCCGGTGATCATGTTCTTGATG	138	58
Ss-Integrin -qPCR	F-CCACTCAAGGACAAGTGCAG R-CCTGTGTACCTTCGAGGACG	101	58

value between the integrin gene and the internal reference EF-1α gene (ΔCt beta integrin- ΔCt EF-1α) was calculated, and the relative expression level of the gene to EF α was determined using $2^{-\Delta\Delta\text{Ct}}$ method proposed by Livak (2001).

Statistical analysis

Statistical processing of gene expression data was executed using SPSS software (v16.0), with differences considered statistically significant at $p < 0.05$. To obtain high quality RNA and account for individual biological variation, tissue and haemolymph samples from three crabs were pooled into a single biological replicate (N=3 pooled biological replicates per treatment group per time point, totalling 9 crabs per group). Relative mRNA levels were quantified using the $2^{-\Delta\Delta\text{Ct}}$ method. Statistical comparisons were performed on normalised ΔCt values, while $2^{-\Delta\Delta\text{Ct}}$ values were utilised for graphical presentation. Differences in transcript levels between control and WSSV-infected groups across post-infection time points (24, 48, and 72 hpi) were determined independently for each tissue using one-way analysis of variance (ANOVA) and Duncan's new multiple-range *post-hoc* test. Results are presented as mean±standard error (SE) of three biological replicates, with distinct superscript letters (a–d) denoting significant inter-group differences.

Results

Sequence information of *S. serrata* integrin (Ss-integrin)

cDNA of *S. serrata* integrin (Ss-integrin) was amplified with different sets of primers, resulting in the amplification of 500, 496, 401, 367, 1200 and 518 bp amplicons, and the PCR products were then purified and sequenced. The partial sequences obtained from each amplicon were verified and assembled to obtain the full-length Ss-integrin cDNA sequence. The open reading frame (ORF) of Ss-integrin cDNA was 2349 nucleotides (nt) long and encodes a putative protein of 782 amino acids (AA) (Fig. 1). The BLAST analysis revealed that the Ss-integrin nucleotide sequence shares 97.06% identity with *S. paramamosain* integrin cDNA. The identified cDNA sequence of Ss-integrin was deposited in GenBank under the Accession no, OR360538.

The protein domain topology prediction of the Ss-integrin deduced amino acid sequence showed a signal peptide domain (1 to 27), a Von Willebrand adhesion factor (136 to 364 aa) and an integrin beta subunit (N terminal portion of extracellular region) (35 to 450 aa) in the extracellular domain, an EGF-like domain (450 to 504 aa), an integrin B tail domain (628 to 710 aa), a single transmembrane domain (711 to 733 aa) and an intracellular cytoplasmic domain (734 to 780 aa) (Fig. 2a).

```

1  ATGAGGAGGAGAGGCGATGGTGGCGCGGTGGTGTGGGGTGGTGTATGCCCGCGGTGTGTGAGGAGCAGCAGAGCCAGGGGCGAGGAGTGT  90
   M R R R R G M V A A V V L G V V I A A V C E E Q Q S Q G Q E C  30
91  AGGGCGAGAGGAGCGTGTGACAAATGCATCCAATCGCAAGGGTGTCTGGTGGTCAATCCCCAGAGCGGGGATCACTGCCAACTTGCC  180
   R A Q R T C D K C I Q S Q G C S W C I N P Q S G D H C Q L A  60
181  GCCAACTTTAATGCATCCTTATGTAAACCCAGTGACATCAGGAACCCACACACCGGTGACACTGGTACAGAATGAGGAGCTAACAGGC  270
   A N F N A S L C N P S D I R N P H N T V T L V Q N E E L T G  90
271  ACAATCCAGAAGAAAGAAAGAAATGGGACAAAAACGCGAGATCTACCAACTAAAGCCACAAAGAGTGAAGCTCCAGCTCGCAAGGGT  315
   T I Q K K R K E W D K N A E I Y Q L K P Q R V K L Q L R K G  120
316  TACCAACTAAAGCCACAAAGATGAAGCTCCAGCTCGCGCAAGGGTATGCCACAGAACTGGACATCAAGTACCGGCGGCCCTTGACTAC  405
   Y Q L K P Q R V K L Q L R K G M P Q K L D I K Y R K A L D Y  150
406  CCCGTCGACCTGTACTACTCTCATGGACCTCTCCAACCTCATGAAGGACGACAGGAGCATCTGGCCAAACGCGGGCGCTATTCTGGCAGAG  495
   P V D L Y Y L M D L S N S M K D D K E H L A N A G A I L A E  180
496  ACCATGAGGAACCTTACACACGCACTTACCTCGGCTTCGGCTCCTTCGTGACAAAGGTATGATGCGCTTTCACCAACACCCAGCTC  585
   T M R N L T T Q P T L G F G S F V D K V M M P P T N T T Q L  210
586  CATAATTACCTCCGTCGCGCGGGTGCAGCCCCCTTACTCCTTCAGGAATGACATGAAGCTTAGTCAGGACCACGCGCTTCACGGAG  675
   H N S P P G C A P P Y S F R N D M K L S Q D H A A F T E  240
676  CAAGTGTCCAACGCCAGATCTCTGGCAACCTGGACCATCCGAGGGCGGGTTGATGCACTGATGCGAGCCATGTGTGCACAGATCAG  765
   Q V S N A L D H P E G G F D A L M Q A M V C T D Q  270
766  ATTGTTGGCGGACAAAGGTGCGCCACATACTCGTCTTCTCCACTGACCGGAAGTTTCATCAGCTGGCGACGGAGCGGCTGCGAGAGTG  855
   I G W R D K V R H I L V F S T D A K F H H A G D G R L A G V  300
856  GTGGAGCCCAATGATGAGAAGTGTCACTGAACGAAAAAACGAGTACACGCGCTTTCAGCAAGTACGACTACCCATCCATTGCACAGATC  945
   V E P N D E K C H L N E K N E Y T A F D K Y D Y P S I A Q I  330
946  AACAAAGTGGCAGAGAAAAGAACATCAACATCATCTCCCGTGTGACAGAACGAGAGACTCTACCGCAACTTCAGAAACAGATCAAA  1035
   N K V A E E K N I N I I F A V L Q N E R L Y R E L Q K Q I K  360
1036  ACTTCTTCTTTGGCTTGTGAGGAAGGATCGGCGAAGCTGGTGAAGCTCATCAAGGATCAGTACAATAGCATCTCGAGCAGCGTGCTCC  1125
   T S S F G L L E E G S A N V V D L I K D Q Y N S I S S S V S  390
1126  CTCACAGACAATAGCACCAGCGCCATCACCATCAAGTACAGTCTCTCTGCAAGGGGAGTGAAGTCAGGGAGACCGAGCTGTGACGGC  1215
   L T D N S A I T I K Y S S C K G S E V R E T R D C D G  420
1216  ATTAGCGAGGACGACGAGGTAACTTCAACCTGGAGATCAGCGCCAAACAGTGTTCGGATGGAGAGCAATCCATTGTGGAGGTGAAGACT  1305
   I S E D D E V T F N L E I T A N T C S D G E Q S I V E V K T  450
1306  CTGGAGGCGAAGCTGATCTTGGATGAACTCTGTGTACTGTGACTGCCCCCATGATGTTCCAGCTTACCGTGCACCAACGCGGCG  1395
   L E A N V I L E I E L L C T C D C P H D V S S L P C K N G G  480
1396  TCCATGATATGCGCGGTGTGCGAGTGTCCGAGTGGCTTTACGGAGAGGAGTGTGAGTGTGCGAGTGGCAGCGTGTCCCAAGATATAAG  1485
   S M I C G V C E C P S G F Y G E E C Q C A S G S V S Q D I S  510
1486  GAGAACCAGGACCAGACAGCAAGTGCATGCGCAGCGAGAGGACACAAAGGCTTGCTCGGGCGCGCGCGCTGCAAGTGTGGCTGTGC  1575
   E N Q D Q T S K C M A S E K D T R P C S G R G A C K C G V C  540
1576  AACTGTTTCAAGCTGAGGAGGTGTCCGGGACTTACTGCCAGTGTAAAGCGGAAATGCATGAGCTCCAACAAGCTGGAGTGCTCGGG  1665
   N C F K P E E V S G T Y C Q C N R R K C M S S N K L E C S G  570
1666  CATGGCACATGTGACTGCAACAGTGTAGATGTAAAGAGGATACTCTGGCGTCCACTGTCTATTGTAAACGACCAAGACTGCATCCCGACA  1755
   H G T C D C N E C R C N E G Y S G V H C H C N D Q D C I P T  600
1756  GGGTCTGAAGAGTGTGCTCGGGCCACGGCAAGTGTGATTGTGGCGGTTGCATCTGTGACCAGCAGAAGAAGCTCACCTACACGGGCCGC  1845
   G S E E V V C S G H G K C D C G R C I C D Q Q K N V T Y E G R  630
1846  TTCTGTGAGGACTGCGCTGAGTTGCGGTGGAGGAAAGTGCACAAAGTTTCAGAGACTGTGTGACAGTGTTCCTACTTTGGGATGGCCACTC  1935
   F C K D C L S C G G G K C K K F R D C V Q C F H F G I G P L  660
1936  AAGGACAACCTGCCAACAGTGTGAATCTCCACAGCTGTGGAGAACTCTGGATGACTACATGTACGAAAACCTCCCGCTGTGTACTCTCGAG  2025
   K D N E Q C E F S T A V E N L D D Y M Y E N S R L C T F E  690
2026  GACGAGGACGCGTGTCTACCTGAATTTACCTACCGCTACCTGGAGGGCCACCAACAAGGACGAGGTGCAGTGCAGGACAGAGAGGAAGTGC  2115
   D E D G C C Y L N F T Y R Y L E A T N K D E V H V G A E R K C  720
2116  CGTGAAAAGCCACCTTGCAGCCATCGTGTTCGCCCTGATCGGCACCATCGTGGGCATCGGCGTGTCTCCCTCATCTCTGGGAAGATC  2160
   R E K P P I A A I V F A L I G T I V G I G V I S L I L W K I  750
2161  ACCATCGTGGGCATCGGCGTCTCTCCCTCATTTCTGTGGAAGATCTTACCACCATCCACGACAAGCGGGAGTACGCCAAGTTCCTGGAG  2250
   F I V G I G V L S L I L W K I T T T I H D K R E Y A K F L E  750
2251  GAGACCAAGCGACCTCAGTTCAGCGCTGAGCAAAACCGCTGTTTATTGACCCCAACACCACACGACGAGAACCCGATGTTTGAAGGGCCAG  2340
   E T K R P Q P S A E Q N P L P I D P N T T I Q N P M P E G G  780
2341  AAACACTGA 2349
   K H * 782

```

Fig. 1. The nucleotide and deduced amino acid sequence of Ss-Integrin. Nucleotides and amino acids are numbered from left margin to right margin. The start codon (ATG) and stop codon (TGA) are marked in bold; Integrin beta subunit is in light grey, Integrin_B_tail is highlighted in dark grey, transmembrane domain is underlined and Integrin_B_cyt domain in deep green

Sequence homology and phylogenetic analysis

BLASTp analysis of the 782 deduced amino acid sequence of Ss-integrin showed the highest sequence similarity with the β -integrin gene of *S. paramamosain* (95.78 % identity), followed by *Portunus trituberculatus* (74.46%), *Eriocheir sinensis* (57.31%), *Pacifastacus leniusculus* (52.81%), *Penaeus chinensis* (51.01%) and *Penaeus japonicus* (50.69%). Eight N-linked glycosylation sites were identified in the coding region of the deduced amino acid sequence (Fig. 2b). The predicted molecular weight of the protein was 87.22 kDa, with an isoelectric point of 5.25. The 3-D structure

of Ss-integrin is shown in Fig. 2c. Phylogenetic analysis revealed the closest relationship between Ss-integrin and *S. paramamosain* integrin (Fig. 3).

Expression pattern of Ss-integrin gene in different tissues

Initially, Ss-integrin mRNA expression was examined in cDNA synthesised from RNA isolated from haemocytes and various tissues such as the heart, gill, hepatopancreas, intestine and muscles of healthy *S. serrata*. Semi-quantitative RT-PCR was




Fig. 2a. Schematic representation of the domain architecture of Ss-Integrin protein

MRRRGMVAAVVLGVVIAAVCEEQQSQGQECRAQRTCDKCIQSGCWCINPQSGDHCQLAANFNASLNCNPSDIRNPHNTV	80
TLVQNEELTGTIQKKRKEWDKNAEIYQLKPQRVKLQLRKGMPPQKLDIKYRRALDYPVDLYYLDLSNSMKDDKEHLANAG	160
AILAETMRNLTTQFTLGFSGFVDKVMMPFTNTQLHNSPPCAGCAPPYSFNRNDMKLSQDHAAFTQVSNQAISGNLDHPE	240
GGFDALMQAMVCTDQIGWRDKVRHILVFSTDAKFHHAGDGRLAGVVEPNDEKCHLNEKNEYTAFDKYDYPISIAQINKVAE	320
EKNINIIFAVLQNERLYRELQKQIKTSSFGLLEEGSANVVDLIKQYNSISSVSLTDNSTSAITIKYSSSCKGSEVRET	400
RDCDGISEDDDEVTFNLEITANTCSDGEQSIVEVKTLEANVILEIELLCTCDCPHDVSSLPCKNNGSMICGVCECPSGFYG	480
EECQCASGSVSQDISENQDQTSKCMASEKDTRPCSGRGACKCGVCNCFKPEEVSGTYQCNRRCMSSNKLECSGHGTCD	560
CNECRCNEGYSGVHCHCNDQDCIPTGSEEVCSGHGKDCGRCICDQQKNVTYTGRCEDCLSCGGGKCKKFRDCVQCFHF	640
GIGPLKDNCQCEFFSTAVENLDDMYENSRLCTFEDEDGCYLNFTYRYLEATNKDEVHVQAERKCREKPPIAAIVFALIG	720
TIVGIGVLSLILWKIFTTIHDKREYAKFLEETKRPQFSAEQNPLFIDPNTTTQNPMFEGQKH	
.....N.....N.....	80
.....	160
.....N.....N.....	240
.....	320
.....N.....	400
.....	480
.....N.....	560
.....N.....	640
.....N.....	720
.....N.....	800

Fig. 2b. Schematic representation of the domain architecture of Ss-Integrin protein




Fig. 2c. Predicted 3-D structure of Ss-Integrin protein by SWISS-MODEL




Fig. 3. Phylogenetic tree of deduced amino acid sequences of Ss-Integrin with other crustaceans

performed using Ss-integrin specific primers, with EF-1 α as the internal control. Ss-integrin transcripts were detected in all the tissues tested (Fig. 4). Subsequently, the transcript levels were estimated using real-time PCR. For this, three pooled samples were examined, and each pool comprised tissues collected from three individual animals. The highest expression of Ss-integrin was noticed in haemocytes, followed by the heart, gills, hepatopancreas, gut and muscle (Fig. 5).

Response of Ss-integrin gene to WSSV infection

Before determining the Ss-integrin gene expression in the WSSV-injected crab, the infection was confirmed in the experimentally infected crab using WSSV-specific PCR (Fig. 6a). Specific PCR amplification was detected in the gill tissue of crabs collected at 24, 48 and 72 hpi. Further, real-time PCR assay showed the highest viral

load (1.64×10^9 copies per 100 ng DNA) noted at 48 hpi compared to 1.6×10^7 and 8.75×10^7 copies per 100 ng DNA) observed at 24 and 72 hpi, respectively (Fig. 6b).

Ss-integrin gene expression kinetics following WSSV infection exhibited distinct tissue-specific patterns. Haemocytes displayed a classic rapid induction of Ss-integrin expression, reaching a maximum of ~ 8 -fold at 24 hpi, followed by a decline of ~ 6 -fold at 72 hpi (Fig. 7a). A similar early response was observed in the heart, where expression peaked at 5.5-fold at 24 hpi, dropped sharply to 2-fold at 48 hpi and remained at a similar level at 72 hpi (Fig. 7b). In contrast, the gill, hepatopancreas, and gut exhibited a delayed response, with Ss-integrin expression peaking at 48 hpi, followed by a decline at 72 hpi (Fig. 7c, d, e). The magnitude of upregulation in these tissues was comparatively lower with maximum increases of approximately 2.7, 1.75, and 2.25-fold in the gill, hepatopancreas and gut, respectively. Among the various tissues examined, no




Fig. 4. Semi-quantitative expression of Ss-Integrin in different tissues of healthy *S. serrata*. (a) Ss-Integrin; (b) EF-1 α . M - 100 bp plus molecular weight marker, Hc - Haemocytes, H - Heart, G - Gill, Hp - Hepatopancreas, M - Muscle, Gt - Gut




Fig. 5. Normal expression profile of Ss-Integrin in different tissues of healthy *S. serrata* estimated using qPCR




Fig. 6a. WSSV infection confirmed by WSSV-specific PCR. (a) WSSV-specific primer, (b) Decapod-specific primer. M - 100 bp plus molecular weight marker; C1-Control 24 hpi, C2-Control 48 hpi, C3-Control 72 hpi, T1-Treatment 24 hpi, T2-Treatment 48 hpi and T3-Treatment 72 hpi




Fig. 6b. Viral load determined in the gill tissue of WSSV-challenged mud crab *S. serrata* at different time points post-challenge

significant changes in Ss-integrin expression were observed in the muscle tissue at any time point following WSSV infection (Fig. 7f).

Discussion

Integrins are transmembrane receptor proteins that primarily recognize ligands found in the extracellular matrix, on cell surfaces, and in some soluble forms. Several researchers have indicated that

the integrin in invertebrates could participate in the regulation of cellular immune reactions such as melanisation, phagocytosis, nodulation and encapsulation in response to infection or injury (Lavine and Strand, 2003; Mamali *et al.*, 2009; Zhuang *et al.*, 2008). Despite this understanding, comprehensive research providing evidence regarding the involvement of invertebrate integrins, especially crustaceans, in innate immunity is yet to be undertaken. Against this background, identifying and characterising the integrin




Fig. 7. Expression profile of *Ss-Integrin* in different tissues of mud crab *S. serrata* at different time-points following viral challenge with WSSV. (a) Haemocytes; (b) Heart; (c) Gill; (d) Hepatopancreas; (e) Gut and (f) Muscle. The expression level in the control group was set as the baseline (1) in the bar graphs. Different letters indicate significant differences ($p < 0.05$) between different time-points within a treatment group compared to the control. Error bars represent $\pm$ SD of three independent PCR amplifications

gene of mud crab *S. serrata*, which is comparatively more resistant to WSSV than other crustaceans like shrimp (Somboonna *et al.*, 2010; Pratapa *et al.*, 2023), and further elucidating the response of the gene to WSSV is significant. Sequencing revealed that the *Ss-integrin* cDNA has a 2349 nt ORF encoding a 782 aa polypeptide. The ORF recorded for *Ss-integrin* is found to be shorter than that reported for the other crab species, *Eriocheir sinensis* (ORF of 2457 nt coding 818 aa) (Huang *et al.*, 2015). However, the domain architecture determined for *Ss-integrin* is similar to that reported for the integrin from *E. sinensis* by Huang *et al.* (2015). At the amino acid level, *Ss-integrin* showed 95.78 % identity, and the phylogenetic tree revealed a close evolutionary relationship between *Ss-integrin* and *S. paramamosain*. However, integrin genes from other crab species, crayfish and shrimp showed lower levels of sequence identity ranging from 50.69 to 74.46 %. The conserved domain architecture and relatively high amino acid sequence identity between *Ss-integrin* and integrin of other species indicate that, *Ss-integrin* may share functional characteristics with other members of the integrin

family, particularly in processes related to signal transduction and immune responses.

Integrins have been reported to be distributed throughout various tissues in crustaceans with varying degrees of expression (Tang *et al.*, 2012; Huang *et al.*, 2015). Analysis of transcript abundance using conventional and qPCR assays showed that *Ss-integrin* was expressed in haemocytes and a range of tissues such as heart, gill, hepatopancreas, intestine and muscle of healthy mud crabs. A similar tissue distribution has been reported in *Fenneropenaeus chinensis* (Tang *et al.*, 2017), suggesting *Ss-integrin* may have multiple biological functions. *Ss-integrin* was found to be highly expressed in haemocytes, consistent with observations in *E. sinensis* (Huang *et al.*, 2015), *P. vannamei* (Lin *et al.*, 2013; Zhang *et al.*, 2012), *P. chinensis* (Tang *et al.*, 2017), *P. monodon* (Shi *et al.*, 2020), the insect *Ostrinia furnacalis* (Hu *et al.*, 2010) and the tobacco hornworm, *Manduca sexta* (Levin *et al.*, 2005). The predominant expression of *Ss-integrin* in haemocytes suggests a potential role in cell-mediated immune

responses in *S. serrata*, given that haemocytes are key effector cells involved in crustacean immunity.

Integrin receptors on host cells have been identified as important targets of WSSV, as several WSSV proteins, including VP187, VP31, VP110, VP33, VP36A, VP36B, and VP281, contain the RGD motif and have been reported to interact with shrimp integrins during infection (Huang *et al.*, 2002; Tsai *et al.*, 2004; Li *et al.*, 2006; Xie *et al.*, 2006). Studies by Tang *et al.* (2012, 2017) in *F. chinensis* and Li *et al.* (2007) in *Marsupenaeus japonicus* showed that β -integrin can specifically bind and interact with WSSV, suggesting a possible role as a cellular receptor or post-attachment receptor or co-receptor during WSSV infection. In the present study, Ss-integrin expression following WSSV challenge, was investigated across various tissues, with the most pronounced and significant upregulation observed in haemocytes at all time points. In contrast, β -integrin was reported to be downregulated at 24 and 96 h post-WSSV challenge in *P. (L.) vannamei* (Lin *et al.*, 2013), indicating species specific differences in the regulation of integrin mediated responses to WSSV infection. The marked upregulation of Ss-integrin in haemocytes may indicate its involvement in the host response to WSSV, potentially through modulation of cell adhesion signal transduction and immune processes. However, whether the observed increase directly contributes to antiviral defence or influences WSSV attachment and entry remains to be experimentally established.

The high expression of Ss-integrin in haemocytes suggests that integrin might play a pivotal role in cell-mediated immune regulation in invertebrates (Zhuang *et al.*, 2007). Haemocytes are considered crucial target cells for WSSV in shrimp and exhibit high expression of immune related genes (Zhan *et al.*, 1999; Wang *et al.*, 2002; Wen *et al.*, 2013). In crustaceans, haemocytes and hepatopancreas are vital tissues involved in humoral immune responses, while the gills are also recognised as an important tissue associated with immune functions (Ji *et al.*, 2009; Garcia-Orozco *et al.*, 2012; Sun *et al.*, 2015). In the present study, Ss-integrin expression in the gills was significantly modulated at all time points following WSSV challenge, whereas the changes observed in the hepatopancreas were not statistically significant. This pattern differs from the observations made in *F. chinensis*, where *Fc β Int* mRNA was significantly upregulated in the gills post-WSSV challenge (Tang *et al.*, 2012; Zhong *et al.*, 2013). Previous studies have suggested that activation of integrin-related signalling pathways following WSSV exposure may facilitate viral infection in hosts by upregulating β -integrin expression (Tang *et al.*, 2017). Silencing of shrimp integrin gene has been reported to reduce WSSV infection and increase host survival following WSSV exposure (Tang *et al.*, 2017). However, whether Ss-integrin plays a similar role in WSSV infection *S. serrata* remains unclear. As *S. serrata* is a comparatively resistant host to WSSV, further functional studies, including integrin knockdown or blocking experiments, are needed to determine whether Ss-integrin contributes to WSSV attachment, entry, immune signalling or host defense.

In conclusion, the present study identified and characterised the integrin gene in *S. serrata* (Ss-integrin) for the first time. Our findings revealed baseline expression of Ss-integrin across multiple tissues with primary abundance in haemocytes and demonstrated significant transcriptional modulation following WSSV infection. While these results highlight Ss-integrin as a key virus-responsive

component of host immune dynamics, transcriptomic observations primarily reflect cellular reactivity rather than direct mechanistic involvement. Further investigations employing targeted functional approaches, such as RNA interference and protein-protein interaction studies, will be essential to elucidate its precise role as a cellular receptor, diagnostic biomarker, or therapeutic target in mud crab antiviral strategies.

Acknowledgements

The first author sincerely thanks ICAR-CIFE, Mumbai, for awarding the Institutional Fellowship during the study period. The authors also express their gratitude to the Director/Vice-Chancellor of ICAR-CIFE for providing the necessary facilities and support to carry out this research.

References

- Cao, J., Lin, Z., Zhang, Y., Zhang, X., Li, S., Zhang, N., Zou, W., Li, Y., Chen, J. and Wang, X. 2013. Cloning and characterization of the SpLRR cDNA from green mud crab, *Scylla paramamosain*. *Fish Shellfish Immunol.*, 34: 129-135. <https://doi.org/10.1016/j.fsi.2012.10.018>.
- Chakraborty, A., Otta, S. K., Joseph, B., Kumar, S. H., Hossain, M., Karunasagar, I., Venugopal, M. N. and Karunasagar, I. 2002. Prevalence of white spot syndrome virus in wild crustaceans along the coast of India. *Curr. Sci.*, 82(11): 1392-1397. <https://www.jstor.org/stable/24106017>.
- Chen, F. and Wang, K. 2019. Characterization of the innate immunity in the mud crab *Scylla paramamosain*. *Fish Shellfish Immunol.*, 93: 436-448. <https://doi.org/10.1016/j.fsi.2019.07.076>.
- Chen, L. L., Lo, C. F., Chiu, Y. L., Chang, C. F. and Kou, G. H. 2000. Natural and experimental infection of white spot syndrome virus (WSSV) in benthic larvae of mud crab *Scylla serrata*. *Dis. Aquat. Org.*, 40: 157-161. <https://doi.org/10.3354/dao040157>.
- Chen, Y., Aweya, J. J., Sun, W., Wei, X., Gong, Y., Ma, H., Zhang, Y., Wen, X. and Li, S. 2018. SpToll1 and SpToll2 modulate the expression of antimicrobial peptides in *Scylla paramamosain*. *Dev. Comp. Immunol.*, 87: 124-136. <https://doi.org/10.1016/j.dci.2018.06.008>.
- Deepika, A., Sreedharan, K., Paria, A., Makesh, M. and Rajendran, K. V. 2014. Toll-pathway in tiger shrimp (*Penaeus monodon*) responds to white spot syndrome virus infection: Evidence through molecular characterization and expression profiles of MyD88, TRAF6 and TLR genes. *Fish Shellfish Immunol.*, 41: 441-454. <https://doi.org/10.1016/j.fsi.2014.09.026>.
- FAO 2022. *The state of world fisheries and aquaculture 2022: Towards blue transformation*. Food and Agriculture Organization of the United Nations, Rome, Italy. <https://doi.org/10.4060/cc0461en>.
- Faull, R. J. and Ginsberg, M. H. 1996. Inside-out signaling through integrins. *J. Am. Soc. Nephrol.*, 7: 1091-1097. <https://doi.org/10.1681/ASN.V781091>.
- Garcia-Orozco, K. D., Sanchez-Paz, A., Aispuro-Hernandez, E., Gomez-Jimenez, S., Lopez-Zavala, A., Araujo-Bernal, S. and Muhlia-Almazan, A. 2012. Gene expression and protein levels of thioredoxin in the gills from the whiteleg shrimp (*Litopenaeus vannamei*) infected with two different viruses: The WSSV or IHHNV. *Fish Shellfish Immunol.*, 32: 1141-1147. <https://doi.org/10.1016/j.fsi.2012.03.020>.
- Holmblad, T., Thornqvist, P. O., Soderhall, K. and Johansson, M. W. 1997. Identification and cloning of an integrin beta subunit from hemocytes of the freshwater crayfish *Pacifastacus leniusculus*. *J. Exp. Zool.*, 277(3): 255-261.
- Hu, J., Zhao, H., Yu, X., Liu, J., Wang, P., Chen, J., Xu, Q. and Zhang, W. 2010. Integrin β 1 subunit from *Ostrinia furnacalis* hemocytes: Molecular characterization,

- expression, and effects on the spreading of plasmatocytes. *J. Insect Physiol.*, 56: 1846-1856. <https://doi.org/10.1016/j.jinsphys.2010.08.001>.
- Huang, C., Zhang, X., Lin, Q., Xu, X. and Hew, C. L. 2002. Characterization of a novel envelope protein (VP281) of shrimp white spot syndrome virus by mass spectrometry. *J. Gen. Virol.*, 83: 2385-2392. <https://doi.org/10.1099/0022-1317-83-10-2385>.
- Huang, Y., Zhao, L. L., Feng, J. L., Zhu, H. X., Huang, X., Ren, Q. and Wang, W. 2015. A novel integrin function in innate immunity from Chinese mitten crab (*Eriocheir sinensis*). *Dev. Comp. Immunol.*, 52: 155-165. <https://doi.org/10.1016/j.dci.2015.05.005>.
- Hynes, R. O. 2002. Integrins: Bidirectional, allosteric signaling machines. *Cell*, 110: 673-687. [https://doi.org/10.1016/S0092-8674\(02\)00971-6](https://doi.org/10.1016/S0092-8674(02)00971-6).
- Ji, P. F., Yao, C. L. and Wang, Z. Y. 2009. Immune response and gene expression in shrimp (*Litopenaeus vannamei*) hemocytes and hepatopancreas against some pathogen-associated molecular patterns. *Fish Shellfish Immunol.*, 27: 563-570. <https://doi.org/10.1016/j.fsi.2009.08.001>.
- Kanchanaphum, P., Wongteerasupaya, C., Sitidilokratana, N., Boonsaeng, V., Panyim, S., Tassanakajon, A., Wityachumnarnkul, B. and Flegel, T. W. 1998. Experimental transmission of white spot syndrome virus (WSSV) from crabs to shrimp *Penaeus monodon*. *Dis. Aquat. Org.*, 34: 1-7. <https://doi.org/10.3354/dao034001>.
- Kong, T., Gong, Y., Liu, Y., Wen, X., Tran, N. T., Aweya, J. J., Zhang, Y., Ma, H., Zheng, H. and Li, S. 2018. Scavenger receptor B promotes bacteria clearance by enhancing phagocytosis and attenuates white spot syndrome virus proliferation in *Scylla paramamosain*. *Fish Shellfish Immunol.*, 78: 79-90. <https://doi.org/10.1016/j.fsi.2018.04.027>.
- Kou, G. H., Peng, S. E., Chiu, Y. L. and Lo, C. F. 1998. Tissue distribution of white spot syndrome virus (WSSV) in shrimp and crabs. In: Flegel, T. W. (Eds.), *Advances in shrimp biotechnology*. National Center for Genetic Engineering and Biotechnology, Bangkok, Thailand, pp. 267-271.
- Kulkarni, A., Krishnan, S., Anand, D., Kokkattunivarthil Uthaman, S., Otta, S. K., Karunasagar, I. and Kooloth Valappil, R. 2021. Immune responses and immunoprotection in crustaceans with special reference to shrimp. *Rev. Aquac.*, 13: 431-459. <https://doi.org/10.1111/raq.12482>.
- Lavilla-Pitogo, C. R. and de la Pena, L. D. 2004. *Diseases in farmed mud crabs Scylla spp.: Diagnosis, prevention, and control*. SEAFDEC Aquaculture Department, Tigbauan, Iloilo, Philippines, 89 p. <http://hdl.handle.net/10862/1709>.
- Lavine, M. D. and Strand, M. R. 2003. Haemocytes from *Pseudoplusia includens* express multiple α and β integrin subunits. *Insect Mol. Biol.*, 12: 441-452. <https://doi.org/10.1046/j.1365-2583.2003.00428.x>.
- Levin, D. M., Breuer, L. N., Zhuang, S., Anderson, S. A., Nardi, J. B. and Kanost, M. R. 2005. A hemocyte-specific integrin required for hemocytic encapsulation in the tobacco hornworm, *Manduca sexta*. *Insect Biochem. Mol. Biol.*, 35: 369-380. <https://doi.org/10.1016/j.ibmb.2005.01.003>.
- Li, D. F., Zhang, M. C., Yang, H. J., Zhu, Y. B. and Xu, X. 2007. β -integrin mediates WSSV infection. *Virology*, 368: 122-132. <https://doi.org/10.1016/j.virol.2007.06.027>.
- Li, H., Zhu, Y., Xie, X. and Yang, F. 2006. Identification of a novel envelope protein (VP187) gene from shrimp white spot syndrome virus. *Virus Res.*, 115: 76-84. <https://doi.org/10.1016/j.virusres.2005.07.007>.
- Li, L., Lin, S. and Yang, F. 2006. Characterization of an envelope protein (VP110) of white spot syndrome virus. *J. Gen. Virol.*, 87: 1909-1915. <https://doi.org/10.1099/vir.0.81730-0>.
- Lin, Y. C., Chen, J. C., Chen, Y. Y., Liu, C. H., Cheng, W., Hsu, C. H. and Tsui, W. C. 2013. Characterization of white shrimp *Litopenaeus vannamei* integrin β and its role in immunomodulation by dsRNA-mediated gene silencing. *Dev. Comp. Immunol.*, 40: 167-179. <https://doi.org/10.1016/j.dci.2013.01.001>.
- Livak, K. J. and Schmittgen, T. D. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25: 402-408. <https://doi.org/10.1006/meth.2001.1262>.
- Lo, C., Ho, C., Peng, S., Chen, C., Hsu, H., Chiu, Y., Chang, C., Liu, K., Su, M., Wang, C. and Kou, G. 1996. White spot syndrome baculovirus (WSBV) detected in cultured and captured shrimp, crabs and other arthropods. *Dis. Aquat. Org.*, 27: 215-225. <https://doi.org/10.3354/dao027215>.
- Mamali, I., Lamprou, I., Karagiannis, F., Karakantza, M., Lampropoulou, M. and Marmaras, V. J. 2009. A β integrin subunit regulates bacterial phagocytosis in medfly haemocytes. *Dev. Comp. Immunol.*, 33: 858-866. <https://doi.org/10.1016/j.dci.2009.02.004>.
- Pratapa, M. G., Kumar, S., Bedekar, M. K., Sanath Kumar, H. and Rajendran, K. V. 2023. Experimental infection reveals mud crab, *Scylla serrata*, is less susceptible than *Scylla olivacea* and shrimp, *Penaeus vannamei*, to white spot syndrome virus (WSSV). *Aquaculture*, 577: 739877. <https://doi.org/10.1016/j.aquaculture.2023.739877>.
- Rajendran, K. V., Vijayan, K. K., Santiago, T. C. and Krol, R. M. 1999. Experimental host range and histopathology of white spot syndrome virus (WSSV) infection in shrimp, prawns, crabs and lobsters from India. *J. Fish Dis.*, 22: 183-191. <https://doi.org/10.1046/j.1365-2761.1999.00162.x>.
- Shi, M., Jiang, S., Li, Y., Yang, Q., Jiang, S., Yang, L., Huang, J. and Zhou, F. 2020. Comprehensive expression analysis of the beta integrin from *Penaeus monodon* indicating its participation in innate immunity and ammonia nitrogen stress response. *Fish Shellfish Immunol.*, 98: 887-898. <https://doi.org/10.1016/j.fsi.2019.11.049>.
- Somboonna, N., Mangkalan, S., Udompetcharaporn, A., Krittanai, C., Sritunyalucksana, K. and Flegel, T. 2010. Mud crab susceptibility to disease from white spot syndrome virus is species-dependent. *BMC Res. Notes*, 3: 315. <https://doi.org/10.1186/1756-0500-3-315>.
- Sun, S., Xuan, F., Fu, H., Zhu, J., Ge, X. and Gu, Z. 2015. Transcriptomic and histological analysis of hepatopancreas, muscle and gill tissues of oriental river prawn (*Macrobrachium nipponense*) in response to chronic hypoxia. *BMC Genomics*, 16: 491. <https://doi.org/10.1186/s12864-015-1701-3>.
- Supamattaya, K., Hoffmann, R., Boonyaratpalin, S. and Kanchanaphum, P. 1998. Experimental transmission of white spot syndrome virus (WSSV) from black tiger shrimp *Penaeus monodon* to the sand crab *Portunus pelagicus*, mud crab *Scylla serrata* and krill *Acetes* sp. *Dis. Aquat. Org.*, 32: 79-85. <https://doi.org/10.3354/dao032079>.
- Tamkun, J. W., DeSimone, D. W., Fonda, D., Patel, R. S., Buck, C., Horwitz, A. F. and Hynes, R. O. 1986. Structure of integrin, a glycoprotein involved in the transmembrane linkage between fibronectin and actin. *Cell*, 46: 271-282. [https://doi.org/10.1016/0092-8674\(86\)90744-0](https://doi.org/10.1016/0092-8674(86)90744-0).
- Tamura, K., Stecher, G. and Kumar, S. 2021. MEGA11: Molecular evolutionary genetics analysis version 11. *Mol. Biol. Evol.*, 38: 3022-3027. <https://doi.org/10.1093/molbev/msab120>.
- Tang, X., Wang, X. and Zhan, W. 2012. An integrin β subunit of Chinese shrimp involved in WSSV infection. *Aquaculture*, 368-369: 1-9. <https://doi.org/10.1016/j.aquaculture.2012.08.004>.
- Tang, X., Zhai, F., Sheng, X., Xing, J. and Zhan, W. 2017. The roles of β -integrin of Chinese shrimp (*Fenneropenaeus chinensis*) in WSSV infection. *Int. J. Mol. Sci.*, 18: 1465. <https://doi.org/10.3390/ijms18071465>.
- Tran, N. T., Kong, T., Zhang, M. and Li, S. 2020. Pattern recognition receptors and their roles in the innate immune system of mud crab (*Scylla paramamosain*). *Dev. Comp. Immunol.*, 102: 103469. <https://doi.org/10.1016/j.dci.2019.103469>.
- Tsai, J. M., Wang, H. C., Leu, J. H., Hsiao, H. H., Wang, A. H. J., Kou, G. H. and Lo, C. F. 2004. Genomic and proteomic analysis of thirty-nine structural proteins of shrimp white spot syndrome virus. *J. Virol.*, 78: 11360-11370. <https://doi.org/10.1128/JVI.78.20.11360-11370.2004>.

- Vidya, R., Makesh, M., Purushothaman, C. S., Chaudhari, A., Gireesh-Babu, P. and Rajendran, K. V. 2016. Report of leucine-rich repeats (LRRs) from *Scylla serrata*: Ontogeny, molecular cloning, characterization and expression analysis following ligand stimulation, and upon bacterial and viral infections. *Gene*, 590: 159-168. <https://doi.org/10.1016/j.gene.2016.06.026>.
- Wang, Y. T., Liu, W., Seah, J. N., Lam, C. S., Xiang, J. H., Korzh, V. and Kwang, J. 2002. White spot syndrome virus (WSSV) infects specific hemocytes of the shrimp *Penaeus merguensis*. *Dis. Aquat. Org.*, 52: 249-259. <https://doi.org/10.3354/dao052249>.
- Wen, R., Li, F., Sun, Z., Li, S. and Xiang, J. 2013. Shrimp MyD88 responsive to bacteria and white spot syndrome virus. *Fish Shellfish Immunol.*, 34: 574-581. <https://doi.org/10.1016/j.fsi.2012.11.034>.
- Xie, X. and Yang, F. 2006. White spot syndrome virus VP24 interacts with VP28 and is involved in virus infection. *J. Gen. Virol.*, 87: 1903-1908. <https://doi.org/10.1099/vir.0.81570-0>.
- Yang, Q. F., Li, S., Feng, G. P., Qin, C., Min, X. W., Fang, W. H., Wu, Y., Zhou, J. and Li, X. C. 2023. A novel C-type lectin (SpccCTL) suppresses MCRV replication by binding viral protein and regulating antiviral peptides in *Scylla paramamosain*. *Fish Shellfish Immunol.*, 142: 109143. <https://doi.org/10.1016/j.fsi.2023.109143>.
- Zhan, W. B., Wang, Y. H., Fryer, J. L., Okubo, K., Fukuda, H., Yu, K. K. and Meng, Q. X. 1999. Production of monoclonal antibodies (MAbs) against white spot syndrome virus (WSSV). *J. Aquat. Anim. Health*, 11: 17-22. [https://doi.org/10.1577/1548-8667\(1999\)011%3C0017:POMAMA%3E2.0.CO;2](https://doi.org/10.1577/1548-8667(1999)011%3C0017:POMAMA%3E2.0.CO;2).
- Zhang, X., Tang, X., Tran, N. T., Huang, Y., Gong, Y., Zhang, Y., Zheng, H., Ma, H. and Li, S. 2019. Innate immune responses and metabolic alterations of mud crab (*Scylla paramamosain*) in response to *Vibrio parahaemolyticus* infection. *Fish Shellfish Immunol.*, 87: 166-177. <https://doi.org/10.1016/j.fsi.2019.01.011>.
- Zhang, Y., Wang, L., Wang, L., Wu, N., Zhou, Z. and Song, L. 2012. An integrin from shrimp *Litopenaeus vannamei* mediated microbial agglutination and cell proliferation. *PLoS ONE*, 7(7): e40615. <https://doi.org/10.1371/journal.pone.0040615>.
- Zhong, R., Tang, X., Zhan, W., Xing, J. and Sheng, X. 2013. Expression kinetics of β -integrin in Chinese shrimp (*Fenneropenaeus chinensis*) hemocytes following infection with white spot syndrome virus. *Fish Shellfish Immunol.*, 35: 539-545. <https://doi.org/10.1016/j.fsi.2013.05.004>.
- Zhou, J., Zhou, J. F., Wang, Y., Feng, G. P., Fang, W. H., Kang, W., Ma, L. B. and Li, X. C. 2021. SpSR-B2 functions as a potential pattern recognition receptor involved in antiviral and antibacterial immune responses of mud crab *Scylla paramamosain*. *Int. J. Biol. Macromol.*, 193: 2173-2182. <https://doi.org/10.1016/j.ijbiomac.2021.11.048>.
- Zhu, J., Carman, C. V., Kim, M., Shimaoka, M., Springer, T. A. and Luo, B. H. 2007. Requirement of α and β subunit transmembrane helix separation for integrin outside-in signaling. *Blood*, 110: 2475-2483. <https://doi.org/10.1182/blood-2007-03-080077>.
- Zhuang, S., Kelo, L., Nardi, J. B. and Kanost, M. R. 2007. An integrin-tetraspanin interaction required for cellular innate immune responses of an insect, *Manduca sexta*. *J. Biol. Chem.*, 282: 22563-22572. <https://doi.org/10.1074/jbc.M700341200>.
- Zhuang, S., Kelo, L., Nardi, J. B. and Kanost, M. R. 2008. Multiple α subunits of integrin are involved in cell-mediated responses of the *Manduca* immune system. *Dev. Comp. Immunol.*, 32: 365-379. <https://doi.org/10.1016/j.dci.2007.07.007>.