



## Analysis of fishery and stock of the portunid crab, *Charybdis feriata* (Linnaeus, 1758) from Veraval waters, north-west coast of India

GYANARANJAN DASH, SWATIPRIYANKA SEN, K. MOHAMMED KOYA, K. R. SREENATH R. THANGAVELU, SURESH KUMAR MOJJADA AND M. S. ZALA

Veraval Regional Centre of Central Marine Fisheries Research Institute, Matsya Bhavan, Bhidia, Veraval - 362 269 Gujarat, India

e-mail: gyanranjandashcmfri@gmail.com

### ABSTRACT

Fishery and stock characteristics of the portunid crab *Charybdis feriata* from the waters off the north-west coast of India was studied based on commercial trawl landings at Veraval during 2009-2011. Growth parameters  $L_{\infty}$ ,  $K$  and  $t_0$  were estimated as 171.5 mm, 0.97 yr<sup>-1</sup> and -0.10 yr respectively. Mortality parameters  $Z$ ,  $M$  and  $F$  were 3.97 yr<sup>-1</sup>, 1.49 yr<sup>-1</sup> and 2.48 yr<sup>-1</sup> respectively. The length at capture ( $L_{50}$ ) and length at maturity ( $L_{M50}$ ) were 95.4 mm and 87.3 mm, respectively which indicates that crabs enter into peak exploitation phase after attaining sexual maturity. The bio-economic stock assessment model showed that the species can be exploited at the present fishing level to maintain the economy of the fishery at maximum level (MEY) or the effort can be increased by 2.4 times to maximise the yield (MSY) depending upon the management objective. However, since the stock recruitment relationship is not defined, as a precautionary approach, the effort should not be increased more than 1.3 times the present level to conserve the spawning stock.

Keywords: *Charybdis feriata*, Fisheries management, Management reference point, Spawning stock biomass, Stock assessment

### Introduction

Stock assessment is the quantitative evaluation of status of exploited living resources and predictions about their responses to alternative management strategies using various statistical models. The analytical models are usually preferred than biomass production models as they provide comparatively reasonable estimation of stock parameters. Similarly, depending upon the management objective (socio-economical, commercial, biological, recreational and conservational), several reference points are used for the management of fishery resources. However, the accuracy of any management reference point is influenced by the level of understanding about stock characteristics such as growth parameters, mortality parameters and spawning stock-recruitment relationships. The crucifix crab, *Charybdis feriata* (Linnaeus, 1758) (Family: Portunidae) is widely distributed throughout the Indo-pacific region (Apel and Spiridonov, 1998; Ng, 1998; Abello and Hispano, 2006). It is a sub-littoral species and found in a wide range of substrata ranging from muddy to rocky bottom (Ng, 1998; Yan, *et al.*, 2004) and is one of

the important edible marine crab resources of India. Crabs are not targeted resources but are mainly landed as bycatch at Veraval of single day and multiday trawl operations along north-west coast of India targeting shrimp and other demersal fishery resources. They are often subjected to sorting onboard and sizes found unsuitable for human consumption are usually discarded at sea or when landed as inedible bycatch, used for fish meal production.

Though there are studies on the fishery (Lalithadevi, 1985; Padayatti, 1990), morphometrics, food and feeding behaviour (Rameshbabu *et al.*, 2002a, b), reproductive biology (Padayatti, 1990; Rameshbabu *et al.*, 2006) and population dynamics (Dineshbabu, 2011) of this crab from different parts of India, there is no earlier report about this resource from Gujarat. Therefore, the present study was conducted to assess the fishery and stock of *C. feriata* along this coast and to judge the possible implication of uncertainty involved in decision making so that precautionary approach can be suggested for sustainable exploitation of the resource.

## Materials and methods

Data on catch, effort and size composition was collected fortnightly for a period of 3 years from January 2009 to December 2011 from commercial trawlers of Veraval Fishing Harbour, India (20°54' 0.50"N; 70°22' 9.60"E). Individual carapace width (CW) between tips of the longest lateral spines across the middle line between the frontal notch and the posterior margin was measured to the nearest mm using a digital caliper (Yamayo Measuring Tools Co. Ltd., Japan) and the individual body weight was recorded to the nearest gram using an electronic weighing machine (Scale-Tec™, India). A total of 1544 fresh specimens of *C. feriata* in the size range of 61-163 mm (CW) were collected randomly for analysis. The von Bertalanffy's growth parameters *viz.*, asymptotic length ( $L_{\infty}$ ) and growth co-efficient (K) were estimated using monthly raised data in the ELEFAN 1 module of FiSAT II (Version 1.2.2) (Gayanillo *et al.*, 2005). Age at length zero ( $t_0$ ) was calculated using formula:

$\log_{10}(-t_0) = -0.3922 - 0.2752 \log_{10} L_{\infty} - 1.038 \log_{10} K$  (Pauly, 1979). Growth performance index ( $\phi$ ) was calculated from formula:  $\phi = \log_{10} K + 2 \log_{10} L_{\infty}$  (Pauly and Munro, 1984). Longevity was estimated from the equation:  $t_{\max} = 3/K + t_0$  (Pauly, 1983a). The CW-weight relationship of *C. feriata* was established following the formula,  $W = aL^b$  (Le Cren, 1951) and similarity of regression lines were tested using analysis of covariance (ANCOVA) test.

The instantaneous total mortality rate (Z) was estimated by FiSAT II package using the length converted catch curve method (Pauly, 1983b). The natural mortality rate (M) was estimated using the formula:  $M = 1.535 K$  (Srinath, 1998), and the fishing mortality rate (F) was obtained as  $F = Z - M$ . The current exploitation rate ( $E_{\text{cur}}$ ) was calculated as  $E = F/Z$  (Ricker, 1975). Length structured virtual population analysis (VPA) of FiSAT II was used to obtain fishing mortalities per length class.

The midpoint of the smallest length group in the catch during the three-year period was taken as length at recruitment (Lr). Length at capture *i.e.*, length at which

50% of the crabs in the stock becomes vulnerable to the gear ( $L_{50}$ ) was estimated by probability of capture routine in the FiSAT-II package. For determining length at maturity ( $L_{M50}$ ), females of *C. feriata* (CW=67-163 mm, n=640) collected during January 2009 to December 2011 were used. Maturity stage of the females were determined by the size of ovary and were classified into 5 stages *viz.*, stage-1: immature, stage-2: early maturing, stage-3: late maturing, stage-4: Mature, and stage-5: Spent (Ryan, 1967). Proportions of mature female (*i.e.*, crabs which have completed stage-3) in sequential length classes (5 mm) were used for the logistic regression analysis following the method described by Ashton (1972):  $P = \frac{e^{a+b(CW)}}{1+e^{a+b(CW)}}$  where P = predicted mature proportion, a and b are the estimated coefficients of the logistic equation and CW = carapace width.  $L_{M50}$  was estimated as the negative ratio of the coefficients ( $-a/b$ ).

Since both male and female crabs inhabit the same fishing ground and are exposed to fishing at the same time, without discrimination, it is impracticable to manage the fishery separately and hence the sex pooled data was used for stock assessment study. The relative yield per recruit ( $Y'/R$ ) and relative biomass per recruit ( $B'/R$ ) at different fishing levels were estimated by FiSAT II package using relative yield per recruit analysis method (Beverton and Holt, 1966). The yield (Y), total biomass (B) and spawning stock biomass (SSB) at different fishing levels were predicted using length based Thompson and Bell bio-economic model (Thompson and Bell, 1934).

## Results and discussion

### Fishery

The annual catch and effort data of *C. feriata* during 2009-2011 is shown in Table 1. An average landing of 541 t per annum was harvested with an average catch rate (CPUE) of 11 kg per boat. During this period, though the contribution of *C. feriata* to the fishery was negligible, it constituted 44% of the annual average edible crab landing. An increase in the annual catch and catch rate of *C. feriata* was observed during the study period (2009-2011).

Table 1. Trawl effort and catch of *C. feriata* landed at Veraval during 2009 to 2011

Year	Efforts (boats)	Total trawl landing (t)	Total commercial crab landing (t)	<i>Charybdis feriata</i>			
				Catch (t)	CPUE (kg boat <sup>-1</sup> )	% to total crab landing	% to total trawl landing
2009	51551	129790	854	349	8	41	0.27
2010	51107	150115	1044	473	9	45	0.32
2011	53081	141433	1780	802	15	45	0.57
Average	51913	140446	1226	541	11	44	0.39

*Stock structure*

The stock structure (Fig. 1.) analysis indicates that samples obtained from the population slightly deviated from the normal distribution. Males constituted 44.75% (n=691) and females 55.25% (n=853) with an overall sex ratio (Male: Female) of 1.3. No significant difference was observed between the mean CW of male populations (109.26 mm) and female population (110.2 mm) (Independent t-test,  $p \leq 0.05$ ). Minimum and maximum CW were 61.0 mm and 161.0 mm in males and 67.0 and 163.0 mm in females. The mean CW of sex pooled data was found to be 109.8 mm. The annual length frequency distribution revealed that the abundant size group in the crab stock mainly consisted of individuals of 110.0 mm to 115.0 mm CW (mode CW = 112.0 mm).

*Carapace width (CW)-weight relationship*

The relationship between CW-body weight for the sex pooled data (Fig. 2.), showed that growth was isometric. The relationship between carapace width and body weight for male and female was derived as:

Male:  $W = 0.0001 CW^{2.94}$  ( $r^2 = 0.93$ ,  $n = 544$ )  
 Female:  $W = 0.00007 CW^{2.97}$  ( $r^2 = 0.96$ ,  $n = 508$ )

where, W is weight in g and CW is carapace width in mm. No significant difference was observed between




Fig. 2. Carapace width and weight relationship of *C. feriata* (sex pooled data) from Veraval waters

the slopes of the regression lines ( $p \geq 0.05$ ) whereas, the intercept of male was found to be significantly higher than the female ( $p \leq 0.05$ ). This suggests that in the present population, the males are heavier than female of similar size. This concurs with the earlier findings where males were found to be heavier than females in *C. feriata* (Dineshbabu, 2011) and other portunid crabs (Laithadevi, 1985, Sukumaran and Neelakantan, 1997; Dineshbabu *et al.*, 2007). This difference may be due to the variation in cheliped size, foraging behaviour and metabolic rate between the sexes (Miyasaka *et al.*, 2007; Thirunavukkarasu and Shanmugam, 2011).




Fig. 1. Population structure of *C. feriata* at Veraval during 2009 and 2011

*Growth parameters*

The estimated growth parameters derived using male, female and sex pooled data is shown in Table 2. The growth estimated by von Bertalanffy's growth equation showed that the male crab attained 113 mm (CW) in 1<sup>st</sup> year, 149 mm (CW) in 2<sup>nd</sup> year and 163 mm (CW) in 3<sup>rd</sup> year, whereas, the female crab reached 111 mm (CW) in 1<sup>st</sup> year, 150 mm (CW) in 2<sup>nd</sup> year and 165 mm (CW) in 3<sup>rd</sup> year. The restructured length frequency distribution and growth parameters of sex pooled data is shown in the Fig. 3. The  $L_{\infty}$  (175.5 mm for female and 171 mm for male) and K (0.91 yr<sup>-1</sup> for female and 0.98 yr<sup>-1</sup>) of *C. feriata* obtained in the present study can be compared with the earlier reported  $L_{\infty}$  (164 mm for female and 173 mm for male) and K (0.89 yr<sup>-1</sup> for female and 0.84 male) obtained from Karnataka waters (Dineshbabu *et al.*, 2011). The estimation of growth parameters by ELEFAN is greatly influenced by  $L_{max}$  (maximum length in sample) and

corresponds to the size ( $L_{max}$  which is approximately 95% of  $L_{\infty}$ ) at which 99% individuals in an unexploited stock dies (Alagaraja, 1989). In the present study, the longevity ( $t_{max}$ ) of the species was estimated to be 3 year.

*Mortality, exploitation and virtual population analysis*

The natural mortality rate (M), fishing mortality rate (F) and the total mortality rate (Z) was found to be higher in males compared to female whereas, the current exploitation rate ( $E_{cur}$ ) was same for both the sexes (Table 2). The M, Z, F and  $E_{cur}$  obtained using the sex pooled data is shown in Fig. 4. The estimated natural mortality rate (M) is important to understand rate of stock decay. But 'M' is a difficult parameter to estimate in an exploited resource (FAO, 1993). The 'F' (2.48 yr<sup>-1</sup>) observed in the present study, was higher than 'M' (1.49 yr<sup>-1</sup>) as a result of which  $E_{cur}$  (0.62) was found to be higher than the optimum exploitation rate of 0.5 (Gulland, 1971).

Table 2. Growth parameters of *C. feriata* from Veraval waters during 2009 and 2011

Sex	von Bertalanffy's growth parameters			Growth performance index	Mortality rates			Exploitation rate
	$L_{\infty}$	K	$t_0$		$\phi$	M	Z	
Male	171.0	0.98	-0.10	2.46	1.50	4.19	2.69	0.64
Female	175.5	0.91	-0.10	2.45	1.40	3.92	2.52	0.64
Pooled	171.5	0.97	-0.10	2.46	1.49	3.97	2.48	0.62




Fig. 3. Restructured growth curve of *C. feriata* from Veraval waters

hence varies from location to location. However, the growth performance index ( $\phi$ ) (female: 2.45 and male: 2.46) obtained in the present study was found to be in the reasonable range within the family ( $2 < \phi < 3$ ) (Pauly and Munro, 1984). The growth rate (K) of male in the present study was found to be comparatively higher than female. As males are not physiologically associated with egg production and incubation, they save energy for somatic growth and show higher growth rate (Hartnoll, 1982). The longevity ( $t_{max}$ ) of a species

Virtual population analysis showed that F changes with crab size (CW) which may be attributed to the variation in selectivity of the fishing gear. F was low ( $< 0.1$ ) for crabs of 65-70 mm CW and it crossed 1.0 when crabs attained CW of 105 mm. However, at  $LM_{50}$  (87.3 mm) the F was below 0.5 and exceeded M




Fig. 4. Estimation of Z of *C. feriata* from Length converted catch curve method (sex pooled data)

when the crabs attained CW of 110 mm. Maximum fishing mortality (2.5) was observed at CW of 120-125 mm (Fig. 5).




Fig. 5. Length structured VPA for *C. feriata* from Veraval waters

*Gear selectivity and maturity*

Length at recruitment ( $L_r$ ) for *C. feriata* was found to be 62.5 mm. Logistic regression of the probability of capture for sequential length classes obtained from length converted catch curve analysis using trawl type selection revealed that 50% of the crabs in stock ( $L_{50}$ ) became vulnerable to gear when they attained CW of 95.4 mm. The  $L_{25}$  and  $L_{75}$  were 90.4 mm and 100.3 mm, respectively (Fig. 6). Mature and berried females were recorded throughout the entire fishing period. Maturity study revealed that, 50% of the crabs in stock ( $L_{m50}$ ) were mature at CW of 87.3 mm (Fig. 7). In the present study, the  $L_{50}$  is found to be higher than  $L_{m50}$  which indicates that crabs enter into peak exploitation phase after attaining sexual maturity which is desirable for the regeneration of




Fig. 6. Length at capture ( $L_{50}$ ) of *C. feriata* from Veraval waters




Fig. 7. Size at maturity ( $L_{m50}$ ) of *C. feriata* from Veraval waters

the stock. The  $L_{m50}$  in the present study is found to be higher than the earlier reported 70 mm from Karnataka waters (Dineshbabu, 2011). Maturity of brachyuran crabs is influenced by environmental factors such as water temperature and salinity that varies from location to location and therefore, shows geographical variations (Hines, 1989; Fisher, 1999). However, the uncertainty factor associated with  $L_{50}$  and  $L_{m50}$  should not be ignored as the analysis does not give any weightage for the smaller sized crabs missing in the samples, which are often discarded at sea.

*Stock assessment*

The relative Y/R and B/R of *C. feriata* were estimated using selection ogive procedure of FiSAT II (Fig. 8). “Selection ogive” was used for the analysis as the “knife edge selection” could lead to substantial bias for the short lived, fast growing tropical species (Pauly and Soriano, 1986; Silvestre, *et al.*, 1991). Moreover, the assumptions




Fig. 8. Relative yield per recruit analysis of *C. feriata* from Veraval waters

in the “knife edge selection” are rarely met in the natural condition.  $L_{50}/L_{\infty}$  as 0.556 and  $M/K$  as 1.535 were used as the input parameters for the analysis. The analysis indicated that, the exploitation rate which maximises yield per recruit ( $E_{\max}$ ) was 0.75. In case of r-selected species maximum yield is achieved at higher rate of exploitation compared to the k-selected species which suggests that r-selected species can sustain higher fishing mortalities (Silvestre, *et al.*, 2004; Clarke, 2005). Since, *C. feriata* is a small, fast growing tropical crab species having higher natural mortality, it can be characterised as r-selected species. But there are certain issues in using  $E_{\max}$  as a management reference point because it tends to be unrealistically high for small tropical species with high natural mortality (Gulland, 1983; Pauly, 1984). Moreover,  $E_{\max}$  usually corresponds to a very low level of biomass. Therefore, it is often recommended to reduce exploitation to a precautionary safe level *i.e.*,  $E_{0.1}$  (an effort level at which the marginal increase in yield per recruit reaches 1/10 of the marginal increase computed at a very low value of  $E$ ).

Both  $E_{\max}$  and  $E_{0.1}$  are often used as proxies for maximum sustainable yield (MSY) and maximum economic yield (MEY), and set as management reference points (Jakubaviciute, *et al.*, 2011). In the present study,  $E_{0.1}$  obtained is equal to  $E_{\text{cur}}$  which indicates that the fishery is running at economic reference point (MEY). Another reference point used for fisheries management is  $E_{0.5}$  which corresponds to exploitation rate at which the B/R reduces by 50% compared to virgin stock *i.e.*, to the level which theoretically maximises surplus production (Pauly, 1984).  $E_{0.5}$  is also used as a proxy for the optimum sustainable yield (OSY) (Dadzie *et al.*, 2005). In the present study, the optimum biomass of *C. feriata* can be maintained at  $E_{0.5}$  of 0.38 whereas, the  $E_{\text{cur}}$  (0.62) is high and reducing the B/R considerably. Since, B/R and  $E$  can be used as proxies for catch rate (CPUE) and effort (F) respectively, it shows that the catch rate decreases with the increase in effort. But this decrease in CPUE should not be necessarily treated as growth overfishing which occurs only at very high fishing effort (usually above  $F_{\text{MSY}}$ ), where growth cannot balance the death process (Sparre and Venema, 1998). Moreover, as far as recruitment overfishing is concerned, it is necessary to understand the relationship between spawning stock biomass (SSB) and recruitment. Since this relationship in the present crab fishery is unknown and we are assuming a constant recruitment over the period in Y/R analysis, it will be difficult to comment

over recruitment over fishing. In such situations, where stock-recruitment relationship cannot be established (Restrepo and Powers, 1999) or the recruitment is independent of stock size (Myers, 2001), reference points such as  $F_{\text{MAX}}$  (fishing effort that gives maximum yield per recruit) or  $F_{0.1}$  (10% gradient in yield per recruit curve) are often recommended.

Spawning stock biomass (SSB) can be used as recruitment fishing reference point which is essential for maintaining the recruitment in the future (Clark, 1991). SSB should be maintained at 20% for stocks having average resilience and at 30% for poorly understood stocks (Mace and Sissenwine, 1993). Therefore, in case of stocks where the relationship between spawning biomass and recruitment cannot be established, it is advisable to use precautionary management reference point such as SSB to prevent recruitment overfishing (Rosenberg and Repestro, 1996). In the present study, Thompson and Bell bio-economic analysis showed that the present fishing level has already achieved the MEY while maintaining the SSB at about 36% of  $\text{SSB}_0$  (Fig. 9).

In this scenario if the management targets economic objective, exploitation can be allowed at the present fishing level. Since the stock recruitment relationship is not clearly defined, as a precautionary approach, SSB at about 30% of  $\text{SSB}_0$  should be considered as a biologically safe management reference point. Considering this, fishing level (F-factor) can be increased by 1.3 times the present level without jeopardising the biological management objective, but this will marginally decrease MEY. The analysis also showed that yield can be increased from 464.4 t (current equilibrium yield) to 513.5 t (MSY) by increasing fishing level (F-factor) by 2.4 times the present value. However, the estimated  $F_{\text{MSY}}$  would deplete the SSB to 20% of virgin spawning stock biomass ( $\text{SSB}_0$ ) and economic yield by 14% of MEY. Poorly understood stock gives rise to uncertainty in the estimation of stock parameters which affects the accuracy of management reference point. For example, maximum sustainable yield (MSY), though has been proposed as a universal objective for the fisheries management, it is not widely adopted as it does not give an accurate estimation of status for many stocks, especially where stock-recruitment relationship is poorly established (Edwards *et al.*, 2012). Therefore, in the event of uncertainty, precautionary approaches should be taken in to account for management decision making (FAO, 1996). Since *C. feriata* is a non-targeted resource in this region, consideration should also be given to the




Fig. 9. Length based Thompson and Bell analysis of *C. feriata* from Veraval waters

demand and exploitation status of other targeted fishery resources while arriving at suitable management reference points.

### Acknowledgements

The authors express sincere gratitude to Dr. G. Syda Rao, Former Director, CMFRI, Cochin for providing facilities and encouragement to carry out the above research work. The authors are also thankful to Dr. G. Maheswarudu, Head, Crustacean Fisheries Division, CMFRI, Cochin and Dr. Subhadeep Ghosh., Senior Scientist, Regional centre of CMFRI, Visakhapatnam for constructive criticism and comments during the preparation of this article.

### References

- Abello, P. and Hispano, C. 2006. The capture of the Indo-Pacific crab *Charybdis feriata* (Linnaeus, 1758) (Brachyura: Portunidae) in the Mediterranean Sea. *Aquatic Invasions*, 1(1): 13-16.
- Alagaraja, K. 1989. Letter to the Editor. *Fishbyte*, 7(3): 2-3.
- Apel, M. and Spiridonov, V. A. 1998. Taxonomy and zoogeography of the portunid crabs (Crustacea: Decapoda: Brachyura: Portunidae) of the Arabian Gulf and adjacent waters. *Fauna of Arabia*, 17: 159-331.
- Ashton, W. D. 1972. *The logit transformation with special reference to its uses in bioassay*. Haffner Publishing Co., Inc., New York, USA, 88 pp.
- Beverton, R. J. H. and Holt, S. J. 1966. Manuals of methods for fish stock assessment: Part II, Tables of yield function, *Fisheries Biology Technical Paper*, 38(4), Ver. 1, FAO, Rome, Italy, 67 pp.
- Clark, W. G. 1991. Groundfish exploitation rates based on life history parameters. *Can. J. Fish. Aqua. Sci.*, 48: 734-750.
- Clarke, M. 2005. A life history approach to the assessment of deepwater fisheries in the North-east Atlantic. Deep Sea 2003: Conference on the Governance and Management of Deep sea Fisheries, Part 1: Conference reports. *FAO Fisheries Proceedings, No. 3/1*. Food and Agriculture Organization of the United Nations, Queenstown, New Zealand, p. 196-210.
- Dadzie, S., Manyala, J. O. and Abou-Seedo, F. 2005. Aspects of the population dynamics of *Liza klunzingeri* in the Kuwait Bay. *Cybum*, 29(1): 13-20.
- Dineshbabu, A. P. 2011. Biology and exploitation of the crucifix crab, *Charybdis feriata* (Linnaeus, 1758) (Brachyura: Portunidae) from Karnataka coast. *Indian J. Fish.*, 58(1): 25-29.
- Dineshbabu, A. P., Sreedhara, B. and Muniyappa, Y. 2007. Fishery and stock assessment of *Portunus sanguinolentus* (Herbst) from south Karnataka coast. *J. Mar. Biol. Ass. India*, 49(2): 134-140.
- Edwards, C. T. T., Hillary, R. M., Levontin, P., Blanchard, J. L. and Lorenzen, K. 2012. Fisheries assessment and management: A synthesis of common approaches with special reference to deepwater and data-poor stocks. *Rev. Fish. Sci.*, 20(3): 136-153.
- FAO 1993. Fisheries management in south-east Lake Malawi, the Upper Shire River and Lake Malombe, with particular reference to the fisheries on chambo (*Oreochromis* spp.). *CIFA Technical Paper no. 21*, FAO, Rome, Italy, 113 pp.

- FAO 1996. *Technical guidelines for responsible fisheries*, Ver. 2. FAO, Rome, Italy, 54 pp.
- Fisher, M. R. 1999. Effect of temperature and salinity on size at maturity of female blue crabs. *Trans. Am. Fish. Soc.*, 128(3): 499-506.
- Gayanillo, Jr. F. C., Sparre, P. and Pauly, D. 2005. *FAO ICLARM Stock assessment tools II (FiSAT II). User's guide. Computerized information Series (Fisheries)*. No. 8, revised version, FAO, Roma, Italy, 126 pp.
- Gulland, J. A. 1971. *The fish resources of the ocean*. Fishing News books Ltd., Farnham, England, 255 pp.
- Gulland, J. A. 1983. *Fish stock assessment - A manual of basic methods*. Wiley Interscience, FAO Wiley series on food and agriculture. Vol. 1, Chichester, U. K., 223 pp.
- Hartnoll, R. G. 1982. Growth. In: Abele, L. G. (Eds.), *The biology of crustacea 2 - Embryology, morphology and genetics*, Academic Press, New York, p. 11 – 196.
- Hines, A. H. 1989. Geographic variation in size at maturity in Brachyuran crabs. *Bull. Mar. Sci.*, 45(2): 356-368.
- Jakubaviciute, E., Putys, Z., Dainys, J. and Lozys, L. 2011. Perch (*Perca fluviatilis*) growth, mortality and stock exploitation by 40-45 mm mesh sized gillnet fishery in the Curonian Lagoon. *Acta Zoologica Lithuana*, 21(3): 215-220.
- Lathadevi, S. 1985. The fishery and biology of crabs of Kakinada region. *Indian J. Fish.*, 32(1): 18-31.
- Le Cren, E. D. 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20(2): 201-219.
- Mace, P. M. and Sissenwine, M. P. 1993. How much spawning per recruit is enough? In: Smith, S. J., Hunt, J. J. and Revered, D. (Eds.), *Risk evaluation and biological reference points for fisheries management*. Canadian Special Publication of Fisheries and Aquatic Sciences, National Research Council of Canada, p. 101-118.
- Miyasaka, H., Genkai-Kato, M., Goda, Y. and Omori, K. 2007. Length-weight relationships of two varunid crab species, *Helice tridens* and *Chasmagnathus convexus* in Japan. *Limnol.*, 8(1): 81-83.
- Myers, R. A. 2001. Stock and recruitment: generalizations about maximum reproductive rate, density dependence, and variability using meta-analytic approaches. *ICES J. Mar. Sci.*, 58: 937-951.
- Ng, P. K. L. 1998. Crabs. In: Carpenter, K. E. and Niem, V. H. (Eds.), *The living marine resources of the Western Central Pacific*, Vol. 2. Food and Agriculture Organization of the United Nations, Rome, Italy, p. 1046-1155.
- Padayatti, P. S. 1990. Notes on population characteristics and reproductive biology of the portunid crab *Charybdis (Charybdis) feriatus* (Linnaeus) at Cochin. *Indian J. Fish.*, 37 (2): 155-158.
- Pauly, D. 1979. Theory and management of tropical multi-species stocks. A review with emphasis on the south-east Asian demersal fisheries. *ICLARM Stud. Rev.*, 1: 35 pp.
- Pauly, D. 1983a. Some simple methods for the assessment of tropical fish stocks. *FAO Fisheries Technical Paper*, No. 243, Rome, Italy, 52 pp.
- Pauly, D. 1983b. Length converted catch curves. A powerful tool for fisheries research in tropics (Part-I). *ICLARM, Fishbyte*, 1(2): 9-13.
- Pauly, D. 1984. Fish population dynamics in tropical waters: a manual for use with programmable calculators. *ICLARM Stud. Rev.*, 8: 325 pp.
- Pauly, D. and Munro, J. L. 1984. Once more, on the composition of growth in fish and invertebrates. *Fishbyte*, 2(1): 21.
- Pauly, D. and Soriano, M. L. 1986. Some practical extensions to Beverton and Holt's Relative Yield-Per-Recruit Model. In: Maclean, J. L., Dizon, L. B. and Hosillo, L. V. (Eds.), *The First Asian Fisheries Forum*, Asian Fisheries Society, Manila, Philippines, p. 491-496.
- Rameshbabu, K. V., Benkappa, S. and Chandra Mohan, K. 2002a. Food and feeding habits of *Charybdis (Charybdis) feriatus* (Linnaeus) from Mangalore region, *Indian Hydrobiol.*, 5(1): 1-7.
- Rameshbabu, K. V., Benkappa, S. and Chandra Mohan, K. 2002b. Length-weight relationship in *Charybdis (Charybdis) feriatus* (Linnaeus) from Mangalore region. *Indian Hydrobiol.*, 5(1): 9-14.
- Rameshbabu, K. V., Benkappa, S. and Chandra Mohan, K. 2006. Breeding biology of *Charybdis (Charybdis) feriatus* (Linnaeus) from Mangalore. *Indian J. Fish.*, 53(2): 182-184.
- Restrepo, V. R. and Powers, J. E. 1999. Precautionary control rules in US fisheries management: specification and performance. *ICES J. Mar. Sci.*, 56: 846-852.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Board Canada*, No. 191, 382 pp.
- Rosenberg, A. A. and Repestro, V. R. 1996. Precautionary management reference points and management strategies, Precautionary approach to fisheries, Part 2, *Fisheries Technical Paper*, No. 350, FAO, Rome, Italy, p. 129-140.
- Ryan, E. P. 1967. Structure and function of the reproductive system of the crab *Portunus sanguinolentus* (Herbst) (Brachura, Portunidae) II. The female system. *Proceedings of the symposium of the Marine Biological Association of India*, Part II, Marine Biological Association of India, Ernakulam, India, p. 522-544.
- Silvestre, G. T. and Garces, L. R. 2004. Population parameters and exploitation rate of demersal fishes in Brunei Darussalam (1989-1990). *Fish. Res.*, 69 (1): 73-90.

- Silvestre, G. T., Soriano, M. and Pauly, D. 1991. Sigmoid selection and the Beverton and Holt yield equation. *Asian Fish. Sci.*, 4(1): 85-98.
- Sparre, P. and Venema, S. C. 1998. Introduction to tropical fish stock assessment (Part-1 manual), *Fisheries Technical Paper*, No. 306 (1), Rev. 2, FAO, Rome, Italy, 407 pp.
- Srinath, M. 1998. Empirical relationships to estimate the instantaneous rate of natural mortality. *Indian J. Fish.*, 45(1): 7-11.
- Sukumaran, K. K. and Neelakantan, B. 1997. Length-weight relationship of two marine crabs, *Portunus sanguinolentus* (Herbst) and *Portunus pelagicus* (Linnaeus). *Indian J. Mar. Sci.*, 26: 39-42.
- Thirunavukkarasu, N. and Shanmugam, A. 2011. Length-weight and width-weight relationships of mud crab *Scylla tranquebarica* (Fabricius, 1798). *Euro. J. Appl. Sci.*, 3(2): 67-70.
- Thompson, W. F. and Bell, F. H. 1934. Biological statistics of the Pacific halibut fishery. Effect of changes in intensity upon total yield and yield per unit of gear. *Report of the International Fisheries (Pacific halibut) Commission*, 8: 49 pp.
- Yan, Y., Huang, L. and Miao, S. 2004. Occurrence of the epizoic barnacle *Octolasmis angulata* on the crab *Charybdis feriatus* from Daya Bay, China. *J. Mar. Biol. Ass. U.K.*, 84: 619-620.

Date of Receipt : 31.01.2013

Date of Acceptance : 03.06.2014