

# ISOLATION AND CHARACTERIZATION OF *Vibrio* spp. FROM SEA FOOD AND ENVIRONMENTAL SAMPLES IN AND AROUND CHENNAI CITY

S.J. Deepak<sup>1</sup>, K. Porteen\*<sup>2</sup>, A. Elango<sup>3</sup>, T.M.A. Senthilkumar<sup>4</sup>,  
R. Narendra Babu<sup>5</sup> and S. Sureshkannan<sup>6</sup>

Department of Veterinary Public Health and Epidemiology  
Madras Veterinary College  
Tamil Nadu Veterinary and Animal Sciences University  
Chennai, Tamil Nadu, India

## ABSTRACT

*Vibrio* is a major sea food pathogen which needs to be monitored regularly for devising appropriate control strategy to ensure food safety among sea food consumers and handlers. The present study aims to ascertain the occurrence of *Vibrio* spp. in seafood viz fishes, prawn/shrimps, crabs and water samples collected from various fish markets in and around Chennai city, Tamil Nadu, India. A total of 175 samples were screened, of which 106 (60.57%) samples were found positive by PCR targeting the 16s rRNA of *Vibrio* genus and all the positive isolates were subjected to *toxR* gene identification by Multiplex PCR for species level identification. At the species level 28(26.41%) were positive for *Vibrio parahemolyticus* (Vp), 6 (5.67%) positive for *Vibrio Cholerae* (Vc), 2 (1.89%) positive for *Vibrio vulnificus* (Vv). The presence of mixed vibrio species were also observed of which 56 samples (43.40%) revealed the presence of Vp and Vc, 7 (6.60%) with Vp and Vv, 3 (2.84%) with Vv and Vc and 14 samples with (13.20%) Vp, Vv and Vc. The present study identified the presence along with mixed *Vibrio* spp. in most of the samples tested from the study area.

**Key words:** Crabs, Fish, Seafood, *Vibrio cholera*, *Vibrio parahemolyticus*, *Vibrio vulnificus*

## INTRODUCTION

*Vibrio* is the genus of Gram-negative rod-shaped bacteria which are catalase

and oxidase positive, motile with bipolar flagella, sensitive to the vibriostatic agent (O/129), require sodium chloride for growth and widespread in the coastal and estuarine ecosystems (Baumann and Schubert, 1984; Farmer III *et al.*, 2005).

This bacterial group has the capacity to cause serious alimentary toxico-infection and watery diarrhoea and is implicated to the consumption of raw or undercooked contaminated fish or shellfish. Pathogenic *Vibrio* species pose a considerable public

Corresponding author, email: rajavet2002@gmail.com

<sup>1</sup>Ph.D Scholar

<sup>2</sup>Assistant Professor

<sup>3</sup>Dean-in-charge, Veterinary College and Research Institute, Tamil Nadu Veterinary and Animal Sciences University, Salem, Tamil Nadu, India

<sup>4</sup>Project Director, Translational Research Platform for Veterinary Biologicals (TRPVB), Tamil Nadu Veterinary and Animal Sciences University, Madhavaram Milk Colony, Chennai, Tamil Nadu, India

<sup>5</sup>Professor and Head, Department of Livestock Products Technology (Meat Science)

<sup>6</sup>Professor and Head

health threat as they result in sporadic and epidemic human infections, and hence represents an important microbial group in the field of food safety and quality.

Major vibrios that cause sea foodborne ichthyozoonoses are *V. cholerae*, *V. mimicus*, *V. parahaemolyticus* and *V. vulnificus* (Oliver, 2006). *Vibrio cholerae* is an important pathogenic *Vibrio* with the serogroups O1 and O139 to be specifically considered. The major virulence factor of *V. cholerae* is the CTX cholera toxin. *Vibrio parahaemolyticus* is a halophilic bacterium which causes acute gastroenteritis on consumption of raw contaminated sea foods and naturally inhabits marine waters. *V. parahaemolyticus* has two major species-specific pathogenic markers the *tdh* that encodes the thermostable direct hemolysin (TDH) which can lyse red blood cells and the *trh* encoding *tdh* related hemolysin (TRH). *Vibrio vulnificus* is a pathogenic species that inhabits in seawater and causes gastroenteritis or septicemia and can result in severe infections and high mortality in immune compromised patients with chronic liver disease.

The phenotypic identification of *Vibrio* species is often problematic as there are only a few common characteristics and owing to the intra-species variation in biochemical characteristics. The application of molecular methods such as PCR may enable easy and rapid identification of the species. The *toxR* gene that encodes the trans-membrane proteins involved in the regulation of virulence-associated genes is an ideal target for a simple multiplex PCR, because it appears to be well conserved

among the bacteria within *Vibrionaceae*, but sufficiently divergent to distinguish between different *Vibrio* spp (Kim *et al.*, 1999; Osorio and Klose, 2000). With this background the present study was envisaged to study the occurrence of *Vibrio* spp. in different sea food and environmental samples collected in and around Chennai city, Tamil Nadu.

## MATERIAL AND METHODS

### Sample collection

A total of 175 sea food and environmental samples were collected from various fish markets in and around Chennai city, Tamil Nadu, India. The sample includes 50 fishes, 60 prawn/shrimps, 30 crabs, 5 shells and 30 water samples.

### Bacterial culture, isolation and identification

The samples were enriched in alkaline peptone water (APW; Himedia, India) at pH 8.5 at 37°C for 6 hours. The enriched samples were plated on selective media thiosulfate citrate bile salts sucrose agar (TCBS; Himedia, India) containing 1 per cent sodium chloride (NaCl), and incubated at 37°C for 18 to 24 hour. The presumptive *V. parahaemolyticus* strains (green with blue tinge on TCBS Agar), presumptive *V. cholerae* strains (yellow on TCBS Agar), and presumptive *V. vulnificus* strains (dark green/yellow on TCBS Agar), were purified on Nutrient agar (NA; Himedia, India) containing 3 per cent NaCl (BAM Protocol, 2014). The presumptive colonies were confirmed by gram staining and biochemical tests.

## DNA preparation

A few presumptive colonies from nutrient agar were transferred to 100 µL of sterile water. The bacterial cells were lysed by boiling at 100°C for 5 min, followed by immediate chilling for 2-3 min at -20°C and then centrifuged at 5000 x g for 15 min at 4°C. The recovered supernatant was used as template in the PCR assay.

## Polymerase chain reaction

The polymerase chain reaction was performed with primers targeting the 16S rRNA for identification of at genus level. Further, multiplex-PCR was performed with primers targeting the *toxR* gene for the identification of *V. parahaemolyticus*, *V. cholera* and *V. vulnificus*. The primer sequences used and the thermal cyclic conditions are provided in Table 1. The PCR

amplicons were electrophoresed in a 1.5 percent agarose gel, stained using ethidium bromide and the results were visualized the Gel documentation system (Bio Rad, India).

## RESULTS

### Isolation and identification of *Vibrio* spp. based on cultural characteristics and PCR

The enrichment with alkaline peptone water for sea food samples was necessary for rejuvenation and enrichment of the *Vibrio* spp. The thiosulphate citrate bile salt sucrose agar (TCBS) is the most widely used media to isolate *Vibrio* spp. from sea food and environmental samples. The colony morphology of the *Vibrio* spp. is as follows: green with blue tinge colony for *Vibrio parahaemolyticus*, yellow color colony for *Vibrio cholera* and dark green/yellow for *Vibrio vulnificus*.

**Table 1. List of primers used in the present study**

Name	Sequence	Product length (bp)	Annealing temperature	Reference
V16S rRNA-F	CGG TGA AAT GCG TAG AGA T	663	57°C	Sudha et al., 2014
V16S rRNA-R	TTA CTA GCG ATT CCG AGT TC			
U-toxR-F	GAS TTT GTT TGG CGY GAR CAA GGT T	Vp-297 Vc-640 vv-435	55°C	Bauer and Rørvik, 2007
vp-toxR-R	GGT TCA ACG ATT GCG TCA GAA G			
vc-toxR-R	GGT TAG CAA CGA TGC GTA AG			
vv-toxR-R	AAC GGA ACT TAG ACT CCG AC			

Out of 175 samples screened in the present study we found that 106 samples were positive for the 16S rRNA specific simplex PCR confirming the genus level identification of *Vibrio* (Fig. 1). The multiplex PCR for the *toxR* gene enabled

differentiation of *Vibrio* species as *Vibrio parahaemolyticus* (297bp), *Vibrio cholera* (640bp) and *Vibrio vulnificus* (435bp). Among the 106 *Vibrio* positive samples we found that 28 (26.41%) positive for *Vibrio parahaemolyticus* (Vp), 6 (5.67%)

positive for *Vibrio cholera* (Vc) and 2 (1.89%) positive for *Vibrio vulnificus* (Vv). The mixed infections were observed in 56 samples (43.40%) with Vp and Vc, 7 (6.60%) with Vp and Vv, 3 (2.84%) with Vv and Vc and 14 (13.20%) with Vp, Vv and Vc (Fig. 2 and Table 2).


## DISCUSSION

The presence of *Vibrio* spp. in sea food is highly variable based on different

seafood samples, time and place of sample collection. From the present study, it is evident that the use of salt at 1-3 per cent is beneficial in enriching the growth of *Vibrio* in laboratory cultural techniques. The prevalence of *Vibrio* spp. in sea food in present study as assessed by the cultural technique was found to be 60.57 per cent (106/175) and confirmation by PCR targeting the 16s rRNA gene, concurred with the report of Azwai *et al.*, (2016), which reported an incidence of *Vibrio* spp.


**Table 2. Prevalence of *Vibrio* spp. indifferent seafood samples**

Organism	Water	Shrimp	Fish	Crab	Total
<i>Vibrio parahaemolyticus</i>	4	8	8	8	28
<i>Vibrio cholerae</i>	0	2	2	2	6
<i>Vibrio vulnificus</i>	0	2	0	0	2
Vp+Vc	8	12	22	14	56
Vp+Vv	0	5	1	1	7
Vv+Vc	0	3	0	0	3
Vp+Vv+Vp	7	7	0	0	14



**Fig. 1. PCR gel image depicting the amplification 16s rRNA gene (663bp)**

[L: DNA Ladder (Genederix 1kb), NC: Negative control, 1-6: sea food samples]



**Fig. 2. PCR gel image depicting the amplification of toxR gene by Multiplex PCR**

[L: DNA Ladder (Genederix 100bp), NC: Negative control, 1,3,4: Vp-297bp+Vc-640bp, 2:Vv-435bp+Vc-640bp amplified from sea food samples]

of 51.6 per cent in seafood, meat and meat products in Libya. Similarly, Alaboudi *et al.*, (2016) detected 66% positivity of typical *Vibrio* isolates from 200 sea food samples. In Indian sub-continent, a study by Sudha *et al.*, (2014), found that 65.5 per cent of the 110 samples screened were reported to be positive for *Vibrio* spp. The *Vibrio* spp. was the major pathogen from seafood to cause diseases in humans and, for this reason, it may be considered as an agent of zoonoses (Austin, 2010).

In the present study, out of 106 positive samples screened for species level identification of *Vibrio* spp. using toxR gene by multiplex PCR was applied by Bauer and Rørvik, (2007) and reported its utility in the reliable identification of *V. parahaemolyticus*, *V. vulnificus* and *V. cholerae*. In the present study, there was a higher frequency of *Vibrio parahaemolyticus* which concurred with results of Yu *et al.*, 2016 who have also reported that *V. parahaemolyticus* in shellfish to be 34.3% (48/140). Xu *et al.*, (2016) also reported that 36.2 % of the samples to be positive in sea food. In India, a study by Sudha *et al.*, (2014), 68.1 per cent samples were found to be positive for *V. parahaemolyticus* from seafoods. The occurrence of *Vibrio cholera* from seafoods was earlier documented by Kumar and Lalitha, (2013) who reported that 39per cent positivity. A study by Hill *et al.*, (2011) and Senderovich *et al.*, (2010) concluded that fish species harvested from Indian coastal water bodies had higher number of *Vibrio* spp. and indicated the risk of this pathogen to consumers.

## CONCLUSION

The occurrence of *Vibrio* spp. in seafood samples may be due to environmental stressors, improper handling and storage and cross contamination of sea food. Hence there is a need for strict hygienic measures to be adopted at various levels of production till it reaches the consumers to ensure safety among sea food consumers and handlers.

## ACKNOWLEDGEMENT

We would thank the support and financial aid provided by Dean, Madras Veterinary College, TANUVAS and Professor and Head, Department of Veterinary Public Health and Epidemiology to carry out special problem.

## REFERENCES

- Akther, F., Neogi, S.B., Chowdhury, W.B., Sadique, A., Islama, A., Akhter, M.Z., Fatema-Tuz, J., Ohnishi, M., Watanabe, H., Boucher, Y. and Alam, M. (2016). Major tdh+ *Vibrio parahaemolyticus* serotype changes temporally in the Bay of Bengal estuary of Bangladesh. *Infection, Genetics and Evolution*, **41**: 153–159.
- Alaboudi, A.R., Ababneh, M., Osaili, T. M. and Al Shloul, K. (2016). Detection, identification, and prevalence of pathogenic *Vibrio parahaemolyticus* in fish and coastal environment in Jordan. *Journal of food science*, **81**(1): 130–134.
- Austin, B. (2010). Vibrios as causal agents of zoonoses. *Veterinary Microbiology*, **140**: 310-317.

- Azwai, S.M., Alfallani, E.A., Abolghait, S.K., Garbaj, A.M., Naas, H.T., Moawad, A.A., Gammoudi, F.T., Rayes, H.M., Barbieri, I. and Eldaghayes, I.M. (2016). Isolation and molecular identification of *Vibrio* spp. by sequencing of 16S rDNA from seafood, meat and meat products in Libya. *Open Veterinary Journal*, **6**(1): 36-43.
- BAM Protocol, (2014) Bacteriological Analytical Manual (BAM), Chapter 9: *Vibrio*. Laboratory Methods (Food), Food and Drug Administration, Eds: Kaysner, C.A., Depaola, A. Jr. and Jones, J. U.S. Food & Drug administration, Accessed: <https://www.fda.gov/food/laboratory-methods-food/bam-chapter-9-vibrio>
- Bauer, A. and Rørvik. L.M. (2007). A novel multiplex PCR for the identification of *Vibrio parahaemolyticus*, *Vibrio cholerae* and *Vibrio vulnificus*. *Letters to Applied Microbiology*, **45**: 371–375.
- Baumann, P. and Schubert. R.H.W. (1984). Section 5. Facultative anaerobic Gram negative rods. Family II Vibrionaceae. In *Bergey's Manual of Systematic Bacteriology* ed Kried, N.R. and Holt, J.G. pp. 516–550 Baltimore/London: Williams & Wilkins.
- Farmer III, J.J., Janda, M., Brenner, F.W., Cameron, D.N. and Birkhead, K.M. (2005) Genus I. *Vibrio* Pacini 1854, 411AL. In: Brenner, D.J., Krieg, N.R., Staley, J.T. (Eds.), *Bergey's Manual of Systematic Bacteriology*, The Proteobacteria. Part B, The Gamma proteobacteria, 2nd edn., vol. 2. Springer, New York, pp. 494–546.
- Hill, V.R., N. Cohen, and A.M. Kahler. (2011). Toxigenic *Vibrio cholerae* O1 in water and seafood, Haiti. *Emerging Infectious Disease*, **17**: 2147–2150.
- Kim, Y.B., Okuda, J., Matsumoto, C., Takahashi, N., Hashimoto, S. and Nishibuchi, M. (1999) Identification of *Vibrio parahaemolyticus* strains at the species level by PCR targeted to the *toxR* gene. *Journal of Clinical Microbiology*, **37**: 1173–1177.
- Kumar, R and K. V. Lalitha. (2013). Prevalence and molecular characterization of *Vibrio cholerae* O1, Non-O1 and Non-O139 in Tropical Seafood in Cochin, India. *Foodborne Pathogens Disease*, **10**(3):278-283.
- Oliver, J.D. (2006). *Vibrio vulnificus*. In: *The Biology of Vibrios*. Thompson, F.L., B. Austin, and J. Swings. (eds), ASM Press, Washington, DC, pp. 349–366.
- Osorio, C.R. and Klose, K.E. (2000) A region of the transmembrane regulatory protein *ToxR* that tethers the transcriptional activation domain to the cytoplasmic membrane displays wide divergence among *Vibrio* species. *Journal of Bacteriology*, **182**: 526–528.
- Senderovich, Y., Izhaki, I. and Halpern, M. (2010). Fish as reservoir and vectors of *Vibrio cholerae*. *PLoS. One*, **5**: 8607.

- Sudha, S., Mridula, C., Silvester, R. and Hatha, A.A.M. (2014). Prevalence and antibiotic resistance of pathogenic Vibrios in shellfishes from Cochin market. *Indian Journal of Geo-Marine Science*, **43**(5): 815-824.
- Xu, X., J. Cheng, Q. Wu, J Zhang and T. Xie. (2016). Prevalence, characterization, and antibiotic susceptibility of *Vibrio parahaemolyticus* isolated from retail aquatic products in North China. *BMC. Microbiology*, **16**: 32.
- Yu, Q., M. Niu, M. Yu, Y. Liu, D. Wang and X. Shi. (2016). Prevalence and antimicrobial susceptibility of *Vibrio parahaemolyticus* isolated from retail shellfish in Shanghai. *Food Control*, **60**: 263-268.