

A1 and A2 milk

– What science claims?

Ayushi Chaudhary¹, Harjeet Singh², Gyanendra S. Senger³ and Umesh Singh⁴

ICAR-Central Institute for Research on Cattle, Meerut 250 001, Uttar Pradesh

Indigenous breeds of cows and buffaloes are of A2 milk type and hence are a source for safe milk. The A1/A2 status of cattle is determined by the two alleles for beta-casein gene i.e. A1 and A2 cattle may be homozygous for one type of protein (A1A1 or A2A2), or heterozygous (A1A2). A1A1 cattle would produce only A1 milk, A2A2 would produce only A2 milk and A1A2 cattle would produce milk with both A1 and A2 beta - caseins. NBAGR also reported moderate to high frequency of A2 allele among the breeding bulls, further supporting the belief that milk being sold in India is safe for human consumption. Considering the lack of concrete scientific evidence for A1/BCM-7 ill-effects on human health, and predominance of A2 milk in Indian cattle there is no need to move to A2 only.

Key words: A1 milk, A2 milk, BCM7, Milk, Protein

MILK is enormously evolved secretion of mammary glands produced by mammals being perfect food for novice. Globally, India is the highest milk producer (176.4 million tonnes) as well as consumer of milk, neither exports nor imports milk. Since long back milk is considered as complete source of nutrients (both micro and macronutrients) providing a range of proteins, carbohydrates and other important components of human food.

Around the World, cow, buffalo, sheep, camel and goat are known as the main producers of milk out of which cow being the major producer of milk i.e. 83% (about 600 million tonnes) of the total milk produced annually. Also, India is the leading producer of milk (176.3 million tonnes approximately) which is uppermost in the world. In relation to India alone, 46.9% milk is yielded by buffalo while 49% of milk is produced by cattle.

The major fraction of milk contains water which constitute about 85% of milk while remaining 15% of milk is composed of sugars

such as lactose (4.6%), milk proteins (2.8%), fats (3.7%), minerals (3.36%) and other component like hormones, immunoglobulins, growth factors, cytokines, nucleotides, peptides, polyamines and enzymes.

Major polymorphs of cows' milk

Milk protein is divided into two major classes i.e. casein and whey protein. The casein may be further sub-divided as alpha, beta and kappa

casein, out of which beta-casein is more ubiquitous containing balanced proportions of essential amino acids. The overall protein makeup of milk is 36% α-Casein, 27% β-Casein, 9% γ-casein whereas other peptides and amino acids comprise about 27% of total milk proteins. Out of all casein proteins in milk β-Casein are of foremost importance as it is the reason for generation of various genetic variants of milk. So far about



Fig. 1. Depiction of BCM 7 (Betacasomorphins-7) breakdown at 67th position of peptide due histidine amino acid in A1 beta-casein variant.

12 genetic variants are known namely A1, A2, A3, B, C, D, E, F, H1, H2, I and G out of which 'A1' and 'A2' are often found. 'C' is the rarest of rare and not so often found in milk proteins. β -Casein of cows' milk contains 209 amino acids. A fluctuation of histidine (A1 type) and proline (A2 type) at position 67 is profoundly observed which give rise to two polymorph of same protein of β -Casein *viz.* variants A1 and A2.

A1 Milk: A valid risk?

β -Casein of cow milk is made up of 209 amino acids. It accounts 24-28% of total milk protein and its molecular weight is about 24 kD. It is position 67 where A1 and A2 variants of cows' milk differ from each other. While A1 contains histidine at this position, A2 have proline instead of histidine. Beta casein releases a range of opioids known as beta casomorphins (BCMs). β -casomorphin-7 (BCM-7), a seven amino acids long bioactive peptide, released in small intestine during digestion of A1 β -Casein which have been found associated as a factor of risk while human health is concerned.

Health problems associated to A1 β -Casein variant

A number of workers had worked upon the problems related to A1 variant of β -Casein of cow milk. It is believed that β -casomorphin-7 (BCM-7), an opioid, have ability to oxidize low dietary lipoproteins which could lead to the formation of arterial plaques. Digestive enzymes lack ability to break A2 protein due to the presence of proline instead of histidine.

Also some of the studies carried out in various European countries such as United Kingdom, Iceland, Norway, Austria as well as in American subcontinent revealed a strong association between β -casein A1 and diabetes mellitus (DM1).

There are three classes of receptors for opioid in brain *viz.* μ (MOP), κ (KOP) and δ (DOP) (IUPHAR, 2008). These receptors are distributed in immune cells, endocrine glands, brain and in intestine. Bovine milk has opioid effect and MOP are the

Table 1. Some of the major health related issues related to A1 β -Casein milk

Health Problem(s)	Target Organ(s)
Type I diabetes	Pancreas
CVD (Cardiovascular disease)	Heart
Arteriosclerosis	Blood vessels and Heart
Schizophrenia	Brain
Autism	Brain
SIDS (Sudden Infant Death Syndrome)	Unexplained
Autistic Spectral Disorder	Brain
Digestive Health	Digestive system
Lactose intolerance	
Magnesium Deficiency and Imbalance	Electrolyte imbalance
Endometriosis	Uterus

receptors through which they interact mostly. It was also found that BCM-7 of bovine milk has stronger affinity as compared to human milk for MOP receptors. Genes producing inflammatory enzymes like myeloperoxidases are found enhanced by BCM-7 that could lead to the expression of MOP receptors in intestine. Though, no report has been encountered related to human brain.

Methodologies available for evaluation of beta-casomorphins in milk

Number of workers in recent past have worked or been working on to identify and quantify the presence of betacasomorphins in milk. Some of the methodologies which are in use are summarized in Table 2.

Aforementioned Table illustrated various methods which are more or less in use to identify or quantify betacasomorphine in milk proteins. Apart from these techniques there are

Table 2. Evaluation of beta-casomorphins in milk

Method/Technique	Theory
Reversed-phase high performance liquid chromatography (RP-HPLC)	It is exercised for separation of peptides and amino acids
RPHPLC- UV	Employed for the separation and quantification of BCM7 present in cheese
Ion-exchange chromatography	Employed for the separation and quantification of BCM7 present in cheese
HPLC-UV	Has been used for determination of BCM7 amount in human milk
Tandem massspectrometry	It helps precise quantification of BCM5 and BCM7 at minor levels
Quadrupole iontrapmass spectrometry (QIT-MS)	Measure BCM5 and BCM7 in single reaction.

several methods which are in use for genotyping various polymorphs of milk proteins in cattle such as PCR-SSR (simple sequence repeat polymorphism or microsatellite markers), PCR-SSCP (Single Strand Conformation Polymorphism), Restriction fragment length polymorphism (PCR-RFLP analysis), Single nucleotide polymorphism (SNP) assay followed by direct sequencing, Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). All these methods are PCR-based methods which can be used to discriminate different variants of milk proteins. Also, there are some techniques *viz.* isoelectric focussing and RT-HPLC which are not dependent of PCR. Some of the major advantages and disadvantages of these methods are summarized in Table 3.

Significance of β -Casein milk variants concerning India

All across the world A1 and A2 alleles of β -Casein gene distributed variably in different cattle breeds. There is a serious debate going on whether A1 milk is not suitable for human health when compared to A2 milk. Some researchers were in support of the opinion that consumption of A1 type milk may be hazardous for human health. Also, some other reports were encountered which recommends that A2 milk is better for human health and crossbreeding should be avoided for overcoming serious consequences on human health. Exotic or crossbreed cattle variants have elevated in number from 14.4 million to 19.42 million

Table 3. Advantage and disadvantage of some method in evaluating milk proteins

Method	Advantage	Disadvantage
IEF	High resolution, resulting in greater separation of solutes Easier, because the placement of sample application is not important The sample and ampholytes can be mixed before application	Carrier ampholyte batch-to-batch reproducibility of the ampholytes Phenomenon of cathodic drift owing to electroendosmosis in the gel causes an eventual breakdown of the basic end of the pH gradient with time
SDS-PAGE	Highly reproducible Inexpensive, fast, sensitive, expressive Economic (no sophisticated equipment required) Easy (clear, one-dimensional separation principle)	Limited pI range (4-8) Proteins >150 kD not seen in 2D gels Difficult to see membrane proteins (>30% of all proteins) Only detects high abundance proteins (top 30% typically) Time consuming
RT-HPLC	Speed (minutes) High resolution Sensitivity (ng to fg) Reproducibility of +/- 1% Accuracy Automation	Cost Complexity Low sensitivity for some compounds Irreversibly adsorbed compounds not detected Coelution difficult to detect
PCR-SSR	Require very little and not necessarily high quality DNA Highly polymorphic Evenly distributed throughout the genome Simple interpretation of results Easily automated, allowing multiplexing Good analytical resolution and high reproducibility	Complex discovery procedure Expensive

(approximately 34.78 %), which is a serious concern as per their opinion.


Genotype project

Studies by the National Bureau of Animal Genetic Resources (NBAGR), Karnal covering 22 *desi* breeds have established that predominant genotype in India's native cattle is A2A2, confirming that our indigenous cows and buffaloes produce A2 milk.

The frequency of A2 allele was 100 per cent in the five high-yielding milk breeds — Red Sindhi, Gir, Rathi, Sahiwal and Tharparkar, meaning that these breeds do not have A1 allele or A1A1/A1A2 genotype. In the remaining breeds, the availability of A2 allele was 94 per cent.

¹Young Professional II CG&B Division, ²SRF, ³Research Associate, ⁴Principal Scientist, Corresponding author's e-mail: ayushichaudhary9502@gmail.com

Textbook on Fundamentals of Rural Sociology and Educational Psychology



India being a land of villages, rural sociology and educational psychology is an essential theme for agricultural students and professionals. Rural sociology deals with rural society and the relations of people who live in villages. Rural sociology presents a scientific picture of rural life.

Extension workers work with farmers very closely in their settings for which their understanding of rural background and farmers' psychology is very essential. Extension professionals should have necessary knowledge of the precise approaches and methods of dealing with farmers. Rural sociology can help professionals in organizing the rural structure in a constructive manner.

This textbook consists of fundamental facts of rural sociology. The book has twenty two chapters covering both rural sociology and educational psychology out of which twelve chapters consist of rural sociology and ten chapters deal with educational psychology. In this text book, emphasis has been given on various aspects of educational psychology also. It provides all the relevant information in the field of rural sociology with special emphasis on Indian culture, merits and demerits of Indian rural societies, the stressful conditions under which farmers live and work. The book will be a good basis of information reference for agriculture students, psychology students besides extension practitioners, research scientists, KVKs and agricultural colleges.

TECHNICAL SPECIFICATIONS

No. of pages : i-viii + 252 • Price : ₹ 450 • ISBN No. : 978-81-7164-181-9

For obtaining copies, please contact:

Business Unit

Directorate of Knowledge Management in Agriculture
Krishi Anusandhan Bhavan-I, Pusa, New Delhi 110 012
Tel : 011-25843657, Fax 91-11-25841282; e-mail : bmicar@gmail.com