

Multiple dimensions of agroecology in sustaining agriculture

A K Srivastava* and Popy Bora

ICAR-Indian Agricultural Research Institute, Gogamukh, Assam 787 035

Sustaining agriculture has become a more daunting challenge amidst multiple nutrient deficiencies triggered by depleting carbon stock of soil coupled with glaring impact of global warming on crop performance. Despite paradigm shift from purely inorganic to either organic or the combination of two, agroecological applications have always occupied a center stage in sustainable agriculture. Development of microbial consortium (microbial concoction), rhizosphere hybridization, identification of microbial corridor and microbial hot spot in a field, besides storing carbon in plant vegetative framework, are some of the upcoming ways to realize the multiple dimensions of agroecology in sustaining the agriculture in years to come with emphasis on on-farm research. These efforts have to be further matched with other natural resource conserving options like conservation agriculture, organic farming and natural farming.

Keywords: Agroecology, Microbial diversity, Natural farming, Sustainable agriculture

A GROECOLOGY works best when natural resources are utilized with on-farm ambit as a driver of sustainability in agriculture, where very often natural farming is linked to give birth to another form of agriculture, conceived as regenerative agriculture having much stronger resilience against climate change-related issues. Agroecological farming like any other sustainable agriculture, integrates five major issues, viz. soil health care, soil biodiversity, crop biodiversity, production stability and environmental health and water quality. Of these, soil health is the central core of sustainable agriculture. Of late, sustainability of agriculture is hugely challenged by global warming triggering unwarranted climate change, the impact of which becomes more discernible, considering strong synergism between CO₂ and nutrients under no water limiting conditions. Several studies revealed that under central Indian conditions recommended dose of fertilizers (RDF) once worked, stands out to be ineffective over a period of 20-years time on account of rise in average temperature by just less than 1°C during flower opening time (such examples can likewise be cited for any other crop), which later necessitated 25% additional K to moderate such temperature stress. How does RDF behave in the long run in different crops when agroecology is integrated with emphasis on microbial turnover of the nutrients to regulate soil fertility functions. Recognition of the importance of soil microorganisms has led to a renewed interest in measuring the quantum of nutrients

held in their biomass and their consequential effect on crop responses on a sustainable basis.

Amidst these facts, a still bigger question emerges, how effectively we can utilise the microbial niches of crop phytobiome to unlock the ongoing plateau in crop production. In this article we will discuss the possible contribution of agroecology beyond crop production sustainability. The current G20 summit under the lead role of India looks forward towards environmentally benign possibilities of sustaining, not only the crop production, but looks far beyond via environmental health for the safeguard of posterity.

Agroecological implications: A regenerative approach

Land degradation (96.4 million ha of degraded land accounting to 29.3% of the country's total geographical area of 328.7 million ha) neutrality has been one of the most vibrant strategies of national agriculture policy where coalition of conventional and traditional farming takes place with singular objectivity of sustainability through agroecology driven regenerative agriculture. India made some modest contribution to realise the strength of sustainable agriculture through: national project on organic farming; systematic rice intensification and zero budget natural farming, all put together under one umbrella towards *Atmanirbhar Bharat*. But, the basics of modern agriculture are never overlooked with regards to core issues like natural resource conservation, soil microbial diversity, resilience

against forging climate change, expanding water intake capacity, scavenging soil contaminants, usage of cover crops (field buffers and plant strips on contours) for reduced run-off loss and maintaining the environmental health. While modern agriculture takes into account the rejuvenation of depleted land from physical, biological and chemical barriers, restricting the targeted optimised crop agronomy and aid further in recuperating the full potential for crop carrying capacity in a farming system module. Hence, agroecology based regenerative agriculture recognizes all sustainable practices those affect the natural systems and uses all the management techniques to restore the system towards improved crop productivity.

While comparing regenerative agriculture with organic agriculture, both are well connected with agroecology, we comprehensively overlook the harmful effects of organic pesticides, which could be even more harmful than synthetic pesticides in organically produced crops. Are natural pesticides safer than artificial pesticides?, the answer could be both in both the ways. Surely, such scientific outcomes put an alarm bell to researchers and policy makers to keep a regular guard on health of agro-ecosystem using agroecology as a complete toolkit.

Intervention of agroecology in conventional agriculture imminent

Concurrent emergence of multiple nutrient deficiencies is one of the prime reasons for fast depleting soil organic carbon stock. No doubt, there has been concerted efforts of measuring carbon pool of Indian soils, including the carbon stock under different land uses. These efforts are more of indicative than suggestive, especially on mitigation terms, with still miniscule information available on horticultural crops based land uses. Lack of concepts like crop grids and land uses -based efforts have put researchers in dilemma to define the upper limits of soil organic carbon as a guarantee to sustained crop production, despite the fact that soil carbon content is the single most important soil property having cascading effects on entire range of soil properties. Our efforts to improve soil organic carbon using various forms of composts and manures have proved futile using citrus as a case study, despite an initial display of increase in organic carbon in the

first –5 years, but later started declining, regardless of fruit yield level. These studies also lend strong support that unless carbon is deposited into passive carbon pool, mere improvement in active pool of carbon will not aid in developing a climate resilient crop production system, unless agroecological intervention is brought into the system. Therefore, an introduction of concept like microbial consortium brought some respite to infuse production sustainability.


Exploiting microbial niche for soil-plant health bioprospecting

Plant phytobiome displaying microbial diversity (culture dependent or culture independent) offer the best opportunity to develop and upscale the combination of rhizo-competent microbes, popularly called as microbial consortium or microbial concoction. The most common objective of developing microbial consortium is to capitalize on both the capabilities of individual microbes and their interactions to create useful systems in tune with enhanced productivity, and soil health improvements through efficient metabolic functionality.

We carried out studies with an aim to develop rhizosphere specific microbial consortium. The efficient microbes, viz. *Aspergillus flavus* (MF113270, P-solubilizer), *Bacillus pseudomyoides* (MF113272, K-solubilizer), *Acinetobacter radioresistens* (MF113273, N-solubilizer), *Micrococcus yunnanensis* (MF113274, P-solubilizer) and *Paenibacillus alvei* (MF113275, P-solubilizer) were finally identified and later tested under nursery as well as field conditions. In the experiment on progressive response of multiple microbes of the microbial consortium was tested without addition of any inorganic fertilizers through soil inoculation, different microbes for the purpose of retrofitting microbes as nutrient replacement. Multiple location testing revealed a significant reduction in mortality of citrus seedlings and budlings. While long term field evaluation (2007–16) of microbial consortium in different combinations with limited chemical fertilizers and vermicompost (Table 1) further showed 30–40% saving on chemical fertilizers without any compromise on fruit yield, fruit quality and soil health including microbial biomass and microbial biomass nutrients, with an additional reduction of carbon dioxide (concurrent increase carbon fixation), indicating recalcitrance nature of carbon lodged into passive pool of soil organic carbon (called legacy carbon). Such recalcitrance of microbes mediated deposited carbon brings crop nutrient holiday more realistic in application than mere speculation.

Rhizosphere hybridization: A rejuvenation for microbial diversity

Of the many novelties, emergence of concept like rhizosphere hybridization (though concept has to still find a scientific acceptance) has provided a strong silver lining as to ensure rhizosphere microbial diversity, a prerequisite for production sustainability. Rhizosphere hybridization is a new concept to modify the rhizosphere ecology to create an optimum environment for harnessing the value added benefits of nutrient-microbe synergy. Our studies on response of different treatments involving rhizosphere soil of three perennial trees, viz. *Ficus racemosa* L. (Umber tree),



Changes in the organic carbon during 8 years of treatment with different organic manures in comparison to organic fertilizers


T₁, Farmyard manure; T₂, Vermicompost; T₃, Poultry manure; T₄, Green manure; T₅, Neem cake; T₆, inorganic fertilizers
Source: Srivastava (2012)

Table 1. Fruit yield and quality parameter of citrus crop in response to different combinations of microbial consortium and vermicompost as varied modules of integrated soil fertility management (Pooled data: 2007–16)

ISFM module	Yield (kg/tree)	Fruit quality parameters			
		Juice content (%)	TSS (°Brix)	Acidity (%)	TSS/Acid ratio
Module I	67.7	40.2	8.4	0.96	8.92
Module II	71.0	39.6	8.6	0.93	9.23
Module III	80.5	41.5	9.5	0.86	10.05
Module IV	83.2	42.7	8.4	0.91	9.56
Module V	88.8	44.2	9.3	0.80	11.62
CD (P=0.05)	2.1	2.1	0.30	0.10	1.10

Module I, 100% RDF; Module II, 75% RDF + 25% Vermicompost; Module III, 75% RDF + 25% Vermicompost + Microbial consortium; Module IV, 50% RDF + 50% Vermicompost; Module V, 50% RDF + 50% Vermicompost + Microbial consortium.

Source: Srivastava *et al.* (2022)



Changes in CO₂ emission rate (mg C/m²/hr) in response to different ISFM modules (Module I-Module V), representing the agroecological consequences in developing carbon neutral agriculture


Source: Srivastava (2022)

Ficus benghalensis L. (Banyan tree) and *Ficus religiosa* L. (Pipal tree) along with rhizosphere soil of healthy and highly productive sweet orange trees in sweet orange buddlings showed differential response in terms of agronomic parameters, changes in soil physical properties, and pool of plant available nutrients. In field, the rhizosphere hybridization can be implemented by collecting rhizosphere soil of healthy trees (along microbial corridor) and injected into weaker trees to rationalize distribution of microbes across field/orchard as a part of agroecological farming.

Agroecology and plant carbon stock

Field crops or horticultural crops are key to any land usage, contributing significantly towards carbon dioxide sinks for tree's permanent woody framework. Storing carbon in plants vegetative framework is a promising alternative to conventional method of carbon sequestration in soil. This is where agroecological applications have great role to play. For example, fruit crops are estimated to sequester 24–109 tons CO₂/ha (depending upon their age, planting distance, soil types, agrotechniques etc.), far higher than many of the tropical rain forests, with an additional ecosystem service in economic terms. Incidentally, the response of different fruit crops under elevated CO₂ condition is a function of nutrition status of the crop, where soil microbial ecology plays a pivotal role. Although, the carbon stocked into the trees is subjected to stiff challenge against long term storage and released into the atmosphere through respiration of soil micro-communities.

The carbon stored into the perennial framework of fruit



Agronomic response of consortium of PGPAs, antagonists, endophytes and nano-composites for improvement in plant nutrition and reduction in HLB soil borne pathogens

Source: A case study of agroecological application, Srivastava *et al.* (2022)

crops is petrified with numerous organic compounds, and then subsequently distributed into various parts of the plants to discharge multiple functions, including their implications on computable economic outcome. While comparing the carbon storage into the soil, which is relatively more unstable, variable, and vulnerable to decomposition, compared to carbon stored into the woody fabric of fruit crops. Therefore, we need a shift of concept, despite the fact that researchers across disciplines are concerned about storing carbon into the soil for an extended period. In either care, recalcitrance of carbon and its residence time would be of paramount importance to exploit atmospheric CO₂. Such a concept is anticipated to be successful with minor modifications in photosynthesis system of field crops and intervention of microbes, especially endophytic in nature to store carbon inside the plants tissues. Considering endophytic microbes as beneficial bioagents and potent colonizers of plant's tissues, we might be able to use the world's tiniest organisms to solve the world's most pressing problem.

SUMMARY

A cultivar evaluated under both intensive farming, organic farming or natural farming system may not perform with similar magnitude of success. The major difference lies with respect to differential soil health indices, which are yet to be streamlined. While talking about Soil Health Card, do we need to breed the fruit crops specifically tailored to such forms of farming, an enigmatic answer still awaits for researchers to either refute such hypothesis or accept with sound scientific database. Hence, linking agroecology with sustainable agriculture is a forward looking agriculture model with more emphasis on genetic, functional and metabolic diversity of soil microorganisms within crop-based rhizosphere. The capacity of soil microbial communities to maintain functional diversity of critical soil processes could ultimately be more important to ecosystem productivity and stability than mere taxonomic diversity.

*Corresponding author's e-mail: aksrivast2007@gmail.com