



Correlation and path analysis studies in some mutant and non-mutant lines of Indian mustard (*Brassica juncea* L. Czern. & Coss.) for seed yield and related traits

Salam Alisha Devi¹, PK Barua¹ and Aradhana Phukan^{2*}

¹Department of Plant Breeding and Genetics, Assam Agricultural University, Jorhat, 785013, Assam

²College of Horticulture and Farming System Research, Nalbari, Assam Agricultural University, 781338, Assam

*Corresponding author: aradhana.phukan@aau.ac.in

(Received: 28 August 2024; Revised: 25 September 2024; Accepted: 23 October 2024)

<https://doi.org/10.56093/JOB.v16i1.17>

Abstract

Twenty-nine genotypes of Indian mustard comprising of twenty-one $M_{4.5}$ mutants of the parent varieties TM-2 and PM-27, five newly developed $F_{6.7}$ lines and three check varieties (TM-2, NRCHB-101 and PM-27) were evaluated in a randomized block design during *rabi* 2019-20 and *rabi* 2020-21 to assess the correlation and path coefficients for seed yield and related traits. The genotypes were evaluated for fourteen quantitative characters. Seed yield per plant was positively correlated with number of siliquae on main shoot at the genotypic level. Number of siliquae on main shoot was positively correlated with plant height, main shoot length, siliqua density and 1000-seed weight. Path analysis based on genotypic correlations showed positive direct effects of number of siliquae on main shoot, number of primary branches and secondary branches per plant and seed yield per plant.

Keywords: Indian mustard, correlation, seed yield, path coefficient analysis

Introduction

Oilseeds play an important role in the human diet and known for their application in industrial use. India is the fifth largest vegetable oil economy in the world, produces 7.4 % oilseeds, 5.8% oils, 6.1% oil meal and 9.3% of edible oils (Jat *et al.*, 2019). Of the total oilseed production, *Brassica* accounts for 24% of the oilseeds acreage and occupies second rank after soybean. Indian mustard ($2n=36$) is a Rabi oilseed crop grown during October to February-March under diverse agro-climatic conditions. It is used for its oil, oil meal and as green leafy vegetables. According to the DRMR, Bharatpur, in the global scenario, rapeseed-mustard crops are cultivated in 35.95 million hectares with a production of 71.49 million tonnes and productivity of 1990 kg/ha as in 2019-20. India holds the position in second after Canada in acreage (19.81%) and fourth in production (10.37%) behind Canada, European Union and China (Jat *et al.*, 2019).

In India, rapeseed-mustard crops occupy 6.86 million hectares, produce 9.12 million tonnes of oilseeds with average productivity of 1331 kg/ha as in 2019-20. The state of Assam is among the six major rapeseed-mustard crop producing states in the country. Assam contributes about 4.5 per cent of area and 2.05 percent of production of total rapeseed-mustard in India. Rapeseed-mustard is grown in substantial area in Assam with a productivity of

647 kg/ha (DRMR, 2022). In India, growing of rapeseed-mustard serves as an important source of income for small and marginal farmers. However, owing to the low availability of edible oils and their continuous import in the country, it is important to increase the yield and production of these crops. As mustard is a long duration crop, toria is more popular in Assam but produces lesser yield than mustard. Therefore, it is important to develop short-duration superior Indian mustard varieties and to increase the average oilseed production in the state. However, yield is a complex polygenic trait and therefore it is important to analyse the relationship between yield and component characters. The association between the yield contributing characters can be determined with the help of correlation analysis. Path analysis helps to identify the direct and indirect effects of different components on seed yield which is required in the selection of high yielding genotypes. With this view, the present investigation was conducted based on correlation and path analysis to study the inter-relationships between the characters.

Materials and Methods

During *rabi* 2019-20 and 2020-21, twenty-one M_4 mutant lines (developed by gamma irradiation of the parent lines TM-2 and PM-27), five newly developed F_6 lines (2019-20) and three check varieties (TM-2, PM-27 and NRCHB-101) of Indian mustard (Table 1) were used for this

experiment. The materials were collected from Department of Plant Breeding and Genetics, AAU, Jorhat. The experimental farm was located at the Department of Plant Breeding & Genetics, Assam Agricultural University, Jorhat, Assam at a latitude of 26° 45' North and a longitude

of 94° 12' East with an elevation of 87 m above mean sea level. The layout was followed in randomized block design with three replications. The agronomic practices were followed according to the standard package and practices of the crop during the cultivation.

Table 1: List of genotypes used in the study

S. No.	Line No.	Designation	S. No.	Line No.	Designation
Mutant lines of TM-2			Mutant lines of TM-2		
1	15	JMM-TM2-15	18	10	JMM-TM2-10
2	17	JMM-TM2-17	19	2	JMM-TM2-2
3	38	JMM-TM2-38	Mutant lines of PM-27		
4	34	JMM-TM2-34	20	1	JMM-PM27-1
5	20	JMM-TM2-20	21	11	JMM-PM27-11
6	47	JMM-TM2-47	Newly-developed lines		
7	160	JMM-TM2-160	22		JM13-2
8	1	JMM-TM2-1	23		JM13-3
9	4	JMM-TM2-4	24		JM13-4
10	51	JMM-TM2-51	25		JM13-5
11	22	JMM-TM2-22	26		JM13-6
12	73	JMM-TM2-73	Check varieties		
13	208	JMM-TM2-208	27		TM-2
14	28	JMM-TM2-28	28		NRCHB-101
15	14	JMM-TM2-14	29		PM-27
16	23	JMM-TM2-23			
17	27	JMM-TM2-27			

Morpho-phenological observations

Fourteen quantitative characters *viz.*, days to 50% flowering (DF), days to maturity (DM), plant height in cm (PH), number of primary branches (PBN) and secondary branches per plant (SBN), main shoot length in cm (MSL), number of siliquae on main shoot (SMS), siliqua density on main shoot in no./cm (SD), number of seeds per siliqua (SPS), number of flowers on terminal main shoot (TF), number of siliqua on terminal main shoot (TS), 1000-seed weight in g (TSW), oil content in % (OC) and seed yield per plant in g (SYP) were recorded according to the standard procedure.

The genotypic and phenotypic correlation coefficients between two characters were calculated by analysis of covariance following the standard procedures by Dabholkar (1999). Path coefficient analysis was carried out using the genotypic correlation coefficients to by following Dewey and Lu (1959) and Dabholkar (1999). The statistical methods were computed using Microsoft Excel 2007.

Results and Discussion

As yield is a complex quantitative trait, it is influenced by various yield attributing characters both at genotypic and phenotypic levels. Correlation studies determine the degree

and direction of association between different yield component traits (Mondal and Khajuria, 2009; Shalini *et al.*, 2000). The genotypic correlations between the characters were studied year-wise, which were presented in Table 2 (2019-20) and Table 3 (2020-21). In the first year, SYP was positively correlated with PH, SMS, SD, 1000-seed weight and PBN at the genotypic level. Swetha *et al.* (2019) and Singh *et al.* (2013) reported positive correlation of seed yield with PH, SMS and 1000-seed weight. Similarly, positive correlation of seed yield with PBN were reported by Nandi *et al.* (2021), Ray *et al.* (2019), Tantuway *et al.* (2018) and Kumari & Kumari (2018).

In the second year, positive correlation was observed between SYP, SPS, TF, TS and SMS. These findings were similar to the results reported by Pandey *et al.* (2020), Tiwari *et al.* (2018) and Kumar *et al.* (2018) for SPS; by Swetha *et al.* (2019), Yadav and Pandey (2018) and Singh *et al.* (2013) for SMS. However, SYP was negatively correlated with DF, which indicates early flowering genotypes might also be higher yielder. Similar results were observed by Saiyad *et al.* (2020) and Kumar *et al.* (2016). Considering both the years, it was observed that SYP was positively correlated with SMS and negatively correlated with DF at the genotypic level. Similarly, TS

Table 2: Genotypic (above diagonal) and phenotypic (below diagonal) correlation coefficients between different characters in Indian mustard during 2019-20

DF	DM	PH	PBN	SBN	MSL	SMS	SD	SPS	TF	TS	TSW	OC	SYP
DF	1	0.820**	0.604**	-0.108	0.053	0.270	0.340	0.383*	0.021	-0.141	0.805**	0.176	0.199
DM	0.165	1	0.713**	-0.324	0.465*	0.670**	0.436*	-0.267	0.257	-0.029	0.157	0.711**	0.241
PH	0.233	0.469*	1	0.181	0.680**	0.806**	0.356	-0.186	0.360	0.093	0.510**	0.337	0.562**
PBN	-0.042	-0.012	0.157	1	0.463*	0.318	0.121	-0.109	0.292	0.215	-0.037	-0.475**	0.418*
SBN	-0.059	-0.100	-0.047	0.508**	1	0.260	-0.162	-0.282	0.192	0.485**	-0.289	-0.693**	-0.079
MSL	0.054	0.397*	0.587**	0.206	0.624**	1	0.750**	-0.261	0.347	0.139	0.407	-0.027	0.105
SMS	0.118	0.412*	0.711**	0.296	0.624**	1	0.548**	-0.335	-0.002	0.408*	0.463*	0.118	0.525**
SD	0.083	0.080	0.218	0.157	-0.333	0.520**	1	-0.145	0.135	0.461*	0.179	0.230	0.574**
SPS	0.138	-0.167	-0.108	-0.058	-0.184	-0.213	-0.047	1	-0.126	-0.296	0.116	0.214	-0.266
TF	0.129	0.303	0.287	0.166	0.189	0.069	0.219	-0.075	1	0.328	-0.260	-0.110	-0.200
TS	0.021	-0.063	0.084	0.204	0.017	0.261	0.290	-0.111	0.291	1	0.191	-0.539**	0.183
TSW	0.185	-0.006	0.363	-0.034	0.260	0.348	0.129	0.066	-0.087	0.279	1	0.132	0.450*
OC	0.135	0.257	0.220	-0.311	-0.007	0.030	0.030	0.109	-0.114	-0.380*	0.037	1	0.291
SYP	0.164	0.263	0.440*	0.260	0.176	0.451*	0.332	-0.065	0.039	0.049	0.170	0.162	1

* Significant at 5 % probability, ** Significant at 1% probability

Table 3: Genotypic (upper diagonal) and phenotypic (lower diagonal) correlation coefficients between different characters in Indian mustard during 2020-21

DF	DM	PH	PBN	SBN	MSL	SMS	SD	SPS	TF	TS	TSW	OC	SYP
DF	1	0.243	0.072	-0.047	-0.030	-0.219	0.273	-0.339	-0.186	-0.343	-0.167	0.402*	-0.399*
DM	0.338	1	0.100	-0.146	0.035	0.186	0.219	-0.208	-0.268	-0.221	-0.151	-0.017	-0.278
PH	0.078	0.036	1	0.230	0.276	0.743**	0.284	0.075	-0.051	0.041	0.265	-0.083	0.263
PBN	-0.112	-0.128	0.238	1	0.114	0.286	0.121	-0.291	0.006	0.153	0.358	-0.301	-0.012
SBN	0.009	0.015	0.283	0.141	1	0.147	-0.006	-0.007	0.397	0.288	-0.123	-0.194	0.236
MSL	-0.225	-0.138	0.430*	0.160	0.179	1	-0.482**	0.314	-0.002	0.101	0.342	0.160	0.366
SMS	-0.018	0.089	0.648**	0.318	0.205	0.471**	0.500**	0.029	-0.002	0.277	0.441*	-0.218	0.367*
SD	0.241	0.223	0.164	0.157	0.003	-0.538**	1	-0.302	0.296	0.232	0.096	-0.451*	0.030
SPS	-0.124	-0.082	0.069	-0.122	0.137	0.156	-0.058	1	0.414*	0.303	0.339	0.043	0.499**
TF	-0.091	-0.132	0.014	0.036	0.305	0.069	0.221	0.250	1	1.020	0.225	-0.403*	0.431*
TS	-0.227	-0.085	0.031	0.051	0.220	0.198	0.017	0.138	0.801**	1	0.157	-0.432*	0.425*
TSW	-0.102	-0.053	0.199	0.260	-0.065	0.394*	0.120	0.195	0.260	0.110	1	0.069	0.225
OC	0.195	0.068	-0.054	-0.260	-0.099	-0.125	-0.141	0.000	-0.254	-0.302	0.062	1	0.022
SYP	-0.384*	-0.238	0.277	0.108	0.240	0.309	0.041	0.414*	0.315	0.228	0.134	-0.089	1

* Significant at 5 % probability, ** Significant at 1% probability

Table 4: Direct and indirect effects of different component characters on seed yield per plant in Indian mustard during 2019-20 (based on genotypic correlation)

	DM	PH	PBN	SBN	MSL	SMS	SPS	TS	TSW	Genotypic correlation with SYP
DM	-0.482	0.463	-0.006	0.026	-0.453	0.538	0.097	0.009	0.048	0.241
PH	-0.344	0.649	0.068	0.008	-0.663	0.647	0.067	-0.027	0.157	0.562**
PBN	0.008	0.117	0.377	-0.038	-0.267	0.255	0.040	-0.063	-0.011	0.418*
SBN	0.156	-0.061	0.174	-0.082	-0.254	0.114	0.102	-0.141	-0.089	-0.079
MSL	-0.224	-0.041	0.103	-0.021	-0.975	0.602	0.095	-0.040	0.125	0.105
SMS	-0.323	0.523	0.120	-0.012	-0.731	0.803	0.122	-0.119	0.142	0.525**
SPS	0.129	-0.121	-0.041	0.023	0.254	-0.269	-0.363	0.086	0.036	-0.266
TS	0.014	0.060	0.081	-0.040	-0.136	0.328	0.107	-0.291	0.059	0.183
TSW	-0.076	0.331	-0.014	0.024	-0.397	0.372	-0.042	-0.056	0.307	0.450*

Residual = 0.295

Direct effect (path coefficient) in bold face on the leading diagonal

Table 5: Direct and indirect effects of different component characters on seed yield per plant in Indian mustard during 2020-21 (based on genotypic correlation)

	DM	PH	PBN	SBN	MSL	SMS	SPS	TS	TSW	Genotypic correlation with SYP
DM	-0.308	-0.017	0.016	0.006	-0.001	0.104	-0.083	-0.015	0.020	-0.278
PH	-0.031	-0.173	-0.026	0.048	0.034	0.413	0.030	0.003	-0.035	0.263
PBN	0.045	-0.040	-0.112	0.020	0.022	0.207	-0.117	0.011	-0.047	-0.012
SBN	-0.011	-0.048	-0.013	0.173	0.011	0.090	-0.003	0.020	0.016	0.236
MSL	0.002	-0.078	-0.032	0.025	0.075	0.285	0.126	0.007	-0.045	0.366
SMS	-0.057	-0.129	-0.042	0.028	0.039	0.556	0.012	0.019	-0.058	0.367*
SPS	0.064	-0.013	0.033	-0.001	0.024	0.016	0.401	0.021	-0.045	0.499**
TS	0.068	-0.007	-0.017	0.050	0.008	0.154	0.121	0.069	-0.021	0.425*
TSW	0.046	-0.046	-0.040	-0.021	0.026	0.245	0.136	0.011	-0.132	0.225

Residual effect = 0.697

Direct effect (path coefficient) in bold face on the leading diagonal

was positively correlated with PH and 1000-seed weight at genotypic level, while it was correlated with MSL and SD at both phenotypic and genotypic levels in both the years. PH showed high

positive phenotypic correlation with MSL and SMS while DF showed negative correlation with SYP. Kumar *et al.*, (2018) also reported negative phenotypic correlation between SYP and DF.

Path coefficient analysis is used to partition the observed correlation coefficient between yield and component traits into direct and indirect effects (Wright, 1921). It helps in identifying the component traits in selection criteria for yield improvement. The path analysis was carried out year-wise as presented in Table 4 and Table 5. Path analysis based on genotypic correlations in both the years revealed that SMS and PBN were important yield attributing traits contributing

highest positive direct effects on yield. Considering both the years, DM showed highest negative direct effect towards SYP. SBN also contributed positively towards seed yield to some extent. In a study conducted by Tripathi *et al.* (2019), PBN was observed to have positive direct effect while DF showed negative direct effect towards SYP. Correlation and path analysis indicated that the number of SMS, PBN and SBN were the major yield attributing characters. Considering these traits, selection would be effective for enhancing seed yield in Indian mustard.

Authors' contribution

Conceptualization of research (PKB, SAD); Designing of the experiments (PKB, SAD); Contribution of experimental materials (PKB, SAD); Execution of field/lab experiments and data collection (SAD); Analysis of data and interpretation (PKB, SAD); Preparation of manuscript (SAD, PKB, AP).

Declaration

The authors declare no conflict of interest.

Acknowledgement

The author is highly grateful to Assam Agricultural University, Jorhat for funding the research work under PG programme.

References

- Dabholkar AR. 1999. Elements of Biometrical Genetics. Concept Publishing Company, New Delhi.
- DACFW. 2020. Agricultural Statistics at a Glance (2020), Department of Agriculture, Cooperation and Farmers Welfare, Directorate of Economics and Statistics, Ministry of Agriculture and Farmers Welfare, Government of India. (<https://eands.dacnet.nic.in/>)
- Devi B. 2018. Correlation and path analysis in Indian mustard (*Brassica juncea* L.) in agro-climatic condition of Jhansi (UP). *J Pharmac Phytochem*, **7**: 1678-1681.
- Dewey DR and Lu KH. 1959. A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron J*, **51**: 515-518.
- DRMR 2022. Technical extension folder, ICAR-DRMR-OPIU (Agri.)-APART Project. Directorate of Rapeseed-Mustard Research, Bharatpur, Rajasthan.
- Jat L, Rai SK, Choudhary JR, Bawa V, Bharti R, Sharma M. and Sharma M. 2019. Phenotypic evaluation of genetic diversity of diverse Indian mustard (*Brassica juncea* L. Czern and Coss) genotypes using correlation and path analysis. *Int J Bio-resource Stress Manag*, **10**: 467-471.
- Kumar A, Singh M, Yadav RK, Singh P and Lallu. 2018. Study of correlation and path coefficient among the characters of Indian mustard. *The Pharma Inno J*, **7**: 412-416.
- Kumar R, Gaurav SS, Jayasudha S and Kumar H. 2016. Study of correlation and path coefficient analysis in germplasm lines of Indian mustard (*Brassica juncea* L.). *Agric Sci Digest-A Res J*, **36**: 92-96.
- Kumari A and Kumari V. 2018. Exploitable genetic variability and character association studies under changing climate in mid-hills of north-west Himalayas in Indian mustard [*Brassica juncea* (L.) Czern. & Coss.]. *J Pharmac Phytochem*, **7**: 372-379.
- Mondal SK and Khajuria MR. 2000. Genetic analysis of yield attributes in mustard. *Environ Ecology*, **18**: 1-5.
- Nandi S, Rout S, Sur B, Chakraborty M, Hijam L, Roy SK and Gupta A. 2021. Assessment of genetic parameters, trait association and diversity analysis in Indian mustard [*Brassica juncea* (L.) Czern and Coss]. *Biological Forum – An Int J*, **13**: 95-109.
- Pandey SK, Srivastava KK, Negi S, Khan NA and Singh RK. 2020. Variability, trait relationship and path analysis for seed yield and seed quality parameters in Indian mustard (*Brassica juncea* L.). *J Oilseed Brassica*, **11**: 69-76.
- Ray J, Singh OP, Verma SP, Pathak VN, Singh B and Jee C. 2019. Characters association studies for yield contributing traits in Indian mustard (*Brassica juncea*). *Environ and Ecology*, **37**: 1497-1500.
- Saiyad AT, Patel PJ, Patel JR, Prajapati KP and Patel BK. 2020. Study of correlation and path coefficient among the quantitative and qualitative characters of Indian mustard [*Brassica juncea* (L.) Czern & Coss]. *Int J Chemical Stud*, **8**: 2112-2116.
- Shalini TS, Sheriff RA, Kulkarni RS and Venkataravana P. 2000. Genetic divergence in Indian mustard (*Brassica juncea* L. Czern and Coss.). *Mysore J Agri Sci*, **34**: 251-256.
- Singh A, Avtar R, Singh D, Sangwan O, Thakral NK, Malik VS, Goyat B and Dalal U. 2013. Combining ability analysis for seed yield and component traits in Indian mustard [*Brassica juncea* (L.) Czern & Coss.]. *Res Plant Biol*, **3**: 26-31.
- Swetha M, Janeja HS, Singh H, Sravani M, Rajaneesh K and Madakemohekar AH. 2019. Genetic evaluation of Indian mustard (*Brassica juncea* L.) genotypes for yield and quality parameters. *Plant Archives* **19**: 413-417.
- Tantuway G, Srivastava K and Tirkey AE. 2018. Character association and path analysis studies of yield and yield-attributing traits in Indian mustard (*Brassica juncea* L. Czern & Coss). *Plant Archives*, **18**: 2245-2251.
- Tiwari AK, Singh SK, Tomar A and Singh M. 2017. Heritability, genetic advance and correlation coefficient analysis in Indian mustard (*Brassica juncea* (L.) Czern & Coss). *J Pharmac Phytochem*, **6**: 356-359.
- Tripathi A, Singh M and Tomar A. 2019. Selection parameter analysis in Indian mustard (*Brassica juncea*) germplasm under normal sown condition. *J Pharmac Phytochem*, **8**: 1551-1553.
- Wright S. 1921. Systems of mating I. The biometric relations between parent and offspring. *Genetics*, **6**: 111-123.
- Yadav S and Pandey A. 2018. Inter relationship and path analysis for seed yield in Indian mustard [*Brassica juncea* (L.) Czern & Coss]. *The Pharma Inno J*, **7**: 152-156.