

Ethanol resistance, as a way to evaluate the stability of leaf cell membranes of cynodon (*Cynodon dactylon*) and 4 wheat genotypes, significantly different in drought resistance and leaf shape

Petar Petrov and Ivan Dimitrov

IFRG-BAS, Bulgaria

Article history:

Received: 17 Aug., 2023

Revised: 19 Nov., 2023

Accepted: 13 Dec., 2023

Citation:

P Petar and I Dimitrov. 2024. Ethanol resistance, as a way to evaluate the stability of leaf cell membranes of cynodon (*Cynodon dactylon*) and 4 wheat genotypes, significantly different in drought resistance and leaf shape. *Journal of Cereal Research* **16** (1): 53-59. <http://doi.org/10.25174/2582-2675/2024/149757>

*Corresponding author:

E-mail: pivanovp@abv.bg

© Society for Advancement of Wheat and Barley Research

Abstract

The stability of leaf cell membranes is one of the factors on which the resistance of agricultural crops to the abiotic factors of the environment depends. Ethanol, as a very good solvent of lipids and highly damaging to cell membranes, was used to test the membrane stability of leaf cells of cynodon (*Cynodon dactylon*) and four genotypes of wheat. The membranes of the cynodon leaf cells are the most resistant. A concentration of 15% ethanol was chosen as the most suitable for testing the wheat genotypes. Of the wheat genotypes, the leaf cells of Katya and Enola have the greatest membrane stability and small leaf dissection index (LDI). The rates of damage to leaf cell membranes are constant in the cynodon and the Katya genotype, while in the other genotypes they increase exponentially. Genotype of the modern selection Prelom has a small LDI and the most unstable membranes.

Keywords: wheat genotypes, drought resistance, membrane stability, ethanol resistance, leaf injury index, leaf shape

1. Introduction

Climatic changes require the selection of wheat genotypes resistant to the abiotic factors of the environment. The stability of cell membranes of leaf cells is key to the tolerance of wheat genotypes to abiotic stresses. Electrolyte leakage method provides an assessment of the stability of cell membranes, i.e. the greater the electrolyte leakage, the more damaged the cell membranes. This method helps in the selection of more cold tolerant (Yazdi-Samadi et al., 2006), more heat tolerant (Dias et al., 2010) and more drought tolerant (Bajji, Mohammed et al., 2002; Ahmadizadeh et al., 2011; Petrov et al., 2018) wheat genotypes. The method is also used to assess salinity tolerance of wheat (Basra et al., 2005).

The organization of cell membranes is very complex. Lipids are a major component of membranes. Ingólfsson et al., 2014 proposed a model of 63 different types of lipids. The lipid bilayer environment surrounding membrane proteins strongly influences their structure and

functions (Tero et al., 2017). The dynamically structured mosaic model proposed by Vereb et al. (2003) suggested a dynamic rearrangement of lipids and proteins through Brownian motion. According to Bhat et al. (2005) biological membranes are not a simple homogeneous layer of proteins and lipids, but rather are organized into discrete regions that can be characterized by different lipid and protein contents. These membrane microdomains are called "lipid rafts". Rafts in membranes are defined by the physical properties of the lipid bilayer and function by selectively partitioning membrane lipids and proteins into membrane domains with specific phase behavior and lipid packing (Harder, 2003).

Lipid solvents could damage the structure of cell membranes and disrupt their functioning. Ethanol, as a very good lipid solvent, has a toxic effect on membranes. Studies of the resistance of cell membranes to ethanol have been carried out in animal and human cells. The fluidity of cell



membranes is important for their functioning and stability. One of the effects that ethanol induces on membranes is to alter their fluidity by altering the fluidity of membrane lipids. Alcohol treatment induces changes in rat synaptic membrane fluidity (Zerouga et al., 1992). Goral et al. (2008) suggested that ethanol may have dose-dependent effects on cell membrane fluidity in human cells. No studies are known to assess the tolerance of plant cell membranes to ethanol. Testing the membrane stability of agricultural crop cells with ethanol solutions, by conductometry with a simplified protocol, is performed in a short time as an express test applicable in breeding practice.

2. Materials and Methods

2.1 Plant material and growing conditions

The cynodon leaves were plucked from plants growing under natural conditions. Leaves of wheat genotypes were plucked from plants grown for 21 days under controlled conditions. The four Bulgarian wheat (*Triticum aestivum* L.) varieties were used, of which 1 old historic varieties and 3 modern releases. The old variety Slomer represent relatively homogeneous selection made within landraces, or early breeding releases. This germplasm is not cultivated anymore and is nearly extinct. Seed samples have been kindly provided by Dr. Andreas Börner, German Federal Genebank, Leibniz Institute of Plant Genetics. Seeds were soaked for 4 h in tap water and planted in 1-kg pots with alluvial meadow soil (pH 6.2). Plants were grown in a climatic chamber with 22/18 °C day/night temperature, 14-h photoperiod, irradiance of 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 70 % relative humidity. Tap water was supplied daily sustaining 60 % of full soil moisture capacity.

2.2 Leaf injury index (LII) and electrolyte leakage kinetics

For determination of LII 15 leaf pieces (2 cm in length) were cut from plants. The solutions was measured with a conductometer Elwro 5721 (Poland). Finally, samples were boiled for 30 min, cooled at room temperature and conductivity was read again. LII was measured by a modification of the Premachandra formula (Premachandra et al., 1992). The original formula is: $I (\%) = [1 - (1 - t_1/t_2)/(1 - c_1/c_2)] \cdot 100$ where t_1 and t_2 are the first and second (after boiling) conductivity measurements of the solutions in which they are treated samples were immersed and c_1 and c_2 are the corresponding values of the controls. In

our case, $t_2 = c_2$, since leaves from untreated plants were used for the experiments. For c_1 , the conductivity values of ethanol solutions with distilled water, which are close to the values of distilled water, are taken.

2.3 Leaf dissection index

Leaf dissection index (LDI) as a function of leaf perimeter and area was calculated by Fourier transformation (Kincaid and Schneider, 1983): $DI = \text{perimeter} / [2\sqrt{(\text{area} \times \pi)}]$

2.4 Statistical analysis

Two independent experiments were conducted and parameters were measured in at least 3 replications. Data are presented as mean values $\pm$ SE

3. Results

3.1 LII of the cynodon (*Cynodon dactylon*) and of the wheat genotype with low drought tolerance (Slomer) treated with a series of ethanol concentrations.

At concentrations of 5 and 10% of the cynodon leaves, no significant electrolyte leakage was recorded until the end of the experiment (fig. 1). At the higher concentrations (15 and 20 %) from the 60th to the 240th minute, a gradual increase in electrolyte leakage up to about 15% was reported (fig. 1).




Fig. 1. Leaf injury index of cynodon (*Cynodon dactylon*) treated with ethanol solutions

After the 120th minute, the lowest concentration of ethanol that causes an increase in the electrolyte leakage from the membranes of the leaf cells of the Slomer wheat genotype is 5% (fig. 2). At the higher concentrations of ethanol, the electrolyte leakage increased from the 30th minute and reached about 20% at the 240th minute at all three concentrations (fig. 2).






Fig. 2. Leaf injury index of the leaves of the Slomer wheat genotype treated with ethanol solutions

3.2 LII of 4 wheat genotypes treated with 15% ethanol.

By the 90th minute, electrolyte leakage increased to the same extent in all four genotypes. After the 120th hour, electrolyte leakage has higher values in the Slomer and Prelom genotypes (fig. 3). At the end of the experiment, the

electrolyte leakage was the highest in the Prelom genotype (fig. 3). The electrolyte leakage of the Slomer genotype has a lower value (fig. 3). The electrolyte leakage is the lowest in the Katya and Enola genotypes (fig. 3).




Fig. 3. Leaf injury index of 4 wheat genotypes treated with 15% ethanol solutions

3.3 Rates of damage to leaf cell membranes

The rates of damage to the cell membranes of cynodon leaves treated with 15 and 20% ethanol solutions were constant (fig. 4).

The rates of damage to the leaf cell membranes of the Enola, Slomer and Prelom genotypes treated in 15% increased exponentially (fig. 4). In the Katya genotype, the rates of leaf cell membrane damage are again constant (fig. 4).






Fig. 4. Rates of damage to cell membranes of leaves of 4 wheat genotypes treated with 15% ethanol solution and cell membranes of cynodon (*Cynodon dactylon*) leaves treated with 15 and 20% ethanol solutions

3.4 Leaf perimeter and LDI

The Slomer genotype has the largest leaf perimeter and LDI (tab. 1).

Table 1.

Genotypes	Leaf perimeter (cm)	Leaf area (cm ²)	Leaf dissection index
Katya	35.1 ^a ± 0.90	7.34 ^a ± 0.37	7.3 ^a ± 0.15
Enola	36.8 ^a ± 0.85	7.09 ^a ± 0.17	7.8 ^a ± 0.19
Slomer	41.2 ^b ± 0.80	7.13 ^a ± 0.33	8.7 ^b ± 0.15
Prelom	36.1 ^a ± 0.91	7.2 ^a ± 0.2	7.6 ^a ± 0.17

4. Discussion

4.1 LII of cynodon after treatment with ethanol

The minimum concentration of ethanol at which significant damage to the cynodon cell membranes is observed is 15% (fig. 1), while it is 10% for the Slomer wheat genotype (fig. 2). As a rule, wild plants are more viable than cultivated ones, which was also observed in our research. These results confirm the workability of our approach.

4.2 Optimization of ethanol concentration for LII testing of wheat genotypes

For the tests, a concentration of ethanol was chosen that was not too low (causing no or little damage to the membranes) or too high (causing strong and equal

damage to the membranes of all genotypes). Slomer is a wheat genotype of old Bulgarian selection. Our previous research showed that the genotype has a low degree of drought resistance and labile membranes of leaf cells (Petrov et al., 2018). Slommer leaves were treated with a series of ethanol concentrations in order to select an appropriate concentration to compare the LII of wheat genotypes (fig. 2). At the end of the experiment, at all three concentrations (10, 15 and 20%), the damage to the leaf cell membranes was significant (about 20% - fig. 2). The lowest concentration at which there is significant damage to cell membranes is 10% (fig. 2). Taking into account the sensitivity of the Slomer genotype and the lack of response of the cynodon to an alcohol concentration of 10% (fig. 1), the next strongest concentration - 15% - was



chosen to compare the membrane stability of the wheat genotypes. An approach similar to ours (via ethanol solutions) has been used to study the ethanol resistance of brewer's yeast (Jones et al., 1987). According to research by Pratt et al. (2003) 10% ethanol inhibited yeast growth and 20% also inhibited their fermentation capacity. The concentrations of ethanol that affect the life processes of the yeast (between 10 and 20%) are same to the test concentration we used (15%).

4.3 LII of 4 wheat genotypes treated with 15% ethanol solution

Differences in electrolyte leakage between genotypes were clearly evident at 210 minutes and persisted at 240 minutes (fig. 3). The maintenance of differences between genotypes in the last two measurements indicates that a treatment period of 4 hours is sufficient to obtain stable results. The average level of electrolyte leakage at the end of the experiment (about 20 %) indicates a strong but not complete damage to the membranes of the leaf cells and allows the achievement of contrasting differences between the genotypes in terms of their membrane stability (fig. 3). Genotype Slomer is an old selection, while genotypes Enola, Prelom and Katya are new selection wheat.

The Katya genotype known for its drought resistance (Vassileva et al., 2011; Vassileva et al., 2009; Petrov et al., 2018; Doneva et al., 2021; Kocheva et al., 2013) is expected to have stable membranes. Our studies confirm the membrane resistance of this genotype, i.e. the degree of damage to its membranes after exposure to ethanol is weak (fig. 3). Unexpectedly, ethanol damaged the leaf cell membranes of the new selection genotype Prelom to a significantly greater extent than the leaf cell membranes of the drought-sensitive genotype old selection Slomer (Fig. 3).

4.4 Rates of damage to leaf cell membranes after treatment with ethanol

The exponential increase in the rate of leaf cell membrane damage in the Prelom, Slomer and Enola genotypes indicates that ethanol causes a time-progressive destruction of the leaf cell membranes largely beyond the control of the cellular mechanisms for maintaining membrane stability (Fig. 4). The constant rate of electrolyte leakage from the leaf cells of the cynodon after exposure to ethanol (with concentrations

of 15 and 20% - fig. 4) leads to the assumption that the cellular mechanisms for maintaining membrane stability work significantly more efficiently in it. The constant rates of electrolyte leakage after exposure to ethanol and in the Katya genotype suggest that cellular mechanisms maintain the stability of cell membranes in it with an efficiency close to that of the cynodon.

4.5 The role of leaf shape and membrane stability in drought

Our previous research showed that under severe drought conditions, the lower the leaf RWC, the greater the membrane damage (Petrov et al., 2018). Severe drought caused in leaves with a more elongated shape (higher LDI) a faster loss of water, up to wateriness values at which the membranes are severely damaged. In this case, the damage to the membranes directly depends on the leaf shape and, respectively, on the water loss. In the case of moderate stress, the leaves do not dehydrate strongly and the tolerance of the plants also depends on the characteristics of the membranes of the respective genotype (Peršić et al., 2022). Our research shows that the genotype of the modern selection Prelom has a small LDI (table 1) and at the same time the most unstable membranes.

Conclusion

LII of cynodon and 4 wheat genotypes was evaluated by conductometric study with 4 hourly experiments. 15% ethanol was chosen as the test concentration. The leaf cells of the cynodon have the most resistant membranes. Among the wheat genotypes with the highest resistance are the membranes of the leaf cells of the genotypes with small LDI - Katya and Enola. Our research shows that the genotype of the modern selection Prelom has a small LDI and at the same time the most unstable membranes.

The approach we used allows with a simple protocol and in a short time to successfully compare the membrane resistance of the leaf cells of the wheat genotypes.

Author contributions

All authors contributed equally for preparing the final version of the manuscript.

Conflict of Interest

Authors declare no conflict of interest.



Ethical Approval

The article doesn't contain any study involving ethical approval.

Acknowledgments

I thank Prof. Svetlana Misheva for kindly providing wheat genotypes

Thanks for the technical assistance of Mrs. A. Petrova

References

1. Ahmadizadeh M, M Valizadeh, M Zaefizadeh and H. Shahbazi. 2011. Antioxidative protection and electrolyte leakage in durum wheat under drought stress condition. *Journal of Applied Sciences Research*, 7(3), pp.236-246.
2. Bajji M, JM Kine and S Lutts. 2002. The use of the electrolyte leakage method for assessing cell membrane stability as a water stress tolerance test in durum wheat. *Plant growth regulation*, 36, pp.61-70. <https://link.springer.com/article/10.1023/A:1014732714549>
3. Basra SMA, I Afzal, RA Rashid and A Hameed. 2005. Inducing salt tolerance in wheat by seed vigor enhancement techniques. *International Journal of Agriculture and Biology*, 2(1), pp.173-179.
4. Bhat RA and R Panstruga. 2005. Lipid rafts in plants. *Planta*, 223, pp.5-19. <https://link.springer.com/article/10.1007/s00425-005-0096-9>
5. Dias AS, MG Barreiro, PS Campos, JC Ramalho and FC Lidon. 2010. Wheat cellular membrane thermotolerance under heat stress. *Journal of Agronomy and Crop Science*, 196(2), pp.100-108. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1439-037X.2009.00398.x>
6. Doneva D, M Pál, L Brankova, G Szalai, J Tajti, R Khalil and B Ivanovska. 2021. The effects of putrescine pre-treatment on osmotic stress responses in drought-tolerant and drought-sensitive wheat seedlings. *Physiologia Plantarum*, 171(2), pp.200-216. <https://onlinelibrary.wiley.com/doi/10.1111/ppl.13150>
7. Goral J, J Karavitis and EJ Kovacs. 2008. Exposure-dependent effects of ethanol on the innate immune system. *Alcohol*, 42(4), pp.237-247. <https://www.sciencedirect.com/science/article/abs/pii/S0741832908000414>
8. Harder T. 2003. Formation of functional cell membrane domains: The interplay of lipid- and protein-mediated interactions. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 358(1433), pp.863-868. <https://royalsocietypublishing.org/doi/10.1098/rstb.2003.1274>
9. Ingólfsson HI, MN Melo, FJ Van Eerden, C Arnarez, CA Lopez, TA Wassenaar, X Periole, AH De Vries, P Tieleman and SJ Marrink. 2014. Lipid organization of the plasma membrane. *Journal of the american chemical society*, 136(41), pp.14554-14559. <https://pubs.acs.org/doi/full/10.1021/ja507832e>
10. Jones RP and PF Greenfield. 1987. Ethanol and the fluidity of the yeast plasma membrane. *Yeast*, 3(4), pp.223-232. <https://doi.org/10.1002/yea.320030403>
11. Kincaid DT and RB Schneider. 1983. Quantification of leaf shape with a microcomputer and Fourier transform. *Canadian Journal of Botany*, 61(9), pp.2333-2342. <https://doi.org/10.1139/b83-256>
12. Kocheva KV, PI Petrov and GI Georgiev. 2013. Physiological and anatomical responses of wheat to induced dehydration and rehydration. *Central European Journal of Biology*, 8, pp.499-503. <https://doi.org/10.2478/s11535-013-0153-x>
13. Peršić V, A Ament, J Antunović Dunić, G Drezner, and V Cesar. 2022. PEG-induced physiological drought for screening winter wheat genotypes sensitivity-integrated biochemical and chlorophyll a fluorescence analysis. *Frontiers in Plant Science*, 13, pp.987702. <https://doi.org/10.3389/fpls.2022.987702>
14. Petrov P, A Petrova, I Dimitrov, T Tashev, K Olsovska, M Brestic and S Misheva. 2018. Relationships between leaf morpho-anatomy, water status and cell membrane stability in leaves of wheat seedlings subjected to severe soil drought. *Journal of Agronomy and Crop Science*, 204(3), pp.219-227. <https://doi.org/10.1111/jac.12255>
15. Pratt PL, JH Bryce and GG Stewart. 2003. The effects of osmotic pressure and ethanol on



- yeast viability and morphology. *Journal of the Institute of Brewing*, 109(3), pp.218-228. <https://doi.org/10.1002/j.2050-0416.2003.tb00162.x>
16. Premachandra GS, H Saneoka, K Fujita and S Ogata. 1992. Leaf water relations, osmotic adjustment, cell membrane stability, epicuticular wax load and growth as affected by increasing water deficits in sorghum. *J Exp Bot*, 43(12), pp.1569–1576. <https://doi.org/10.1093/jxb/43.12.1569>
17. Tero R, K Fukumoto, T Motegi, M Yoshida, M Niwano and A Hirano-Iwata. 2017. Formation of cell membrane component domains in artificial lipid bilayer. *Scientific reports*, 7, pp.1-10. http://space.bas.bg/SES/archive/SES%202023_DOKLADI/2_Aerospace%20Technologies/5_Petrov.pdf
18. Vassileva V, C Signarbieux, I Anders and U Feller. 2011. Genotypic variation in drought stress response and subsequent recovery of wheat (*Triticum aestivum* L.). *Journal of Plant Research*, 124, pp.147-154. <https://link.springer.com/article/10.1007/s10265-010-0340-7>
19. Vassileva V, L Simova-Stoilova, K Demirevska and U Feller. 2009. Variety-specific response of wheat (*Triticum aestivum* L.) leaf mitochondria to drought stress. *Journal of plant research*, 122, pp.445-454. <https://link.springer.com/article/10.1007/s10265-009-0225-9>
20. Vereb G, J Szöllősi, J Matko, P Nagy, T Farkas, L. Vigh, L Matyus, TA Waldmann and S Damjanovich. 2003. Dynamic, yet structured: the cell membrane three decades after the Singer–Nicolson model. *Proceedings of the national academy of sciences*, 100, pp.8053-8058. <https://doi.org/10.1073/pnas.1332550100>
21. Yazdi-Samadi B, H Dashti, A Nabipour and A Sarrafi. 2006. Reciprocal substitutions analysis of freezing resistance in wheat (*Triticum aestivum* L.). *Plant breeding*, 125(3), pp.217-220. <https://doi.org/10.1111/j.1439-0523.2006.01234.x>
22. Zerouga M and F Beaugé. 1992. Rat synaptic membrane fluidity parameters after intermittent exposures to ethanol in vivo. *Alcohol*, 9(4), pp.311-315. [https://doi.org/10.1016/0741-8329\(92\)90072-I](https://doi.org/10.1016/0741-8329(92)90072-I)

